


A Solution Manual For

**Differential equations and the calculus of
variations by L. ELSGOLTS. MIR
PUBLISHERS, MOSCOW, Third
printing 1977.**



Nasser M. Abbasi

May 15, 2024

Contents

1	Chapter 1, First-Order Differential Equations. Problems page 88	2
2	Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172	509
3	Chapter 3, SYSTEMS OF DIFFERENTIAL EQUATIONS. Problems page 209	892

1 Chapter 1, First-Order Differential Equations.

Problems page 88

1.1	problem Problem 1	4
1.2	problem Problem 2	16
1.3	problem Problem 3	28
1.4	problem Problem 4	36
1.5	problem Problem 5	50
1.6	problem Problem 6	61
1.7	problem Problem 7	74
1.8	problem Problem 8	87
1.9	problem Problem 9	100
1.10	problem Problem 10	113
1.11	problem Problem 11	126
1.12	problem Problem 12	130
1.13	problem Problem 13	134
1.14	problem Problem 14	145
1.15	problem Problem 15	148
1.16	problem Problem 16	165
1.17	problem Problem 17	170
1.18	problem Problem 18	182
1.19	problem Problem 26	185
1.20	problem Problem 28	189
1.21	problem Problem 29	200
1.22	problem Problem 30	216
1.23	problem Problem 31	221
1.24	problem Problem 35	225
1.25	problem Problem 36	229
1.26	problem Problem 37	237
1.27	problem Problem 39	241
1.28	problem Problem 40	246
1.29	problem Problem 42	251
1.30	problem Problem 43	259
1.31	problem Problem 45	277
1.32	problem Problem 46	292
1.33	problem Problem 47	312
1.34	problem Problem 48	317
1.35	problem Problem 49	322
1.36	problem Problem 50	330

1.37	problem Problem 51	343
1.38	problem Problem 52	355
1.39	problem Problem 53	366
1.40	problem Problem 54	370
1.41	problem Problem 55	377
1.42	problem Problem 56	385
1.43	problem Problem 57	403
1.44	problem Problem 58	418
1.45	problem Problem 59	435
1.46	problem Problem 60	447
1.47	problem Problem 61	455
1.48	problem Problem 62	466
1.49	problem Problem 63	480
1.50	problem Problem 64	494
1.51	problem Problem 65	498
1.52	problem Problem 66	504

1.1 problem Problem 1

1.1.1	Solving as separable ode	4
1.1.2	Solving as first order ode lie symmetry lookup ode	6
1.1.3	Solving as exact ode	10
1.1.4	Maple step by step solution	14

Internal problem ID [12112]

Internal file name [OUTPUT/10764_Monday_September_11_2023_12_50_28_AM_43300585/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 1.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact", "separable", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_separable]`

$$\tan(y) - y' \cot(x) = 0$$

1.1.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{\tan(y)}{\cot(x)}\end{aligned}$$

Where $f(x) = \frac{1}{\cot(x)}$ and $g(y) = \tan(y)$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\tan(y)} dy &= \frac{1}{\cot(x)} dx \\ \int \frac{1}{\tan(y)} dy &= \int \frac{1}{\cot(x)} dx \\ \ln(\sin(y)) &= -\ln(\cos(x)) + c_1\end{aligned}$$

Raising both side to exponential gives

$$\sin(y) = e^{-\ln(\cos(x))+c_1}$$

Which simplifies to

$$\sin(y) = \frac{c_2}{\cos(x)}$$

Summary

The solution(s) found are the following

$$y = \arcsin\left(\frac{c_2 e^{c_1}}{\cos(x)}\right) \quad (1)$$




Figure 1: Slope field plot

Verification of solutions

$$y = \arcsin\left(\frac{c_2 e^{c_1}}{\cos(x)}\right)$$

Verified OK.

1.1.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{\tan(y)}{\cot(x)}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 1: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= \cot(x) \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{\cot(x)} dx\end{aligned}$$

Which results in

$$S = -\ln(\cos(x))$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{\tan(y)}{\cot(x)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 0 \\R_y &= 1 \\S_x &= \tan(x) \\S_y &= 0\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \cot(y) \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \cot(R)$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \ln(\sin(R)) + c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\ln(\cos(x)) = \ln(\sin(y)) + c_1$$



Which simplifies to

$$-\ln(\cos(x)) = \ln(\sin(y)) + c_1$$

Which gives

$$y = \arcsin\left(\frac{e^{-c_1}}{\cos(x)}\right)$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{\tan(y)}{\cot(x)}$ 	$R = y$ $S = -\ln(\cos(x))$	$\frac{dS}{dR} = \cot(R)$ 

Summary

The solution(s) found are the following

$$y = \arcsin\left(\frac{e^{-c_1}}{\cos(x)}\right) \tag{1}$$




Figure 2: Slope field plot

Verification of solutions

$$y = \arcsin\left(\frac{e^{-c_1}}{\cos(x)}\right)$$

Verified OK.

1.1.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}\left(\frac{1}{\tan(y)}\right) dy &= \left(\frac{1}{\cot(x)}\right) dx \\ \left(-\frac{1}{\cot(x)}\right) dx + \left(\frac{1}{\tan(y)}\right) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -\frac{1}{\cot(x)} \\ N(x, y) &= \frac{1}{\tan(y)}\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{\cot(x)}\right) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{1}{\tan(y)} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{\cot(x)} dx \\ \phi &= \ln(\cos(x)) + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{\tan(y)}$. Therefore equation (4) becomes

$$\frac{1}{\tan(y)} = 0 + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$\begin{aligned}f'(y) &= \frac{1}{\tan(y)} \\ &= \cot(y)\end{aligned}$$

Integrating the above w.r.t y results in

$$\int f'(y) dy = \int (\cot(y)) dy$$

$$f(y) = \ln(\sin(y)) + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \ln(\cos(x)) + \ln(\sin(y)) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \ln(\cos(x)) + \ln(\sin(y))$$

Summary

The solution(s) found are the following

$$\ln(\cos(x)) + \ln(\sin(y)) = c_1 \tag{1}$$




Figure 3: Slope field plot

Verification of solutions

$$\ln(\cos(x)) + \ln(\sin(y)) = c_1$$

Verified OK.

1.1.4 Maple step by step solution

Let's solve

$$\tan(y) - y' \cot(x) = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'}{\tan(y)} = \frac{1}{\cot(x)}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\tan(y)} dx = \int \frac{1}{\cot(x)} dx + c_1$$

- Evaluate integral

$$\ln(\sin(y)) = -\ln(\cos(x)) + c_1$$

- Solve for y

$$y = \arcsin\left(\frac{e^{c_1}}{\cos(x)}\right)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.203 (sec). Leaf size: 9

```
dsolve(tan(y(x))-cot(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \arcsin(\sec(x) c_1)$$

✓ Solution by Mathematica

Time used: 4.745 (sec). Leaf size: 19

```
DSolve[Tan[y[x]]-Cot[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \arcsin\left(\frac{1}{2}c_1 \sec(x)\right)$$

$$y(x) \rightarrow 0$$

1.2 problem Problem 2

- 1.2.1 Solving as homogeneousTypeMapleC ode 16
- 1.2.2 Solving as first order ode lie symmetry calculated ode 20

Internal problem ID [12113]

Internal file name [OUTPUT/10765_Monday_September_11_2023_12_50_29_AM_54692774/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 2.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"homogeneousTypeMapleC", "first_order_ode_lie_symmetry_calculated"**

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$6y + (5x + 2y - 3)y' = -12x + 9$$

1.2.1 Solving as homogeneousTypeMapleC ode

Let $Y = y + y_0$ and $X = x + x_0$ then the above is transformed to new ode in $Y(X)$

$$\frac{d}{dX}Y(X) = -\frac{3(4X + 4x_0 + 2Y(X) + 2y_0 - 3)}{5X + 5x_0 + 2Y(X) + 2y_0 - 3}$$

Solving for possible values of x_0 and y_0 which makes the above ode a homogeneous ode results in

$$x_0 = 0$$

$$y_0 = \frac{3}{2}$$

Using these values now it is possible to easily solve for $Y(X)$. The above ode now becomes

$$\frac{d}{dX}Y(X) = -\frac{3(4X + 2Y(X))}{5X + 2Y(X)}$$

In canonical form, the ODE is

$$\begin{aligned} Y' &= F(X, Y) \\ &= -\frac{6(2X + Y)}{5X + 2Y} \end{aligned} \quad (1)$$

An ode of the form $Y' = \frac{M(X, Y)}{N(X, Y)}$ is called homogeneous if the functions $M(X, Y)$ and $N(X, Y)$ are both homogeneous functions and of the same order. Recall that a function $f(X, Y)$ is homogeneous of order n if

$$f(t^n X, t^n Y) = t^n f(X, Y)$$

In this case, it can be seen that both $M = -12X - 6Y$ and $N = 5X + 2Y$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{Y}{X}$, or $Y = uX$. Hence

$$\frac{dY}{dX} = \frac{du}{dX}X + u$$

Applying the transformation $Y = uX$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dX}X + u &= \frac{-6u - 12}{2u + 5} \\ \frac{du}{dX} &= \frac{\frac{-6u(X)-12}{2u(X)+5} - u(X)}{X} \end{aligned}$$

Or

$$\frac{d}{dX}u(X) - \frac{\frac{-6u(X)-12}{2u(X)+5} - u(X)}{X} = 0$$

Or

$$2\left(\frac{d}{dX}u(X)\right)Xu(X) + 5\left(\frac{d}{dX}u(X)\right)X + 2u(X)^2 + 11u(X) + 12 = 0$$

Or

$$12 + X(2u(X) + 5)\left(\frac{d}{dX}u(X)\right) + 2u(X)^2 + 11u(X) = 0$$

Which is now solved as separable in $u(X)$. Which is now solved in $u(X)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(X, u) \\ &= f(X)g(u) \\ &= -\frac{2u^2 + 11u + 12}{X(2u + 5)} \end{aligned}$$

Where $f(X) = -\frac{1}{X}$ and $g(u) = \frac{2u^2+11u+12}{2u+5}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{2u^2+11u+12}{2u+5}} du &= -\frac{1}{X} dX \\ \int \frac{1}{\frac{2u^2+11u+12}{2u+5}} du &= \int -\frac{1}{X} dX \\ \frac{2 \ln(2u+3)}{5} + \frac{3 \ln(u+4)}{5} &= -\ln(X) + c_2\end{aligned}$$

The above can be written as

$$\begin{aligned}\frac{2 \ln(2u+3) + 3 \ln(u+4)}{5} &= -\ln(X) + c_2 \\ 2 \ln(2u+3) + 3 \ln(u+4) &= (5)(-\ln(X) + c_2) \\ &= -5 \ln(X) + 5c_2\end{aligned}$$

Raising both side to exponential gives

$$e^{2 \ln(2u+3)+3 \ln(u+4)} = e^{-5 \ln(X)+5c_2}$$

Which simplifies to

$$\begin{aligned}(2u+3)^2 (u+4)^3 &= \frac{5c_2}{X^5} \\ &= \frac{c_3}{X^5}\end{aligned}$$

Which simplifies to

$$u(X) = \text{RootOf} \left(4_Z^5 + 60_Z^4 + 345_Z^3 - \frac{c_3 e^{5c_2}}{X^5} + 940_Z^2 + 1200_Z + 576 \right)$$

Now u in the above solution is replaced back by Y using $u = \frac{Y}{X}$ which results in the solution

$$Y(X) = X \text{RootOf} (4_Z^5 X^5 + 60_Z^4 X^5 + 345_Z^3 X^5 + 940_Z^2 X^5 - c_3 e^{5c_2} + 1200_Z X^5 + 576 X^5)$$

Using the solution for $Y(X)$

$$Y(X) = X \text{RootOf} (4_Z^5 X^5 + 60_Z^4 X^5 + 345_Z^3 X^5 + 940_Z^2 X^5 - c_3 e^{5c_2} + 1200_Z X^5 + 576 X^5)$$

And replacing back terms in the above solution using

$$Y = y + y_0$$

$$X = x + x_0$$

Or

$$Y = y + \frac{3}{2}$$

$$X = x$$

Then the solution in y becomes

$$y - \frac{3}{2} = x \text{RootOf}(4_Z^5 x^5 + 60_Z^4 x^5 + 345x^5_Z^3 + 940x^5_Z^2 - c_3 e^{5c_2} + 1200x^5_Z + 576x^5)$$

Summary

The solution(s) found are the following

$$y - \frac{3}{2} = x \text{RootOf}(4_Z^5 x^5 + 60_Z^4 x^5 + 345x^5_Z^3 + 940x^5_Z^2 - c_3 e^{5c_2} + 1200x^5_Z + 576x^5) \quad (1)$$




Figure 4: Slope field plot

Verification of solutions

$$y - \frac{3}{2} = x \text{RootOf} (4_Z^5 x^5 + 60_Z^4 x^5 + 345x^5_Z^3 + 940x^5_Z^2 - c_3 e^{5c_2} + 1200x^5_Z + 576x^5)$$

Verified OK.

1.2.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = -\frac{3(4x + 2y - 3)}{5x + 2y - 3}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (\text{1E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 - \frac{3(4x + 2y - 3)(b_3 - a_2)}{5x + 2y - 3} - \frac{9(4x + 2y - 3)^2 a_3}{(5x + 2y - 3)^2}$$

$$- \left(-\frac{12}{5x + 2y - 3} + \frac{60x + 30y - 45}{(5x + 2y - 3)^2} \right) (xa_2 + ya_3 + a_1) \quad (\text{5E})$$

$$- \left(-\frac{6}{5x + 2y - 3} + \frac{24x + 12y - 18}{(5x + 2y - 3)^2} \right) (xb_2 + yb_3 + b_1) = 0$$

Putting the above in normal form gives

$$60x^2 a_2 - 144x^2 a_3 + 31x^2 b_2 - 60x^2 b_3 + 48xy a_2 - 144xy a_3 + 20xy b_2 - 48xy b_3 + 12y^2 a_2 - 42y^2 a_3 + 4y^2 b_2$$

$$= 0$$

Setting the numerator to zero gives

$$\begin{aligned}
&60x^2a_2 - 144x^2a_3 + 31x^2b_2 - 60x^2b_3 + 48xya_2 - 144xya_3 + 20xyb_2 - 48xyb_3 \quad (6E) \\
&+ 12y^2a_2 - 42y^2a_3 + 4y^2b_2 - 12y^2b_3 - 72xa_2 + 216xa_3 + 6xb_1 - 30xb_2 + 81xb_3 \\
&- 6ya_1 - 36ya_2 + 117ya_3 - 12yb_2 + 36yb_3 + 9a_1 + 27a_2 - 81a_3 + 9b_2 - 27b_3 = 0
\end{aligned}$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned}
&60a_2v_1^2 + 48a_2v_1v_2 + 12a_2v_2^2 - 144a_3v_1^2 - 144a_3v_1v_2 - 42a_3v_2^2 \quad (7E) \\
&+ 31b_2v_1^2 + 20b_2v_1v_2 + 4b_2v_2^2 - 60b_3v_1^2 - 48b_3v_1v_2 - 12b_3v_2^2 \\
&- 6a_1v_2 - 72a_2v_1 - 36a_2v_2 + 216a_3v_1 + 117a_3v_2 + 6b_1v_1 - 30b_2v_1 \\
&- 12b_2v_2 + 81b_3v_1 + 36b_3v_2 + 9a_1 + 27a_2 - 81a_3 + 9b_2 - 27b_3 = 0
\end{aligned}$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned}
&(60a_2 - 144a_3 + 31b_2 - 60b_3)v_1^2 + (48a_2 - 144a_3 + 20b_2 - 48b_3)v_1v_2 \quad (8E) \\
&+ (-72a_2 + 216a_3 + 6b_1 - 30b_2 + 81b_3)v_1 + (12a_2 - 42a_3 + 4b_2 - 12b_3)v_2^2 \\
&+ (-6a_1 - 36a_2 + 117a_3 - 12b_2 + 36b_3)v_2 + 9a_1 + 27a_2 - 81a_3 + 9b_2 - 27b_3 = 0
\end{aligned}$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned}
&12a_2 - 42a_3 + 4b_2 - 12b_3 = 0 \\
&48a_2 - 144a_3 + 20b_2 - 48b_3 = 0 \\
&60a_2 - 144a_3 + 31b_2 - 60b_3 = 0 \\
&-6a_1 - 36a_2 + 117a_3 - 12b_2 + 36b_3 = 0 \\
&9a_1 + 27a_2 - 81a_3 + 9b_2 - 27b_3 = 0 \\
&-72a_2 + 216a_3 + 6b_1 - 30b_2 + 81b_3 = 0
\end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned}
 a_1 &= -\frac{3a_3}{2} \\
 a_2 &= \frac{11a_3}{2} + b_3 \\
 a_3 &= a_3 \\
 b_1 &= -\frac{3b_3}{2} \\
 b_2 &= -6a_3 \\
 b_3 &= b_3
 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}
 \xi &= x \\
 \eta &= -\frac{3}{2} + y
 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}
 \eta &= \eta - \omega(x, y) \xi \\
 &= -\frac{3}{2} + y - \left(-\frac{3(4x + 2y - 3)}{5x + 2y - 3} \right) (x) \\
 &= \frac{24x^2 + 22xy + 4y^2 - 33x - 12y + 9}{10x + 4y - 6} \\
 \xi &= 0
 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{24x^2 + 22xy + 4y^2 - 33x - 12y + 9}{10x + 4y - 6}} dy \end{aligned}$$

Which results in

$$S = \frac{2 \ln(3x + 2y - 3)}{5} + \frac{3 \ln(8x + 2y - 3)}{5}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{3(4x + 2y - 3)}{5x + 2y - 3}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{6}{15x + 10y - 15} + \frac{24}{40x + 10y - 15} \\ S_y &= \frac{4}{15x + 10y - 15} + \frac{6}{40x + 10y - 15} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{2 \ln(3x + 2y - 3)}{5} + \frac{3 \ln(8x + 2y - 3)}{5} = c_1$$

Which simplifies to

$$\frac{2 \ln(3x + 2y - 3)}{5} + \frac{3 \ln(8x + 2y - 3)}{5} = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{3(4x+2y-3)}{5x+2y-3}$ 	$R = x$ $S = \frac{2 \ln(3x + 2y - 3)}{5} +$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$\frac{2 \ln(3x + 2y - 3)}{5} + \frac{3 \ln(8x + 2y - 3)}{5} = c_1 \quad (1)$$




Figure 5: Slope field plot

Verification of solutions

$$\frac{2 \ln(3x + 2y - 3)}{5} + \frac{3 \ln(8x + 2y - 3)}{5} = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous C  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.89 (sec). Leaf size: 44

```
dsolve((12*x+6*y(x)-9)+(5*x+2*y(x)-3)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\text{RootOf}(128_Z^{25}c_1x^5 + 640_Z^{20}c_1x^5 + 800_Z^{15}c_1x^5 - 1)^5 x - 4x + \frac{3}{2}$$

✓ Solution by Mathematica

Time used: 60.12 (sec). Leaf size: 1121

`DSolve[(12*x+6*y[x]-9)+(5*x+2*y[x]-3)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{1}{2}(3 - 5x)$$

$$+ \frac{1}{2\text{Root}[\#1^{10}(11664x^{10} + 11664e^{60c_1}) - 9720\#1^8x^8 - 1080\#1^7x^7 + 3105\#1^6x^6 + 666\#1^5x^5 - 425\#1^4x^4 - 90\#1^3x^3 + 15\#1^2x^2 - 3\#1x + 1]}{1}$$

$$y(x) \rightarrow \frac{1}{2}(3 - 5x)$$

$$+ \frac{1}{2\text{Root}[\#1^{10}(11664x^{10} + 11664e^{60c_1}) - 9720\#1^8x^8 - 1080\#1^7x^7 + 3105\#1^6x^6 + 666\#1^5x^5 - 425\#1^4x^4 - 90\#1^3x^3 + 15\#1^2x^2 - 3\#1x + 1]}$$

$$y(x) \rightarrow \frac{1}{2}(3 - 5x)$$

$$+ \frac{1}{2\text{Root}[\#1^{10}(11664x^{10} + 11664e^{60c_1}) - 9720\#1^8x^8 - 1080\#1^7x^7 + 3105\#1^6x^6 + 666\#1^5x^5 - 425\#1^4x^4 - 90\#1^3x^3 + 15\#1^2x^2 - 3\#1x + 1]}$$

$$y(x) \rightarrow \frac{1}{2}(3 - 5x)$$

$$+ \frac{1}{2\text{Root}[\#1^{10}(11664x^{10} + 11664e^{60c_1}) - 9720\#1^8x^8 - 1080\#1^7x^7 + 3105\#1^6x^6 + 666\#1^5x^5 - 425\#1^4x^4 - 90\#1^3x^3 + 15\#1^2x^2 - 3\#1x + 1]}$$

$$y(x) \rightarrow \frac{1}{2}(3 - 5x)$$

$$+ \frac{1}{2\text{Root}[\#1^{10}(11664x^{10} + 11664e^{60c_1}) - 9720\#1^8x^8 - 1080\#1^7x^7 + 3105\#1^6x^6 + 666\#1^5x^5 - 425\#1^4x^4 - 90\#1^3x^3 + 15\#1^2x^2 - 3\#1x + 1]}$$

$$y(x) \rightarrow \frac{1}{2}(3 - 5x)$$

$$+ \frac{1}{2\text{Root}[\#1^{10}(11664x^{10} + 11664e^{60c_1}) - 9720\#1^8x^8 - 1080\#1^7x^7 + 3105\#1^6x^6 + 666\#1^5x^5 - 425\#1^4x^4 - 90\#1^3x^3 + 15\#1^2x^2 - 3\#1x + 1]}$$

$$y(x) \rightarrow \frac{1}{2}(3 - 5x)$$

$$+ \frac{1}{2\text{Root}[\#1^{10}(11664x^{10} + 11664e^{60c_1}) - 9720\#1^8x^8 - 1080\#1^7x^7 + 3105\#1^6x^6 + 666\#1^5x^5 - 425\#1^4x^4 - 90\#1^3x^3 + 15\#1^2x^2 - 3\#1x + 1]}$$

$$y(x) \rightarrow \frac{1}{2}(3 - 5x)$$

$$+ \frac{1}{2\text{Root}[\#1^{10}(11664x^{10} + 11664e^{60c_1}) - 9720\#1^8x^8 - 1080\#1^7x^7 + 3105\#1^6x^6 + 666\#1^5x^5 - 425\#1^4x^4 - 90\#1^3x^3 + 15\#1^2x^2 - 3\#1x + 1]}$$

$$y(x) \rightarrow \frac{1}{2}(3 - 5x)$$

$$+ \frac{1}{2\text{Root}[\#1^{10}(11664x^{10} + 11664e^{60c_1}) - 9720\#1^8x^8 - 1080\#1^7x^7 + 3105\#1^6x^6 + 666\#1^5x^5 - 425\#1^4x^4 - 90\#1^3x^3 + 15\#1^2x^2 - 3\#1x + 1]}$$

$$y(x) \rightarrow \frac{1}{2}(3 - 5x)$$

$$+ \frac{1}{2\text{Root}[\#1^{10}(11664x^{10} + 11664e^{60c_1}) - 9720\#1^8x^8 - 1080\#1^7x^7 + 3105\#1^6x^6 + 666\#1^5x^5 - 425\#1^4x^4 - 90\#1^3x^3 + 15\#1^2x^2 - 3\#1x + 1]}$$

1.3 problem Problem 3

1.3.1 Solving as first order ode lie symmetry calculated ode 28

Internal problem ID [12114]

Internal file name [OUTPUT/10766_Monday_September_11_2023_12_50_32_AM_68973649/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 3.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"first_order_ode_lie_symmetry_calculated"**

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _dAlembert]
```

$$y'x - y - \sqrt{x^2 + y^2} = 0$$

1.3.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{y + \sqrt{x^2 + y^2}}{x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \tag{1E}$$

$$\eta = xb_2 + yb_3 + b_1 \tag{2E}$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} & b_2 + \frac{(y + \sqrt{x^2 + y^2})(b_3 - a_2)}{x} - \frac{(y + \sqrt{x^2 + y^2})^2 a_3}{x^2} \\ & - \left(\frac{1}{\sqrt{x^2 + y^2}} - \frac{y + \sqrt{x^2 + y^2}}{x^2} \right) (xa_2 + ya_3 + a_1) \\ & - \frac{\left(1 + \frac{y}{\sqrt{x^2 + y^2}}\right) (xb_2 + yb_3 + b_1)}{x} = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\begin{aligned} & \frac{(x^2 + y^2)^{\frac{3}{2}} a_3 + x^3 a_2 - x^3 b_3 + 2x^2 y a_3 + x^2 y b_2 + y^3 a_3 + \sqrt{x^2 + y^2} x b_1 - \sqrt{x^2 + y^2} y a_1 + x y b_1 - y^2 a_1}{\sqrt{x^2 + y^2} x^2} \\ & = 0 \end{aligned}$$

Setting the numerator to zero gives

$$\begin{aligned} & -(x^2 + y^2)^{\frac{3}{2}} a_3 - x^3 a_2 + x^3 b_3 - 2x^2 y a_3 - x^2 y b_2 - y^3 a_3 \\ & - \sqrt{x^2 + y^2} x b_1 + \sqrt{x^2 + y^2} y a_1 - x y b_1 + y^2 a_1 = 0 \end{aligned} \quad (6E)$$

Simplifying the above gives

$$\begin{aligned} & -(x^2 + y^2)^{\frac{3}{2}} a_3 + (x^2 + y^2) x b_3 - (x^2 + y^2) y a_3 - x^3 a_2 - x^2 y a_3 - x^2 y b_2 \\ & - x y^2 b_3 + (x^2 + y^2) a_1 - \sqrt{x^2 + y^2} x b_1 + \sqrt{x^2 + y^2} y a_1 - x^2 a_1 - x y b_1 = 0 \end{aligned} \quad (6E)$$

Since the PDE has radicals, simplifying gives

$$\begin{aligned} & -x^3 a_2 + x^3 b_3 - x^2 \sqrt{x^2 + y^2} a_3 - 2x^2 y a_3 - x^2 y b_2 - \sqrt{x^2 + y^2} y^2 a_3 \\ & - y^3 a_3 - \sqrt{x^2 + y^2} x b_1 - x y b_1 + \sqrt{x^2 + y^2} y a_1 + y^2 a_1 = 0 \end{aligned}$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y, \sqrt{x^2 + y^2}\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\left\{ x = v_1, y = v_2, \sqrt{x^2 + y^2} = v_3 \right\}$$

The above PDE (6E) now becomes

$$\begin{aligned} -v_1^3 a_2 - 2v_1^2 v_2 a_3 - v_1^2 v_3 a_3 - v_2^3 a_3 - v_3 v_2^2 a_3 - v_1^2 v_2 b_2 \\ + v_1^3 b_3 + v_2^2 a_1 + v_3 v_2 a_1 - v_1 v_2 b_1 - v_3 v_1 b_1 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3\}$$

Equation (7E) now becomes

$$\begin{aligned} (b_3 - a_2) v_1^3 + (-2a_3 - b_2) v_1^2 v_2 - v_1^2 v_3 a_3 - v_1 v_2 b_1 \\ - v_3 v_1 b_1 - v_2^3 a_3 - v_3 v_2^2 a_3 + v_2^2 a_1 + v_3 v_2 a_1 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} a_1 &= 0 \\ -a_3 &= 0 \\ -b_1 &= 0 \\ -2a_3 - b_2 &= 0 \\ b_3 - a_2 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= b_3 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}\xi &= x \\ \eta &= y\end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}\eta &= \eta - \omega(x, y) \xi \\ &= y - \left(\frac{y + \sqrt{x^2 + y^2}}{x} \right) (x) \\ &= -\sqrt{x^2 + y^2} \\ \xi &= 0\end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{-\sqrt{x^2 + y^2}} dy\end{aligned}$$

Which results in

$$S = -\ln \left(y + \sqrt{x^2 + y^2} \right)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y) S_y}{R_x + \omega(x, y) R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y + \sqrt{x^2 + y^2}}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{x}{\sqrt{x^2 + y^2} (y + \sqrt{x^2 + y^2})} \\ S_y &= -\frac{1}{\sqrt{x^2 + y^2}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -\frac{2(\sqrt{x^2 + y^2} y + x^2 + y^2)}{x\sqrt{x^2 + y^2} (y + \sqrt{x^2 + y^2})} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{2}{R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -2 \ln(R) + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\ln(y + \sqrt{x^2 + y^2}) = -2 \ln(x) + c_1$$



Which simplifies to

$$-\ln(y + \sqrt{x^2 + y^2}) = -2 \ln(x) + c_1$$

Which gives

$$y = -\frac{e^{-c_1}(e^{2c_1} - x^2)}{2}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y + \sqrt{x^2 + y^2}}{x}$ 	$R = x$ $S = -\ln\left(y + \sqrt{x^2 + y^2}\right)$	$\frac{dS}{dR} = -\frac{2}{R}$ 

Summary

The solution(s) found are the following

$$y = -\frac{e^{-c_1}(e^{2c_1} - x^2)}{2} \tag{1}$$




Figure 6: Slope field plot

Verification of solutions

$$y = -\frac{e^{-c_1}(e^{2c_1} - x^2)}{2}$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying homogeneous types:
trying homogeneous G
1st order, trying the canonical coordinates of the invariance group
<- 1st order, canonical coordinates successful
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 26

```
dsolve(x*diff(y(x),x)=y(x)+sqrt(x^2+y(x)^2),y(x), singsol=all)
```

$$\frac{-c_1 x^2 + y(x) + \sqrt{y(x)^2 + x^2}}{x^2} = 0$$

✓ Solution by Mathematica

Time used: 0.603 (sec). Leaf size: 27

```
DSolve[x*y'[x]==y[x]+Sqrt[x^2+y[x]^2],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^{-c_1} (-1 + e^{2c_1} x^2)$$

1.4 problem Problem 4

1.4.1	Solving as linear ode	36
1.4.2	Solving as differentialType ode	38
1.4.3	Solving as first order ode lie symmetry lookup ode	40
1.4.4	Solving as exact ode	44
1.4.5	Maple step by step solution	48

Internal problem ID [12115]

Internal file name [OUTPUT/10767_Monday_September_11_2023_12_50_34_AM_17149971/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 4.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exact"**, **"linear"**, **"differentialType"**, **"first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_linear]`

$$y'x + y = x^3$$

1.4.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{1}{x}$$
$$q(x) = x^2$$

Hence the ode is

$$y' + \frac{y}{x} = x^2$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{1}{x} dx} \\ &= x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (x^2) \\ \frac{d}{dx}(xy) &= (x) (x^2) \\ d(xy) &= x^3 dx\end{aligned}$$

Integrating gives

$$\begin{aligned}xy &= \int x^3 dx \\ xy &= \frac{x^4}{4} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = x$ results in

$$y = \frac{x^3}{4} + \frac{c_1}{x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^3}{4} + \frac{c_1}{x} \tag{1}$$




Figure 7: Slope field plot

Verification of solutions

$$y = \frac{x^3}{4} + \frac{c_1}{x}$$

Verified OK.

1.4.2 Solving as differentialType ode

Writing the ode as

$$y' = \frac{x^3 - y}{x} \tag{1}$$

Which becomes

$$0 = (-x) dy + (x^3 - y) dx \tag{2}$$

But the RHS is complete differential because

$$(-x) dy + (x^3 - y) dx = d\left(\frac{1}{4}x^4 - xy\right)$$

Hence (2) becomes

$$0 = d\left(\frac{1}{4}x^4 - xy\right)$$

Integrating both sides gives gives these solutions

$$y = \frac{x^4 + 4c_1}{4x} + c_1$$

Summary

The solution(s) found are the following

$$y = \frac{x^4 + 4c_1}{4x} + c_1 \tag{1}$$




Figure 8: Slope field plot

Verification of solutions

$$y = \frac{x^4 + 4c_1}{4x} + c_1$$

Verified OK.

1.4.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{-x^3 + y}{x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 4: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{1}{x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{1}{x}} dy\end{aligned}$$

Which results in

$$S = xy$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y}\tag{2}$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{-x^3 + y}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 1 \\R_y &= 0 \\S_x &= y \\S_y &= x\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = x^3 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = R^3$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{R^4}{4} + c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$yx = \frac{x^4}{4} + c_1$$



Which simplifies to

$$yx = \frac{x^4}{4} + c_1$$

Which gives

$$y = \frac{x^4 + 4c_1}{4x}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{-x^3+y}{x}$ 	$R = x$ $S = xy$	$\frac{dS}{dR} = R^3$ 

Summary

The solution(s) found are the following

$$y = \frac{x^4 + 4c_1}{4x} \tag{1}$$




Figure 9: Slope field plot

Verification of solutions

$$y = \frac{x^4 + 4c_1}{4x}$$

Verified OK.

1.4.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(x) dy &= (x^3 - y) dx \\ (-x^3 + y) dx + (x) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -x^3 + y \\ N(x, y) &= x\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x^3 + y) \\ &= 1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x) \\ &= 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int -x^3 + y dx$$

$$\phi = -\frac{1}{4}x^4 + xy + f(y) \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = x + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = x$. Therefore equation (4) becomes

$$x = x + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{1}{4}x^4 + xy + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{1}{4}x^4 + xy$$

The solution becomes

$$y = \frac{x^4 + 4c_1}{4x}$$

Summary

The solution(s) found are the following

$$y = \frac{x^4 + 4c_1}{4x} \tag{1}$$




Figure 10: Slope field plot

Verification of solutions

$$y = \frac{x^4 + 4c_1}{4x}$$

Verified OK.

1.4.5 Maple step by step solution

Let's solve

$$y'x + y = x^3$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -\frac{y}{x} + x^2$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + \frac{y}{x} = x^2$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' + \frac{y}{x} \right) = \mu(x) x^2$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' + \frac{y}{x} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{\mu(x)}{x}$$

- Solve to find the integrating factor

$$\mu(x) = x$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \mu(x) x^2 dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \mu(x) x^2 dx + c_1$$

- Solve for y

$$y = \frac{\int \mu(x) x^2 dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = x$

$$y = \frac{\int x^3 dx + c_1}{x}$$

- Evaluate the integrals on the rhs

$$y = \frac{\frac{x^4}{4} + c_1}{x}$$

- Simplify

$$y = \frac{x^4 + 4c_1}{4x}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(x*diff(y(x),x)+y(x)=x^3,y(x), singsol=all)
```

$$y(x) = \frac{x^4 + 4c_1}{4x}$$

✓ Solution by Mathematica

Time used: 0.048 (sec). Leaf size: 19

```
DSolve[x*y'[x]+y[x]==x^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{4} + \frac{c_1}{x}$$

1.5 problem Problem 5

- 1.5.1 Solving as first order ode lie symmetry calculated ode 50
- 1.5.2 Solving as exact ode 55

Internal problem ID [12116]

Internal file name [OUTPUT/10768_Monday_September_11_2023_12_50_35_AM_98222464/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 5.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exactWithIntegrationFactor", "first_order_ode_lie_symmetry_calculated"**

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _rational, [_Abel, `2nd type`, `class B`]]
```

$$y - y'x - y'yx^2 = 0$$

1.5.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{y}{x(xy + 1)}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \tag{1E}$$

$$\eta = xb_2 + yb_3 + b_1 \tag{2E}$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 + \frac{y(b_3 - a_2)}{x(xy + 1)} - \frac{y^2 a_3}{x^2(xy + 1)^2} \\ - \left(-\frac{y}{x^2(xy + 1)} - \frac{y^2}{x(xy + 1)^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(\frac{1}{x(xy + 1)} - \frac{y}{(xy + 1)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{x^4 y^2 b_2 + 2x^3 y b_2 + x^2 y^2 a_2 + x^2 y^2 b_3 + 2x y^3 a_3 + 2x y^2 a_1 - x b_1 + y a_1}{x^2 (xy + 1)^2} = 0$$

Setting the numerator to zero gives

$$x^4 y^2 b_2 + 2x^3 y b_2 + x^2 y^2 a_2 + x^2 y^2 b_3 + 2x y^3 a_3 + 2x y^2 a_1 - x b_1 + y a_1 = 0 \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$b_2 v_1^4 v_2^2 + a_2 v_1^2 v_2^2 + 2a_3 v_1 v_2^3 + 2b_2 v_1^3 v_2 + b_3 v_1^2 v_2^2 + 2a_1 v_1 v_2^2 + a_1 v_2 - b_1 v_1 = 0 \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$b_2 v_1^4 v_2^2 + 2b_2 v_1^3 v_2 + (a_2 + b_3) v_1^2 v_2^2 + 2a_3 v_1 v_2^3 + 2a_1 v_1 v_2^2 - b_1 v_1 + a_1 v_2 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} a_1 &= 0 \\ b_2 &= 0 \\ 2a_1 &= 0 \\ 2a_3 &= 0 \\ -b_1 &= 0 \\ 2b_2 &= 0 \\ a_2 + b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= -b_3 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the ansatz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= -x \\ \eta &= y \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= y - \left(\frac{y}{x(xy+1)} \right) (-x) \\ &= \frac{xy^2 + 2y}{xy+1} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{xy^2+2y}{xy+1}} dy \end{aligned}$$

Which results in

$$S = \frac{\ln(y(xy+2))}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y}{x(xy+1)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{y}{2xy+4} \\ S_y &= \frac{xy+1}{y(xy+2)} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{1}{2x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{2R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{\ln(R)}{2} + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{\ln(y)}{2} + \frac{\ln(yx + 2)}{2} = \frac{\ln(x)}{2} + c_1$$

Which simplifies to

$$\frac{\ln(y)}{2} + \frac{\ln(yx + 2)}{2} = \frac{\ln(x)}{2} + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y}{x(xy+1)}$ 	$R = x$ $S = \frac{\ln(y)}{2} + \frac{\ln(xy + 2)}{2}$	$\frac{dS}{dR} = \frac{1}{2R}$ 

Summary

The solution(s) found are the following

$$\frac{\ln(y)}{2} + \frac{\ln(yx + 2)}{2} = \frac{\ln(x)}{2} + c_1 \quad (1)$$




Figure 11: Slope field plot

Verification of solutions

$$\frac{\ln(y)}{2} + \frac{\ln(yx + 2)}{2} = \frac{\ln(x)}{2} + c_1$$

Verified OK.

1.5.2 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (-x^2y - x) dy &= (-y) dx \\ (y) dx + (-x^2y - x) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= y \\ N(x, y) &= -x^2y - x \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(y) \\ &= 1 \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(-x^2y - x) \\ &= -2xy - 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= -\frac{1}{x(xy+1)}((1) - (-2xy - 1)) \\ &= -\frac{2}{x}\end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dx} \\ &= e^{\int -\frac{2}{x} dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{-2\ln(x)} \\ &= \frac{1}{x^2}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= \frac{1}{x^2}(y) \\ &= \frac{y}{x^2}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{1}{x^2}(-x^2y - x) \\ &= \frac{-xy - 1}{x}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(\frac{y}{x^2}\right) + \left(\frac{-xy-1}{x}\right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{y}{x^2} dx \\ \phi &= -\frac{y}{x} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = -\frac{1}{x} + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{-xy-1}{x}$. Therefore equation (4) becomes

$$\frac{-xy-1}{x} = -\frac{1}{x} + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -y$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int (-y) dy \\ f(y) &= -\frac{y^2}{2} + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{y}{x} - \frac{y^2}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{y}{x} - \frac{y^2}{2}$$

Summary

The solution(s) found are the following

$$-\frac{y}{x} - \frac{y^2}{2} = c_1 \tag{1}$$




Figure 12: Slope field plot

Verification of solutions

$$-\frac{y}{x} - \frac{y^2}{2} = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous G  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 49

```
dsolve(y(x)-x*diff(y(x),x)=x^2*y(x)*diff(y(x),x),y(x), singsol=all)
```

$$y(x) = \frac{-c_1 + \sqrt{c_1^2 + x^2}}{c_1 x}$$
$$y(x) = \frac{-c_1 - \sqrt{c_1^2 + x^2}}{c_1 x}$$

✓ Solution by Mathematica

Time used: 0.786 (sec). Leaf size: 68

```
DSolve[y[x]-x*y'[x]==x^2*y[x]*y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1 + \sqrt{\frac{1}{x^2}x\sqrt{1 + c_1x^2}}}{x}$$
$$y(x) \rightarrow -\frac{1}{x} + \sqrt{\frac{1}{x^2}\sqrt{1 + c_1x^2}}$$
$$y(x) \rightarrow 0$$

1.6 problem Problem 6

1.6.1	Solving as linear ode	61
1.6.2	Solving as first order ode lie symmetry lookup ode	63
1.6.3	Solving as exact ode	67
1.6.4	Maple step by step solution	71

Internal problem ID [12117]

Internal file name [OUTPUT/10769_Monday_September_11_2023_12_50_36_AM_52281106/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 6.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$x' + 3x = e^{2t}$$

1.6.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$x' + p(t)x = q(t)$$

Where here

$$p(t) = 3$$
$$q(t) = e^{2t}$$

Hence the ode is

$$x' + 3x = e^{2t}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int 3dt} \\ &= e^{3t}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dt}(\mu x) &= (\mu) (e^{2t}) \\ \frac{d}{dt}(e^{3t}x) &= (e^{3t}) (e^{2t}) \\ d(e^{3t}x) &= e^{5t} dt\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{3t}x &= \int e^{5t} dt \\ e^{3t}x &= \frac{e^{5t}}{5} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{3t}$ results in

$$x = \frac{e^{-3t}e^{5t}}{5} + e^{-3t}c_1$$

which simplifies to

$$x = \frac{(e^{5t} + 5c_1) e^{-3t}}{5}$$

Summary

The solution(s) found are the following

$$x = \frac{(e^{5t} + 5c_1) e^{-3t}}{5} \tag{1}$$




Figure 13: Slope field plot

Verification of solutions

$$x = \frac{(e^{5t} + 5c_1) e^{-3t}}{5}$$

Verified OK.

1.6.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$\begin{aligned} x' &= -3x + e^{2t} \\ x' &= \omega(t, x) \end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_t + \omega(\eta_x - \xi_t) - \omega^2 \xi_x - \omega_t \xi - \omega_x \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 7: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(t, x) &= 0 \\ \eta(t, x) &= e^{-3t}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(t, x) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dt}{\xi} = \frac{dx}{\eta} = dS\tag{1}$$

The above comes from the requirements that $(\xi \frac{\partial}{\partial t} + \eta \frac{\partial}{\partial x})S(t, x) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = t$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-3t}} dy \end{aligned}$$

Which results in

$$S = e^{3t} x$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_t + \omega(t, x)S_x}{R_t + \omega(t, x)R_x} \quad (2)$$

Where in the above R_t, R_x, S_t, S_x are all partial derivatives and $\omega(t, x)$ is the right hand side of the original ode given by

$$\omega(t, x) = -3x + e^{2t}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_t &= 1 \\ R_x &= 0 \\ S_t &= 3e^{3t}x \\ S_x &= e^{3t} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = e^{5t} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for t, x in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = e^{5R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{e^{5R}}{5} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to t, x coordinates. This results in

$$x e^{3t} = \frac{e^{5t}}{5} + c_1$$



Which simplifies to

$$x e^{3t} = \frac{e^{5t}}{5} + c_1$$

Which gives

$$x = \frac{(e^{5t} + 5c_1) e^{-3t}}{5}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in t, x coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dx}{dt} = -3x + e^{2t}$ 	$R = t$ $S = e^{3t}x$	$\frac{dS}{dR} = e^{5R}$ 

Summary

The solution(s) found are the following

$$x = \frac{(e^{5t} + 5c_1) e^{-3t}}{5} \quad (1)$$




Figure 14: Slope field plot

Verification of solutions

$$x = \frac{(e^{5t} + 5c_1) e^{-3t}}{5}$$

Verified OK.

1.6.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(t, x) dt + N(t, x) dx = 0 \tag{1A}$$

Therefore

$$\begin{aligned}dx &= (-3x + e^{2t}) dt \\ (3x - e^{2t}) dt + dx &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(t, x) &= 3x - e^{2t} \\ N(t, x) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial x} = \frac{\partial N}{\partial t}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial x} &= \frac{\partial}{\partial x}(3x - e^{2t}) \\ &= 3\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial t} &= \frac{\partial}{\partial t}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial x} \neq \frac{\partial N}{\partial t}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial x} - \frac{\partial N}{\partial t} \right) \\ &= 1((3) - (0)) \\ &= 3 \end{aligned}$$

Since A does not depend on x , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dt} \\ &= e^{\int 3 dt} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{3t} \\ &= e^{3t} \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= e^{3t}(3x - e^{2t}) \\ &= (3x - e^{2t}) e^{3t} \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= e^{3t}(1) \\ &= e^{3t} \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dx}{dt} &= 0 \\ ((3x - e^{2t}) e^{3t}) + (e^{3t}) \frac{dx}{dt} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(t, x)$

$$\frac{\partial \phi}{\partial t} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial x} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. t gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial t} dt &= \int \bar{M} dt \\ \int \frac{\partial \phi}{\partial t} dt &= \int (3x - e^{2t}) e^{3t} dt \\ \phi &= -\frac{e^{5t}}{5} + e^{3t}x + f(x)\end{aligned}\quad (3)$$

Where $f(x)$ is used for the constant of integration since ϕ is a function of both t and x . Taking derivative of equation (3) w.r.t x gives

$$\frac{\partial \phi}{\partial x} = e^{3t} + f'(x) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial x} = e^{3t}$. Therefore equation (4) becomes

$$e^{3t} = e^{3t} + f'(x) \quad (5)$$

Solving equation (5) for $f'(x)$ gives

$$f'(x) = 0$$

Therefore

$$f(x) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(x)$ into equation (3) gives ϕ

$$\phi = -\frac{e^{5t}}{5} + e^{3t}x + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{e^{5t}}{5} + e^{3t}x$$

The solution becomes

$$x = \frac{(e^{5t} + 5c_1) e^{-3t}}{5}$$

Summary

The solution(s) found are the following

$$x = \frac{(e^{5t} + 5c_1) e^{-3t}}{5} \quad (1)$$




Figure 15: Slope field plot

Verification of solutions

$$x = \frac{(e^{5t} + 5c_1) e^{-3t}}{5}$$

Verified OK.

1.6.4 Maple step by step solution

Let's solve

$$x' + 3x = e^{2t}$$

- Highest derivative means the order of the ODE is 1

$$x'$$

- Isolate the derivative

$$x' = -3x + e^{2t}$$

- Group terms with x on the lhs of the ODE and the rest on the rhs of the ODE

$$x' + 3x = e^{2t}$$

- The ODE is linear; multiply by an integrating factor $\mu(t)$

$$\mu(t)(x' + 3x) = \mu(t)e^{2t}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dt}(\mu(t)x)$

$$\mu(t)(x' + 3x) = \mu'(t)x + \mu(t)x'$$

- Isolate $\mu'(t)$

$$\mu'(t) = 3\mu(t)$$

- Solve to find the integrating factor

$$\mu(t) = e^{3t}$$

- Integrate both sides with respect to t

$$\int \left(\frac{d}{dt}(\mu(t)x) \right) dt = \int \mu(t)e^{2t} dt + c_1$$

- Evaluate the integral on the lhs

$$\mu(t)x = \int \mu(t)e^{2t} dt + c_1$$

- Solve for x

$$x = \frac{\int \mu(t)e^{2t} dt + c_1}{\mu(t)}$$

- Substitute $\mu(t) = e^{3t}$

$$x = \frac{\int e^{2t}e^{3t} dt + c_1}{e^{3t}}$$

- Evaluate the integrals on the rhs

$$x = \frac{\frac{e^{5t}}{5} + c_1}{e^{3t}}$$

- Simplify

$$x = \frac{(e^{5t} + 5c_1)e^{-3t}}{5}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(x(t),t)+3*x(t)=exp(2*t),x(t), singsol=all)
```

$$x(t) = \frac{(e^{5t} + 5c_1)e^{-3t}}{5}$$

✓ Solution by Mathematica

Time used: 0.07 (sec). Leaf size: 23

```
DSolve[x'[t]+3*x[t]==Exp[2*t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{e^{2t}}{5} + c_1 e^{-3t}$$

1.7 problem Problem 7

1.7.1	Solving as linear ode	74
1.7.2	Solving as first order ode lie symmetry lookup ode	76
1.7.3	Solving as exact ode	80
1.7.4	Maple step by step solution	84

Internal problem ID [12118]

Internal file name [OUTPUT/10770_Tuesday_September_12_2023_08_51_39_AM_75656588/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 7.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_linear]`

$$\sin(x)y + \cos(x)y' = 1$$

1.7.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \tan(x)$$

$$q(x) = \sec(x)$$

Hence the ode is

$$y' + \tan(x)y = \sec(x)$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \tan(x) dx} \\ &= \frac{1}{\cos(x)}\end{aligned}$$

Which simplifies to

$$\mu = \sec(x)$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) (\sec(x)) \\ \frac{d}{dx}(\sec(x) y) &= (\sec(x)) (\sec(x)) \\ d(\sec(x) y) &= \sec(x)^2 dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\sec(x) y &= \int \sec(x)^2 dx \\ \sec(x) y &= \tan(x) + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \sec(x)$ results in

$$y = \cos(x) \tan(x) + c_1 \cos(x)$$

which simplifies to

$$y = c_1 \cos(x) + \sin(x)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(x) + \sin(x) \tag{1}$$




Figure 16: Slope field plot

Verification of solutions

$$y = c_1 \cos(x) + \sin(x)$$

Verified OK.

1.7.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{\sin(x)y - 1}{\cos(x)}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 10: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \cos(x)\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\cos(x)} dy \end{aligned}$$

Which results in

$$S = \frac{y}{\cos(x)}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{\sin(x)y - 1}{\cos(x)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \sec(x) \tan(x) y \\ S_y &= \sec(x) \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \sec(x)^2 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \sec(R)^2$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \tan(R) + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\sec(x) y = \tan(x) + c_1$$



Which simplifies to

$$\sec(x) y = \tan(x) + c_1$$

Which gives

$$y = \frac{\tan(x) + c_1}{\sec(x)}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{\sin(x)y-1}{\cos(x)}$ 	$R = x$ $S = \sec(x) y$	$\frac{dS}{dR} = \sec(R)^2$ 

Summary

The solution(s) found are the following

$$y = \frac{\tan(x) + c_1}{\sec(x)} \quad (1)$$




Figure 17: Slope field plot

Verification of solutions

$$y = \frac{\tan(x) + c_1}{\sec(x)}$$

Verified OK.

1.7.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(\cos(x)) dy &= (-\sin(x)y + 1) dx \\ (\sin(x)y - 1) dx + (\cos(x)) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= \sin(x)y - 1 \\ N(x, y) &= \cos(x)\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(\sin(x)y - 1) \\ &= \sin(x)\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(\cos(x)) \\ &= -\sin(x)\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \sec(x) ((\sin(x)) - (-\sin(x))) \\ &= 2 \tan(x) \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A \, dx} \\ &= e^{\int 2 \tan(x) \, dx} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-2 \ln(\cos(x))} \\ &= \sec(x)^2 \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= \sec(x)^2 (\sin(x) y - 1) \\ &= (\sin(x) y - 1) \sec(x)^2 \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= \sec(x)^2 (\cos(x)) \\ &= \sec(x) \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ ((\sin(x) y - 1) \sec(x)^2) + (\sec(x)) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int (\sin(x)y - 1) \sec(x)^2 dx \\ \phi &= \sec(x)y - \tan(x) + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \sec(x) + f'(y)\quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \sec(x)$. Therefore equation (4) becomes

$$\sec(x) = \sec(x) + f'(y)\quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \sec(x)y - \tan(x) + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \sec(x)y - \tan(x)$$

The solution becomes

$$y = \frac{\tan(x) + c_1}{\sec(x)}$$

Summary

The solution(s) found are the following

$$y = \frac{\tan(x) + c_1}{\sec(x)} \quad (1)$$




Figure 18: Slope field plot

Verification of solutions

$$y = \frac{\tan(x) + c_1}{\sec(x)}$$

Verified OK.

1.7.4 Maple step by step solution

Let's solve

$$\sin(x)y + \cos(x)y' = 1$$

- Highest derivative means the order of the ODE is 1
- y'
- Isolate the derivative

$$y' = -\frac{\sin(x)y}{\cos(x)} + \frac{1}{\cos(x)}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + \frac{\sin(x)y}{\cos(x)} = \frac{1}{\cos(x)}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' + \frac{\sin(x)y}{\cos(x)} \right) = \frac{\mu(x)}{\cos(x)}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' + \frac{\sin(x)y}{\cos(x)} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{\mu(x)\sin(x)}{\cos(x)}$$

- Solve to find the integrating factor

$$\mu(x) = \frac{1}{\cos(x)}$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \frac{\mu(x)}{\cos(x)} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \frac{\mu(x)}{\cos(x)} dx + c_1$$

- Solve for y

$$y = \frac{\int \frac{\mu(x)}{\cos(x)} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = \frac{1}{\cos(x)}$

$$y = \cos(x) \left(\int \frac{1}{\cos(x)^2} dx + c_1 \right)$$

- Evaluate the integrals on the rhs

$$y = \cos(x) (\tan(x) + c_1)$$

- Simplify

$$y = c_1 \cos(x) + \sin(x)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(y(x)*sin(x)+diff(y(x),x)*cos(x)=1,y(x), singsol=all)
```

$$y(x) = c_1 \cos(x) + \sin(x)$$

✓ Solution by Mathematica

Time used: 0.066 (sec). Leaf size: 13

```
DSolve[y[x]*Sin[x]+y'[x]*Cos[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \sin(x) + c_1 \cos(x)$$

1.8 problem Problem 8

1.8.1	Solving as separable ode	87
1.8.2	Solving as first order special form ID 1 ode	89
1.8.3	Solving as first order ode lie symmetry lookup ode	90
1.8.4	Solving as exact ode	94
1.8.5	Maple step by step solution	98

Internal problem ID [12119]

Internal file name [OUTPUT/10771_Tuesday_September_12_2023_08_51_40_AM_30239279/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 8.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "separable", "first order special form ID 1", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y' - e^{-y+x} = 0$$

1.8.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= e^{-y}e^x\end{aligned}$$

Where $f(x) = e^x$ and $g(y) = e^{-y}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{e^{-y}} dy &= e^x dx \\ \int \frac{1}{e^{-y}} dy &= \int e^x dx \\ e^y &= e^x + c_1\end{aligned}$$

Which results in

$$y = \ln(e^x + c_1)$$

Summary

The solution(s) found are the following

$$y = \ln(e^x + c_1) \tag{1}$$




Figure 19: Slope field plot

Verification of solutions

$$y = \ln(e^x + c_1)$$

Verified OK.

1.8.2 Solving as first order special form ID 1 ode

Writing the ode as

$$y' = e^{-y+x} \quad (1)$$

And using the substitution $u = e^y$ then

$$u' = y'e^y$$

The above shows that

$$\begin{aligned} y' &= u'(x) e^{-y} \\ &= \frac{u'(x)}{u} \end{aligned}$$

Substituting this in (1) gives

$$\frac{u'(x)}{u} = \frac{e^x}{u}$$

The above simplifies to

$$u'(x) = e^x \quad (2)$$

Now ode (2) is solved for $u(x)$ Integrating both sides gives

$$\begin{aligned} u(x) &= \int e^x dx \\ &= e^x + c_1 \end{aligned}$$

Substituting the solution found for $u(x)$ in $u = e^y$ gives

$$\begin{aligned} y &= \ln(u(x)) \\ &= \ln(e^x + c_1) \\ &= \ln(e^x + c_1) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \ln(e^x + c_1) \quad (1)$$




Figure 20: Slope field plot

Verification of solutions

$$y = \ln(e^x + c_1)$$

Verified OK.

1.8.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = e^{-y+x}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 13: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= e^{-x} \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{e^{-x}} dx \end{aligned}$$

Which results in

$$S = e^x$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = e^{-y+x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 0 \\ R_y &= 1 \\ S_x &= e^x \\ S_y &= 0 \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = e^y \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = e^R$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = e^R + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$e^x = e^y + c_1$$



Which simplifies to

$$e^x = e^y + c_1$$

Which gives

$$y = \ln(e^x - c_1)$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = e^{-y+x}$ 	$R = y$ $S = e^x$	$\frac{dS}{dR} = e^R$ 

Summary

The solution(s) found are the following

$$y = \ln(e^x - c_1) \quad (1)$$




Figure 21: Slope field plot

Verification of solutions

$$y = \ln(e^x - c_1)$$

Verified OK.

1.8.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}(e^y) dy &= (e^x) dx \\ (-e^x) dx + (e^y) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -e^x \\ N(x, y) &= e^y\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-e^x) \\ &= 0\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(e^y) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int -e^x dx$$

$$\phi = -e^x + f(y) \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = e^y$. Therefore equation (4) becomes

$$e^y = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = e^y$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int (e^y) dy$$

$$f(y) = e^y + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -e^x + e^y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -e^x + e^y$$

The solution becomes

$$y = \ln(e^x + c_1)$$

Summary

The solution(s) found are the following

$$y = \ln(e^x + c_1) \tag{1}$$




Figure 22: Slope field plot

Verification of solutions

$$y = \ln(e^x + c_1)$$

Verified OK.

1.8.5 Maple step by step solution

Let's solve

$$y' - e^{-y+x} = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$y'e^y = e^x$$

- Integrate both sides with respect to x

$$\int y'e^y dx = \int e^x dx + c_1$$

- Evaluate integral

$$e^y = e^x + c_1$$

- Solve for y

$$y = \ln(e^x + c_1)$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
<- separable successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 9

```
dsolve(diff(y(x),x)=exp(x-y(x)),y(x), singsol=all)
```

$$y(x) = \ln(e^x + c_1)$$

✓ Solution by Mathematica

Time used: 1.307 (sec). Leaf size: 12

```
DSolve[y'[x]==Exp[x-y[x]],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \log(e^x + c_1)$$

1.9 problem Problem 9

1.9.1	Solving as linear ode	100
1.9.2	Solving as first order ode lie symmetry lookup ode	102
1.9.3	Solving as exact ode	106
1.9.4	Maple step by step solution	110

Internal problem ID [12120]

Internal file name [OUTPUT/10772_Tuesday_September_12_2023_08_51_41_AM_93042087/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 9.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$-x + x' = \sin(t)$$

1.9.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$x' + p(t)x = q(t)$$

Where here

$$p(t) = -1$$

$$q(t) = \sin(t)$$

Hence the ode is

$$-x + x' = \sin(t)$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int(-1)dt} \\ &= e^{-t}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dt}(\mu x) &= (\mu) (\sin(t)) \\ \frac{d}{dt}(e^{-t}x) &= (e^{-t}) (\sin(t)) \\ d(e^{-t}x) &= (e^{-t} \sin(t)) dt\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{-t}x &= \int e^{-t} \sin(t) dt \\ e^{-t}x &= -\frac{e^{-t} \cos(t)}{2} - \frac{e^{-t} \sin(t)}{2} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{-t}$ results in

$$x = e^t \left(-\frac{e^{-t} \cos(t)}{2} - \frac{e^{-t} \sin(t)}{2} \right) + c_1 e^t$$

which simplifies to

$$x = c_1 e^t - \frac{\sin(t)}{2} - \frac{\cos(t)}{2}$$

Summary

The solution(s) found are the following

$$x = c_1 e^t - \frac{\sin(t)}{2} - \frac{\cos(t)}{2} \tag{1}$$




Figure 23: Slope field plot

Verification of solutions

$$x = c_1 e^t - \frac{\sin(t)}{2} - \frac{\cos(t)}{2}$$

Verified OK.

1.9.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$\begin{aligned} x' &= x + \sin(t) \\ x' &= \omega(t, x) \end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_t + \omega(\eta_x - \xi_t) - \omega^2 \xi_x - \omega_t \xi - \omega_x \eta = 0 \tag{A}$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 16: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(t, x) &= 0 \\ \eta(t, x) &= e^t\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(t, x) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dt}{\xi} = \frac{dx}{\eta} = dS\tag{1}$$

The above comes from the requirements that $(\xi \frac{\partial}{\partial t} + \eta \frac{\partial}{\partial x}) S(t, x) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = t$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^t} dy \end{aligned}$$

Which results in

$$S = e^{-t}x$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_t + \omega(t, x)S_x}{R_t + \omega(t, x)R_x} \quad (2)$$

Where in the above R_t, R_x, S_t, S_x are all partial derivatives and $\omega(t, x)$ is the right hand side of the original ode given by

$$\omega(t, x) = x + \sin(t)$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_t &= 1 \\ R_x &= 0 \\ S_t &= -e^{-t}x \\ S_x &= e^{-t} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = e^{-t} \sin(t) \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for t, x in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = e^{-R} \sin(R)$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 - \frac{e^{-R}(\cos(R) + \sin(R))}{2} \quad (4)$$

To complete the solution, we just need to transform (4) back to t, x coordinates. This results in

$$e^{-t}x = -\frac{(\cos(t) + \sin(t))e^{-t}}{2} + c_1$$



Which simplifies to

$$e^{-t}x = -\frac{(\cos(t) + \sin(t))e^{-t}}{2} + c_1$$

Which gives

$$x = -\frac{e^t(e^{-t}\sin(t) + e^{-t}\cos(t) - 2c_1)}{2}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in t, x coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dx}{dt} = x + \sin(t)$ 	$R = t$ $S = e^{-t}x$	$\frac{dS}{dR} = e^{-R} \sin(R)$ 

Summary

The solution(s) found are the following

$$x = -\frac{e^t(e^{-t}\sin(t) + e^{-t}\cos(t) - 2c_1)}{2} \quad (1)$$




Figure 24: Slope field plot

Verification of solutions

$$x = -\frac{e^t(e^{-t} \sin(t) + e^{-t} \cos(t) - 2c_1)}{2}$$

Verified OK.

1.9.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(t, x) dt + N(t, x) dx = 0 \tag{1A}$$

Therefore

$$\begin{aligned}dx &= (x + \sin(t)) dt \\ (-x - \sin(t)) dt + dx &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(t, x) &= -x - \sin(t) \\ N(t, x) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial x} = \frac{\partial N}{\partial t}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial x} &= \frac{\partial}{\partial x}(-x - \sin(t)) \\ &= -1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial t} &= \frac{\partial}{\partial t}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial x} \neq \frac{\partial N}{\partial t}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial x} - \frac{\partial N}{\partial t} \right) \\ &= 1((-1) - (0)) \\ &= -1 \end{aligned}$$

Since A does not depend on x , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dt} \\ &= e^{\int -1 dt} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-t} \\ &= e^{-t} \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \bar{M} &= \mu M \\ &= e^{-t}(-x - \sin(t)) \\ &= -e^{-t}(x + \sin(t)) \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= e^{-t}(1) \\ &= e^{-t} \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \bar{M} + \bar{N} \frac{dx}{dt} &= 0 \\ (-e^{-t}(x + \sin(t))) + (e^{-t}) \frac{dx}{dt} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(t, x)$

$$\frac{\partial \phi}{\partial t} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial x} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. t gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial t} dt &= \int \bar{M} dt \\ \int \frac{\partial \phi}{\partial t} dt &= \int -e^{-t}(x + \sin(t)) dt \\ \phi &= \frac{(2x + \cos(t) + \sin(t)) e^{-t}}{2} + f(x)\end{aligned}\quad (3)$$

Where $f(x)$ is used for the constant of integration since ϕ is a function of both t and x . Taking derivative of equation (3) w.r.t x gives

$$\frac{\partial \phi}{\partial x} = e^{-t} + f'(x) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial x} = e^{-t}$. Therefore equation (4) becomes

$$e^{-t} = e^{-t} + f'(x) \quad (5)$$

Solving equation (5) for $f'(x)$ gives

$$f'(x) = 0$$

Therefore

$$f(x) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(x)$ into equation (3) gives ϕ

$$\phi = \frac{(2x + \cos(t) + \sin(t)) e^{-t}}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{(2x + \cos(t) + \sin(t)) e^{-t}}{2}$$

The solution becomes

$$x = -\frac{e^t(e^{-t} \sin(t) + e^{-t} \cos(t) - 2c_1)}{2}$$

Summary

The solution(s) found are the following

$$x = -\frac{e^t(e^{-t} \sin(t) + e^{-t} \cos(t) - 2c_1)}{2} \quad (1)$$




Figure 25: Slope field plot

Verification of solutions

$$x = -\frac{e^t(e^{-t} \sin(t) + e^{-t} \cos(t) - 2c_1)}{2}$$

Verified OK.

1.9.4 Maple step by step solution

Let's solve

$$-x + x' = \sin(t)$$

- Highest derivative means the order of the ODE is 1
- x'
- Isolate the derivative

$$x' = x + \sin(t)$$

- Group terms with x on the lhs of the ODE and the rest on the rhs of the ODE

$$-x + x' = \sin(t)$$

- The ODE is linear; multiply by an integrating factor $\mu(t)$

$$\mu(t)(-x + x') = \mu(t)\sin(t)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dt}(\mu(t)x)$

$$\mu(t)(-x + x') = \mu'(t)x + \mu(t)x'$$

- Isolate $\mu'(t)$

$$\mu'(t) = -\mu(t)$$

- Solve to find the integrating factor

$$\mu(t) = e^{-t}$$

- Integrate both sides with respect to t

$$\int \left(\frac{d}{dt}(\mu(t)x) \right) dt = \int \mu(t)\sin(t) dt + c_1$$

- Evaluate the integral on the lhs

$$\mu(t)x = \int \mu(t)\sin(t) dt + c_1$$

- Solve for x

$$x = \frac{\int \mu(t)\sin(t)dt + c_1}{\mu(t)}$$

- Substitute $\mu(t) = e^{-t}$

$$x = \frac{\int e^{-t}\sin(t)dt + c_1}{e^{-t}}$$

- Evaluate the integrals on the rhs

$$x = \frac{-\frac{e^{-t}\sin(t)}{2} - \frac{e^{-t}\cos(t)}{2} + c_1}{e^{-t}}$$

- Simplify

$$x = c_1 e^t - \frac{\sin(t)}{2} - \frac{\cos(t)}{2}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 17

```
dsolve(diff(x(t),t)=x(t)+sin(t),x(t), singsol=all)
```

$$x(t) = -\frac{\cos(t)}{2} - \frac{\sin(t)}{2} + c_1 e^t$$

✓ Solution by Mathematica

Time used: 0.073 (sec). Leaf size: 24

```
DSolve[x'[t]==x[t]+Sin[t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow -\frac{\sin(t)}{2} - \frac{\cos(t)}{2} + c_1 e^t$$

1.10 problem Problem 10

- 1.10.1 Solving as first order ode lie symmetry calculated ode 113
- 1.10.2 Solving as exact ode 119

Internal problem ID [12121]

Internal file name [OUTPUT/10773_Tuesday_September_12_2023_08_51_42_AM_66891701/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 10.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exactByInspection", "first_order_ode_lie_symmetry_calculated"

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _dAlembert]
```

$$x(\ln(x) - \ln(y))y' - y = 0$$

1.10.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{y}{x(\ln(x) - \ln(y))}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \tag{1E}$$

$$\eta = xb_2 + yb_3 + b_1 \tag{2E}$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 + \frac{y(b_3 - a_2)}{x(\ln(x) - \ln(y))} - \frac{y^2 a_3}{x^2(\ln(x) - \ln(y))^2} \\ - \left(-\frac{y}{x^2(\ln(x) - \ln(y))} - \frac{y}{x^2(\ln(x) - \ln(y))^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(\frac{1}{x(\ln(x) - \ln(y))} + \frac{1}{x(\ln(x) - \ln(y))^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{\ln(x)^2 x^2 b_2 - 2 \ln(x) \ln(y) x^2 b_2 + \ln(y)^2 x^2 b_2 - \ln(x) x^2 b_2 + \ln(x) y^2 a_3 + \ln(y) x^2 b_2 - \ln(y) y^2 a_3 - \ln(x) x b_1 + \ln(x) y a_1 + \ln(y) x b_1 - \ln(y) y a_1 - b_2 x^2 + x y a_2 - x y b_3 - x b_1 + y a_1}{x^2 (\ln(x) - \ln(y))^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} \ln(x)^2 x^2 b_2 - 2 \ln(x) \ln(y) x^2 b_2 + \ln(y)^2 x^2 b_2 - \ln(x) x^2 b_2 \\ + \ln(x) y^2 a_3 + \ln(y) x^2 b_2 - \ln(y) y^2 a_3 - \ln(x) x b_1 + \ln(x) y a_1 \\ + \ln(y) x b_1 - \ln(y) y a_1 - b_2 x^2 + x y a_2 - x y b_3 - x b_1 + y a_1 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y, \ln(x), \ln(y)\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2, \ln(x) = v_3, \ln(y) = v_4\}$$

The above PDE (6E) now becomes

$$\begin{aligned} v_3^2 v_1^2 b_2 - 2 v_3 v_4 v_1^2 b_2 + v_4^2 v_1^2 b_2 + v_3 v_2^2 a_3 - v_4 v_2^2 a_3 - v_3 v_1^2 b_2 + v_4 v_1^2 b_2 + v_3 v_2 a_1 \\ - v_4 v_2 a_1 + v_1 v_2 a_2 - v_3 v_1 b_1 + v_4 v_1 b_1 - b_2 v_1^2 - v_1 v_2 b_3 + v_2 a_1 - v_1 b_1 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3, v_4\}$$

Equation (7E) now becomes

$$\begin{aligned} v_3^2 v_1^2 b_2 - 2v_3 v_4 v_1^2 b_2 - v_3 v_1^2 b_2 + v_4^2 v_1^2 b_2 + v_4 v_1^2 b_2 - b_2 v_1^2 + (-b_3 + a_2) v_1 v_2 & \quad (8E) \\ -v_3 v_1 b_1 + v_4 v_1 b_1 - v_1 b_1 + v_3 v_2^2 a_3 - v_4 v_2^2 a_3 + v_3 v_2 a_1 - v_4 v_2 a_1 + v_2 a_1 = 0 \end{aligned}$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} a_1 &= 0 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ -a_1 &= 0 \\ -a_3 &= 0 \\ -b_1 &= 0 \\ -2b_2 &= 0 \\ -b_2 &= 0 \\ -b_3 + a_2 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= b_3 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= x \\ \eta &= y \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}
 \eta &= \eta - \omega(x, y) \xi \\
 &= y - \left(\frac{y}{x(\ln(x) - \ln(y))} \right) (x) \\
 &= \frac{-y + y \ln(x) - \ln(y) y}{\ln(x) - \ln(y)} \\
 \xi &= 0
 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}
 S &= \int \frac{1}{\eta} dy \\
 &= \int \frac{1}{\frac{-y + y \ln(x) - \ln(y) y}{\ln(x) - \ln(y)}} dy
 \end{aligned}$$

Which results in

$$S = \ln(y) - \ln(-1 + \ln(x) - \ln(y))$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y) S_y}{R_x + \omega(x, y) R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y}{x(\ln(x) - \ln(y))}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{1}{x(-1 + \ln(x) - \ln(y))} \\ S_y &= \frac{1}{y} + \frac{1}{y(-1 + \ln(x) - \ln(y))} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\ln(y) - \ln(-1 + \ln(x) - \ln(y)) = c_1$$



Which simplifies to

$$\ln(y) - \ln(-1 + \ln(x) - \ln(y)) = c_1$$

Which gives

$$y = e^{-\text{LambertW}(e^{-1-c_1}x)-1}x$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y}{x(\ln(x) - \ln(y))}$ 	$R = x$ $S = \ln(y) - \ln(-1 + \ln(x))$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$y = e^{-\text{LambertW}(e^{-1-c_1 x}) - 1} x \tag{1}$$




Figure 26: Slope field plot

Verification of solutions

$$y = e^{-\text{LambertW}(e^{-1-c_1x})-1}x$$

Verified OK.

1.10.2 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned}(x(\ln(x) - \ln(y))) dy &= (y) dx \\ (-y) dx + (x(\ln(x) - \ln(y))) dy &= 0\end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -y \\ N(x, y) &= x(\ln(x) - \ln(y))\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-y) \\ &= -1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x(\ln(x) - \ln(y))) \\ &= \ln(x) - \ln(y) + 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. By inspection $\frac{1}{xy^2}$ is an integrating factor. Therefore by multiplying $M = -y$ and $N = x(\ln(x) - \ln(y))$ by this integrating factor the ode becomes exact. The new M, N are

$$M = -\frac{1}{yx}$$

$$N = \frac{\ln(x) - \ln(y)}{y^2}$$

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\frac{\partial \phi}{\partial x} = M$$

$$\frac{\partial \phi}{\partial y} = N$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} \left(\frac{\ln(x) - \ln(y)}{y^2}\right) dy &= \left(\frac{1}{xy}\right) dx \\ \left(-\frac{1}{xy}\right) dx + \left(\frac{\ln(x) - \ln(y)}{y^2}\right) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\frac{1}{xy} \\ N(x, y) &= \frac{\ln(x) - \ln(y)}{y^2} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{xy}\right) \\ &= \frac{1}{x y^2} \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{\ln(x) - \ln(y)}{y^2}\right) \\ &= \frac{1}{x y^2} \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{xy} dx \\ \phi &= -\frac{\ln(x)}{y} + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \frac{\ln(x)}{y^2} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{\ln(x) - \ln(y)}{y^2}$. Therefore equation (4) becomes

$$\frac{\ln(x) - \ln(y)}{y^2} = \frac{\ln(x)}{y^2} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -\frac{\ln(y)}{y^2}$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int \left(-\frac{\ln(y)}{y^2} \right) dy$$

$$f(y) = \frac{\ln(y)}{y} + \frac{1}{y} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{\ln(x)}{y} + \frac{\ln(y)}{y} + \frac{1}{y} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{\ln(x)}{y} + \frac{\ln(y)}{y} + \frac{1}{y}$$

The solution becomes

$$y = -\frac{\text{LambertW}(-c_1 e^{-1}x)}{c_1}$$

Summary

The solution(s) found are the following

$$y = -\frac{\text{LambertW}(-c_1 e^{-1}x)}{c_1} \quad (1)$$




Figure 27: Slope field plot

Verification of solutions

$$y = -\frac{\text{LambertW}(-c_1 e^{-1}x)}{c_1}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous G  
1st order, trying the canonical coordinates of the invariance group  
<- 1st order, canonical coordinates successful  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 14

```
dsolve(x*(ln(x)-ln(y(x)))*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\text{LambertW}(c_1 x e^{-1})}{c_1}$$

✓ Solution by Mathematica

Time used: 7.587 (sec). Leaf size: 37

```
DSolve[x*(Log[x]-Log[y[x]])*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned}y(x) &\rightarrow -e^{c_1} W(-e^{-1-c_1} x) \\y(x) &\rightarrow 0 \\y(x) &\rightarrow \frac{x}{e}\end{aligned}$$

1.11 problem Problem 11

1.11.1 Maple step by step solution 128

Internal problem ID [12122]

Internal file name [OUTPUT/10774_Tuesday_September_12_2023_08_51_44_AM_24737299/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 11.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "differentialType", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$xyy'^2 - (x^2 + y^2)y' + yx = 0$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{y}{x} \tag{1}$$

$$y' = \frac{x}{y} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{y}{x} \end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= \frac{1}{x} dx \\ \int \frac{1}{y} dy &= \int \frac{1}{x} dx \\ \ln(y) &= \ln(x) + c_1 \\ y &= e^{\ln(x)+c_1} \\ &= c_1 x\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x \tag{1}$$

Verification of solutions

$$y = c_1 x$$

Verified OK.

Solving equation (2)

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{x}{y}\end{aligned}$$

Where $f(x) = x$ and $g(y) = \frac{1}{y}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= x dx \\ \int \frac{1}{y} dy &= \int x dx \\ \frac{y^2}{2} &= \frac{x^2}{2} + c_2\end{aligned}$$

Which results in

$$\begin{aligned}y &= \sqrt{x^2 + 2c_2} \\ y &= -\sqrt{x^2 + 2c_2}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \sqrt{x^2 + 2c_2} \quad (1)$$

$$y = -\sqrt{x^2 + 2c_2} \quad (2)$$

Verification of solutions

$$y = \sqrt{x^2 + 2c_2}$$

Verified OK.

$$y = -\sqrt{x^2 + 2c_2}$$

Verified OK.

1.11.1 Maple step by step solution

Let's solve

$$xyy'^2 - (x^2 + y^2)y' + yx = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{y} = \frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int \frac{1}{x} dx + c_1$$

- Evaluate integral

$$\ln(y) = \ln(x) + c_1$$

- Solve for y

$$y = e^{c_1} x$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful  
Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 29

```
dsolve(x*y(x)*diff(y(x),x)^2-(x^2+y(x)^2)*diff(y(x),x)+x*y(x)=0,y(x), singsol=all)
```

$$\begin{aligned}y(x) &= c_1 x \\ y(x) &= \sqrt{x^2 + c_1} \\ y(x) &= -\sqrt{x^2 + c_1}\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.17 (sec). Leaf size: 55

```
DSolve[x*y[x]*y'[x]^2-(x^2+y[x]^2)*y'[x]+x*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned}y(x) &\rightarrow c_1 x \\ y(x) &\rightarrow -\sqrt{x^2 + 2c_1} \\ y(x) &\rightarrow \sqrt{x^2 + 2c_1} \\ y(x) &\rightarrow -x \\ y(x) &\rightarrow x\end{aligned}$$

1.12 problem Problem 12

1.12.1 Maple step by step solution 131

Internal problem ID [12123]

Internal file name [OUTPUT/10775_Tuesday_September_12_2023_08_51_44_AM_49389396/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 12.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y'^2 - 9y^4 = 0$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = -3y^2 \tag{1}$$

$$y' = 3y^2 \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int -\frac{1}{3y^2} dy = x + c_1$$
$$\frac{1}{3y} = x + c_1$$

Solving for y gives these solutions

$$y_1 = \frac{1}{3c_1 + 3x}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{3c_1 + 3x} \quad (1)$$

Verification of solutions

$$y = \frac{1}{3c_1 + 3x}$$

Verified OK.

Solving equation (2)

Integrating both sides gives

$$\int \frac{1}{3y^2} dy = x + c_2$$
$$-\frac{1}{3y} = x + c_2$$

Solving for y gives these solutions

$$y_1 = -\frac{1}{3(x + c_2)}$$

Summary

The solution(s) found are the following

$$y = -\frac{1}{3(x + c_2)} \quad (1)$$

Verification of solutions

$$y = -\frac{1}{3(x + c_2)}$$

Verified OK.

1.12.1 Maple step by step solution

Let's solve

$$y'^2 - 9y^4 = 0$$

- Highest derivative means the order of the ODE is 1
- y'
- Separate variables

$$\frac{y'}{y^2} = 3$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y^2} dx = \int 3 dx + c_1$$

- Evaluate integral

$$-\frac{1}{y} = 3x + c_1$$

- Solve for y

$$y = -\frac{1}{3x+c_1}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful  
Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)^2=9*y(x)^4,y(x), singsol=all)
```

$$y(x) = \frac{1}{c_1 - 3x}$$

$$y(x) = \frac{1}{3x + c_1}$$

✓ Solution by Mathematica

Time used: 0.263 (sec). Leaf size: 34

```
DSolve[y'[x]^2==9*y[x]^4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{1}{3x + c_1}$$

$$y(x) \rightarrow \frac{1}{3x - c_1}$$

$$y(x) \rightarrow 0$$

1.13 problem Problem 13

1.13.1 Solving as homogeneousTypeD ode	134
1.13.2 Solving as homogeneousTypeD2 ode	136
1.13.3 Solving as first order ode lie symmetry lookup ode	138

Internal problem ID [12124]

Internal file name [OUTPUT/10776_Tuesday_September_12_2023_08_51_44_AM_80572458/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 13.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"homogeneousTypeD", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _dAlembert]
```

$$x' - e^{\frac{x}{t}} - \frac{x}{t} = 0$$

1.13.1 Solving as homogeneousTypeD ode

Writing the ode as

$$x' = e^{\frac{x}{t}} + \frac{x}{t} \tag{A}$$

The given ode has the form

$$y' = \frac{y}{x} + g(x) f\left(b\frac{y}{x}\right)^{\frac{n}{m}} \tag{1}$$

Where b is scalar and $g(x)$ is function of x and n, m are integers. The solution is given in Kamke page 20. Using the substitution $y(x) = u(x)x$ then

$$\frac{dy}{dx} = \frac{du}{dx}x + u$$

Hence the given ode becomes

$$\begin{aligned}\frac{du}{dx}x + u &= u + g(x) f(bu)^{\frac{n}{m}} \\ u' &= \frac{1}{x}g(x) f(bu)^{\frac{n}{m}}\end{aligned}\quad (2)$$

The above ode is always separable. This is easily solved for u assuming the integration can be resolved, and then the solution to the original ode becomes $y = ux$. Comparing the given ode (A) with the form (1) shows that

$$\begin{aligned}g(t) &= 1 \\ b &= 1 \\ f\left(\frac{bt}{x}\right) &= e^{\frac{x}{t}}\end{aligned}$$

Substituting the above in (2) results in the $u(t)$ ode as

$$u'(t) = \frac{e^{u(t)}}{t}$$

Which is now solved as separable In canonical form the ODE is

$$\begin{aligned}u' &= F(t, u) \\ &= f(t)g(u) \\ &= \frac{e^u}{t}\end{aligned}$$

Where $f(t) = \frac{1}{t}$ and $g(u) = e^u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{e^u} du &= \frac{1}{t} dt \\ \int \frac{1}{e^u} du &= \int \frac{1}{t} dt \\ -e^{-u} &= \ln(t) + c_1\end{aligned}$$

The solution is

$$-e^{-u(t)} - \ln(t) - c_1 = 0$$

Therefore the solution is found using $x = ut$. Hence

$$-e^{-\frac{x}{t}} - \ln(t) - c_1 = 0$$

Summary

The solution(s) found are the following

$$-e^{-\frac{x}{t}} - \ln(t) - c_1 = 0 \quad (1)$$




Figure 28: Slope field plot

Verification of solutions

$$-e^{-\frac{x}{t}} - \ln(t) - c_1 = 0$$

Verified OK.

1.13.2 Solving as homogeneous Type D2 ode

Using the change of variables $x = u(t)t$ on the above ode results in new ode in $u(t)$

$$u'(t)t - e^{u(t)} = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(t, u) \\ &= f(t)g(u) \\ &= \frac{e^u}{t} \end{aligned}$$

Where $f(t) = \frac{1}{t}$ and $g(u) = e^u$. Integrating both sides gives

$$\begin{aligned}\frac{1}{e^u} du &= \frac{1}{t} dt \\ \int \frac{1}{e^u} du &= \int \frac{1}{t} dt \\ -e^{-u} &= \ln(t) + c_2\end{aligned}$$

The solution is

$$-e^{-u(t)} - \ln(t) - c_2 = 0$$

Replacing $u(t)$ in the above solution by $\frac{x}{t}$ results in the solution for x in implicit form

$$\begin{aligned}-e^{-\frac{x}{t}} - \ln(t) - c_2 &= 0 \\ -e^{-\frac{x}{t}} - \ln(t) - c_2 &= 0\end{aligned}$$

Summary

The solution(s) found are the following

$$-e^{-\frac{x}{t}} - \ln(t) - c_2 = 0 \tag{1}$$




Figure 29: Slope field plot

Verification of solutions

$$-e^{-\frac{x}{t}} - \ln(t) - c_2 = 0$$

Verified OK.

1.13.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$x' = \frac{e^{\frac{x}{t}}t + x}{t}$$
$$x' = \omega(t, x)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_t + \omega(\eta_x - \xi_t) - \omega^2\xi_x - \omega_t\xi - \omega_x\eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **homogeneous Type D**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 21: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(t, x) &= t^2 \\ \eta(t, x) &= tx\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(t, x) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dt}{\xi} = \frac{dx}{\eta} = dS\tag{1}$$

The above comes from the requirements that $(\xi \frac{\partial}{\partial t} + \eta \frac{\partial}{\partial x})S(t, x) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Therefore

$$\begin{aligned}\frac{dx}{dt} &= \frac{\eta}{\xi} \\ &= \frac{tx}{t^2} \\ &= \frac{x}{t}\end{aligned}$$

This is easily solved to give

$$x = c_1 t$$

Where now the coordinate R is taken as the constant of integration. Hence

$$R = \frac{x}{t}$$

And S is found from

$$\begin{aligned}dS &= \frac{dt}{\xi} \\ &= \frac{dt}{t^2}\end{aligned}$$

Integrating gives

$$\begin{aligned}S &= \int \frac{dt}{T} \\ &= -\frac{1}{t}\end{aligned}$$

Where the constant of integration is set to zero as we just need one solution. Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_t + \omega(t, x)S_x}{R_t + \omega(t, x)R_x} \quad (2)$$

Where in the above R_t, R_x, S_t, S_x are all partial derivatives and $\omega(t, x)$ is the right hand side of the original ode given by

$$\omega(t, x) = \frac{e^{\frac{x}{t}} t + x}{t}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_t &= -\frac{x}{t^2} \\ R_x &= \frac{1}{t} \\ S_t &= \frac{1}{t^2} \\ S_x &= 0\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{e^{-\frac{x}{t}}}{t} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for t, x in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -S(R) e^{-R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 e^{e^{-R}} \quad (4)$$

To complete the solution, we just need to transform (4) back to t, x coordinates. This results in

$$-\frac{1}{t} = c_1 e^{-\frac{x}{t}}$$



Which simplifies to

$$-\frac{1}{t} = c_1 e^{-\frac{x}{t}}$$

Which gives

$$x = -\ln \left(\ln \left(-\frac{1}{c_1 t} \right) \right) t$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in t, x coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dx}{dt} = \frac{e^{\frac{x}{t}} t + x}{t}$ 	$R = \frac{x}{t}$ $S = -\frac{1}{t}$	$\frac{dS}{dR} = -S(R) e^{-R}$ 

Summary

The solution(s) found are the following

$$x = -\ln \left(\ln \left(-\frac{1}{c_1 t} \right) \right) t \tag{1}$$




Figure 30: Slope field plot

Verification of solutions

$$x = -\ln\left(\ln\left(-\frac{1}{c_1 t}\right)\right) t$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous D
<- homogeneous successful`

```


✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 15

```
dsolve(diff(x(t),t)=exp(x(t)/t)+x(t)/t,x(t), singsol=all)
```

$$x(t) = \ln\left(-\frac{1}{\ln(t) + c_1}\right) t$$

✓ Solution by Mathematica

Time used: 0.54 (sec). Leaf size: 18

```
DSolve[x'[t]==Exp[x[t]/t]+x[t]/t,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow -t \log(-\log(t) - c_1)$$

1.14 problem Problem 14

1.14.1 Maple step by step solution 146

Internal problem ID [12125]

Internal file name [OUTPUT/10777_Tuesday_September_12_2023_08_51_46_AM_91798504/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 14.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "quadrature"

Maple gives the following as the ode type

[_quadrature]

$$y'^2 = -x^2 + 1$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{-x^2 + 1} \tag{1}$$

$$y' = -\sqrt{-x^2 + 1} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned} y &= \int \sqrt{-x^2 + 1} \, dx \\ &= \frac{\sqrt{-x^2 + 1} x}{2} + \frac{\arcsin(x)}{2} + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{\sqrt{-x^2 + 1} x}{2} + \frac{\arcsin(x)}{2} + c_1 \tag{1}$$

Verification of solutions

$$y = \frac{\sqrt{-x^2 + 1} x}{2} + \frac{\arcsin(x)}{2} + c_1$$

Verified OK.

Solving equation (2)

Integrating both sides gives

$$\begin{aligned} y &= \int -\sqrt{-x^2 + 1} dx \\ &= -\frac{\sqrt{-x^2 + 1} x}{2} - \frac{\arcsin(x)}{2} + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = -\frac{\sqrt{-x^2 + 1} x}{2} - \frac{\arcsin(x)}{2} + c_2 \quad (1)$$

Verification of solutions

$$y = -\frac{\sqrt{-x^2 + 1} x}{2} - \frac{\arcsin(x)}{2} + c_2$$

Verified OK.

1.14.1 Maple step by step solution

Let's solve

$$y'^2 = -x^2 + 1$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int y'^2 dx = \int (-x^2 + 1) dx + c_1$$

- Cannot compute integral

$$\int y'^2 dx = x - \frac{1}{3}x^3 + c_1$$

Maple trace

```
`Methods for first order ODEs:  
-> Solving 1st order ODE of high degree, 1st attempt  
trying 1st order WeierstrassP solution for high degree ODE  
trying 1st order WeierstrassPPrime solution for high degree ODE  
trying 1st order JacobiSN solution for high degree ODE  
trying 1st order ODE linearizable_by_differentiation  
trying differential order: 1; missing variables  
<- differential order: 1; missing y(x) successful`
```

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 43

```
dsolve(x^2+diff(y(x),x)^2=1,y(x), singsol=all)
```

$$y(x) = \frac{x\sqrt{-x^2+1}}{2} + \frac{\arcsin(x)}{2} + c_1$$
$$y(x) = -\frac{x\sqrt{-x^2+1}}{2} - \frac{\arcsin(x)}{2} + c_1$$

✓ Solution by Mathematica

Time used: 0.07 (sec). Leaf size: 85

```
DSolve[x^2+y'[x]^2==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\arctan\left(\frac{\sqrt{1-x^2}}{x+1}\right) + \frac{1}{2}\sqrt{1-x^2}x + c_1$$
$$y(x) \rightarrow \arctan\left(\frac{\sqrt{1-x^2}}{x+1}\right) - \frac{1}{2}\sqrt{1-x^2}x + c_1$$

1.15 problem Problem 15

1.15.1 Solving as separable ode	148
1.15.2 Solving as homogeneousTypeD2 ode	150
1.15.3 Solving as first order ode lie symmetry lookup ode	152
1.15.4 Solving as bernoulli ode	156
1.15.5 Solving as exact ode	159
1.15.6 Maple step by step solution	163

Internal problem ID [12126]

Internal file name [OUTPUT/10778_Tuesday_September_12_2023_08_51_46_AM_73155439/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 15.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "bernoulli", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y - y'x - \frac{1}{y} = 0$$

1.15.1 Solving as separable ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{y^2 - 1}{xy}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(y) = \frac{y^2-1}{y}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{y^2-1}{y}} dy &= \frac{1}{x} dx \\ \int \frac{1}{\frac{y^2-1}{y}} dy &= \int \frac{1}{x} dx \\ \frac{\ln(y-1)}{2} + \frac{\ln(y+1)}{2} &= \ln(x) + c_1\end{aligned}$$

The above can be written as

$$\begin{aligned}\left(\frac{1}{2}\right) (\ln(y-1) + \ln(y+1)) &= \ln(x) + 2c_1 \\ \ln(y-1) + \ln(y+1) &= (2) (\ln(x) + 2c_1) \\ &= 2\ln(x) + 4c_1\end{aligned}$$

Raising both side to exponential gives

$$e^{\ln(y-1)+\ln(y+1)} = e^{2\ln(x)+2c_1}$$

Which simplifies to

$$\begin{aligned}y^2 - 1 &= 2c_1x^2 \\ &= c_2x^2\end{aligned}$$

The solution is

$$y^2 - 1 = c_2x^2$$

Summary

The solution(s) found are the following

$$y^2 - 1 = c_2x^2 \tag{1}$$




Figure 31: Slope field plot

Verification of solutions

$$y^2 - 1 = c_2 x^2$$

Verified OK.

1.15.2 Solving as homogeneous Type D2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u(x)x - (u'(x)x + u(x))x - \frac{1}{u(x)x} = 0$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{1}{u x^3} \end{aligned}$$

Where $f(x) = -\frac{1}{x^3}$ and $g(u) = \frac{1}{u}$. Integrating both sides gives

$$\frac{1}{u} du = -\frac{1}{x^3} dx$$

$$\int \frac{1}{\frac{1}{u}} du = \int -\frac{1}{x^3} dx$$

$$\frac{u^2}{2} = \frac{1}{2x^2} + c_2$$

The solution is

$$\frac{u(x)^2}{2} - \frac{1}{2x^2} - c_2 = 0$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$\frac{y^2}{2x^2} - \frac{1}{2x^2} - c_2 = 0$$

$$\frac{y^2}{2x^2} - \frac{1}{2x^2} - c_2 = 0$$

Summary

The solution(s) found are the following

$$\frac{y^2}{2x^2} - \frac{1}{2x^2} - c_2 = 0 \tag{1}$$




Figure 32: Slope field plot

Verification of solutions

$$\frac{y^2}{2x^2} - \frac{1}{2x^2} - c_2 = 0$$

Verified OK.

1.15.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{y^2 - 1}{xy}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **separable**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 24: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= x \\ \eta(x, y) &= 0\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\eta = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\xi} dx \\ &= \int \frac{1}{x} dx \end{aligned}$$

Which results in

$$S = \ln(x)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y^2 - 1}{xy}$$

Evaluating all the partial derivatives gives

$$R_x = 0$$

$$R_y = 1$$

$$S_x = \frac{1}{x}$$

$$S_y = 0$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{y}{y^2 - 1} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{R}{R^2 - 1}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{\ln(R-1)}{2} + \frac{\ln(R+1)}{2} + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\ln(x) = \frac{\ln(y-1)}{2} + \frac{\ln(y+1)}{2} + c_1$$

Which simplifies to

$$\ln(x) = \frac{\ln(y-1)}{2} + \frac{\ln(y+1)}{2} + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y^2-1}{xy}$ 	$R = y$ $S = \ln(x)$	$\frac{dS}{dR} = \frac{R}{R^2-1}$ 

Summary

The solution(s) found are the following

$$\ln(x) = \frac{\ln(y-1)}{2} + \frac{\ln(y+1)}{2} + c_1 \quad (1)$$




Figure 33: Slope field plot

Verification of solutions

$$\ln(x) = \frac{\ln(y-1)}{2} + \frac{\ln(y+1)}{2} + c_1$$

Verified OK.

1.15.4 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{y^2 - 1}{xy} \end{aligned}$$

This is a Bernoulli ODE.

$$y' = \frac{1}{x}y - \frac{1}{x}\frac{1}{y} \tag{1}$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \tag{2}$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \tag{3}$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned} f_0(x) &= \frac{1}{x} \\ f_1(x) &= -\frac{1}{x} \\ n &= -1 \end{aligned}$$

Dividing both sides of ODE (1) by $y^n = \frac{1}{y}$ gives

$$y'y = \frac{y^2}{x} - \frac{1}{x} \quad (4)$$

Let

$$\begin{aligned} w &= y^{1-n} \\ &= y^2 \end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = 2yy' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} \frac{w'(x)}{2} &= \frac{w(x)}{x} - \frac{1}{x} \\ w' &= \frac{2w}{x} - \frac{2}{x} \end{aligned} \quad (7)$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$\begin{aligned} p(x) &= -\frac{2}{x} \\ q(x) &= -\frac{2}{x} \end{aligned}$$

Hence the ode is

$$w'(x) - \frac{2w(x)}{x} = -\frac{2}{x}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\frac{2}{x} dx} \\ &= \frac{1}{x^2}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu w) &= (\mu) \left(-\frac{2}{x}\right) \\ \frac{d}{dx} \left(\frac{w}{x^2}\right) &= \left(\frac{1}{x^2}\right) \left(-\frac{2}{x}\right) \\ d\left(\frac{w}{x^2}\right) &= \left(-\frac{2}{x^3}\right) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}\frac{w}{x^2} &= \int -\frac{2}{x^3} dx \\ \frac{w}{x^2} &= \frac{1}{x^2} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x^2}$ results in

$$w(x) = c_1 x^2 + 1$$

Replacing w in the above by y^2 using equation (5) gives the final solution.

$$y^2 = c_1 x^2 + 1$$

Solving for y gives

$$\begin{aligned}y(x) &= \sqrt{c_1 x^2 + 1} \\ y(x) &= -\sqrt{c_1 x^2 + 1}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \sqrt{c_1 x^2 + 1} \tag{1}$$

$$y = -\sqrt{c_1 x^2 + 1} \tag{2}$$




Figure 34: Slope field plot

Verification of solutions

$$y = \sqrt{c_1 x^2 + 1}$$

Verified OK.

$$y = -\sqrt{c_1 x^2 + 1}$$

Verified OK.

1.15.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} \left(\frac{y}{y^2 - 1}\right) dy &= \left(\frac{1}{x}\right) dx \\ \left(-\frac{1}{x}\right) dx + \left(\frac{y}{y^2 - 1}\right) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -\frac{1}{x} \\ N(x, y) &= \frac{y}{y^2 - 1} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(-\frac{1}{x}\right) \\ &= 0 \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{y}{y^2 - 1} \right) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{x} dx \\ \phi &= -\ln(x) + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 0 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{y}{y^2 - 1}$. Therefore equation (4) becomes

$$\frac{y}{y^2 - 1} = 0 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{y}{y^2 - 1}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{y}{y^2 - 1} \right) dy \\ f(y) &= \frac{\ln(y - 1)}{2} + \frac{\ln(y + 1)}{2} + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x) + \frac{\ln(y-1)}{2} + \frac{\ln(y+1)}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x) + \frac{\ln(y-1)}{2} + \frac{\ln(y+1)}{2}$$

Summary

The solution(s) found are the following

$$-\ln(x) + \frac{\ln(y-1)}{2} + \frac{\ln(y+1)}{2} = c_1 \quad (1)$$




Figure 35: Slope field plot

Verification of solutions

$$-\ln(x) + \frac{\ln(y-1)}{2} + \frac{\ln(y+1)}{2} = c_1$$

Verified OK.

1.15.6 Maple step by step solution

Let's solve

$$y - y'x - \frac{1}{y} = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'}{-y + \frac{1}{y}} = -\frac{1}{x}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{-y + \frac{1}{y}} dx = \int -\frac{1}{x} dx + c_1$$

- Evaluate integral

$$-\frac{\ln(y-1)}{2} - \frac{\ln(y+1)}{2} = -\ln(x) + c_1$$

- Solve for y

$$\left\{ y = \frac{(e^{c_1})^2 - \sqrt{(e^{c_1})^4 + (e^{c_1})^2 x^2 + x^2}}{(e^{c_1})^2 - \sqrt{(e^{c_1})^4 + (e^{c_1})^2 x^2}}, y = \frac{(e^{c_1})^2 + \sqrt{(e^{c_1})^4 + (e^{c_1})^2 x^2 + x^2}}{(e^{c_1})^2 + \sqrt{(e^{c_1})^4 + (e^{c_1})^2 x^2}} \right\}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 27

```
dsolve(y(x)=x*diff(y(x),x)+1/y(x),y(x), singsol=all)
```

$$y(x) = \sqrt{c_1 x^2 + 1}$$
$$y(x) = -\sqrt{c_1 x^2 + 1}$$

✓ Solution by Mathematica

Time used: 0.43 (sec). Leaf size: 53

```
DSolve[y[x]==x*y'[x]+1/y[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\sqrt{1 + e^{2c_1}x^2}$$

$$y(x) \rightarrow \sqrt{1 + e^{2c_1}x^2}$$

$$y(x) \rightarrow -1$$

$$y(x) \rightarrow 1$$

1.16 problem Problem 16

1.16.1 Maple step by step solution 168

Internal problem ID [12127]

Internal file name [OUTPUT/10779_Tuesday_September_12_2023_08_51_48_AM_22632786/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 16.

ODE order: 1.

ODE degree: 3.

The type(s) of ODE detected by this program : **"quadrature"**

Maple gives the following as the ode type

[_quadrature]

$$\boxed{-y'^3 + y' = -x + 2}$$

Solving the given ode for y' results in 3 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}}{6} + \frac{2}{(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} \quad (1)$$

$$y' = -\frac{(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}}{12} - \frac{1}{(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} + \frac{i\sqrt{3}}{(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} \quad (2)$$

$$y' = -\frac{(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}}{12} - \frac{1}{(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} - \frac{i\sqrt{3}}{(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} \quad (3)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{2}{3}} + 12}{6(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} dx \\ &= \int \frac{(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{2}{3}} + 12}{6(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} dx + c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \int \frac{(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{2}{3}} + 12}{6(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} dx + c_1 \quad (1)$$

Verification of solutions

$$y = \int \frac{(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{2}{3}} + 12}{6(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} dx + c_1$$

Verified OK.

Solving equation (2)

Integrating both sides gives

$$\begin{aligned} y &= \int \frac{i(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{2}{3}} \sqrt{3} - 12i\sqrt{3} - (-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}}{12(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} dx \\ &= \int \frac{i(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{2}{3}} \sqrt{3} - 12i\sqrt{3} - (-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}}{12(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} dx \end{aligned}$$

Summary

The solution(s) found are the following

$$\begin{aligned} y &= \int \frac{i(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{2}{3}} \sqrt{3} - 12i\sqrt{3} - (-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}}{12(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} dx + c_2 \end{aligned} \quad (1)$$

Verification of solutions

$$y = \int \frac{i(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{2}{3}} \sqrt{3} - 12i\sqrt{3} - (-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}}{12(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} dx + c_2$$

Verified OK.

Solving equation (3)

Integrating both sides gives

$$y = \int -\frac{i(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{2}{3}} \sqrt{3} - 12i\sqrt{3} + (-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}}{12(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} dx + c_3$$
$$= \int -\frac{i(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{2}{3}} \sqrt{3} - 12i\sqrt{3} + (-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}}{12(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} dx + c_3$$

Summary

The solution(s) found are the following

$$y = \int \frac{i(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{2}{3}} \sqrt{3} - 12i\sqrt{3} + (-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}}{12(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} dx + c_3 \quad (1)$$

Verification of solutions

$$y = \int \frac{i(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{2}{3}} \sqrt{3} - 12i\sqrt{3} + (-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}}{12(-216 + 108x + 12\sqrt{81x^2 - 324x + 312})^{\frac{1}{3}}} dx + c_3$$

Verified OK.

1.16.1 Maple step by step solution

Let's solve

$$-y'^3 + y' = -x + 2$$

- Highest derivative means the order of the ODE is 1

y'

- Integrate both sides with respect to x

$$\int (-y'^3 + y') dx = \int (-x + 2) dx + c_1$$

- Cannot compute integral

$$\int (-y'^3 + y') dx = -\frac{1}{2}x^2 + 2x + c_1$$

Maple trace

```
`Methods for first order ODEs:  
-> Solving 1st order ODE of high degree, 1st attempt  
trying 1st order WeierstrassP solution for high degree ODE  
trying 1st order WeierstrassPPrime solution for high degree ODE  
trying 1st order JacobiSN solution for high degree ODE  
trying 1st order ODE linearizable_by_differentiation  
trying differential order: 1; missing variables  
<- differential order: 1; missing y(x) successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 211

```
dsolve(x=diff(y(x),x)^3-diff(y(x),x)+2,y(x), singsol=all)
```

$$y(x) = -\frac{\left(\int \frac{i\sqrt{3}(-216+108x+12\sqrt{81x^2-324x+312})^{\frac{2}{3}}-12i\sqrt{3}+(-216+108x+12\sqrt{81x^2-324x+312})^{\frac{2}{3}}+12}{(-216+108x+12\sqrt{81x^2-324x+312})^{\frac{1}{3}}} dx\right)}{12} + c_1$$
$$y(x) = \frac{\left(\int \frac{(i\sqrt{3}-1)(-216+108x+12\sqrt{81x^2-324x+312})^{\frac{2}{3}}-12i\sqrt{3}-12}{(-216+108x+12\sqrt{81x^2-324x+312})^{\frac{1}{3}}} dx\right)}{12} + c_1$$
$$y(x) = \frac{\left(\int \frac{(-216+108x+12\sqrt{81x^2-324x+312})^{\frac{2}{3}}+12}{(-216+108x+12\sqrt{81x^2-324x+312})^{\frac{1}{3}}} dx\right)}{6} + c_1$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[x==y'[x]^3-y'[x]+2,y[x],x,IncludeSingularSolutions -> True]
```

Timed out

1.17 problem Problem 17

- 1.17.1 Solving as first order ode lie symmetry calculated ode 170
- 1.17.2 Solving as exact ode 175

Internal problem ID [12128]

Internal file name [OUTPUT/10780_Tuesday_September_12_2023_08_51_49_AM_54326988/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 17.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exactWithIntegrationFactor", "first_order_ode_lie_symmetry_calculated"**

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _rational]
```

$$y' - \frac{y}{x + y^3} = 0$$

1.17.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{y}{y^3 + x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \tag{1E}$$

$$\eta = xb_2 + yb_3 + b_1 \tag{2E}$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 + \frac{y(b_3 - a_2)}{y^3 + x} - \frac{y^2 a_3}{(y^3 + x)^2} + \frac{y(xa_2 + ya_3 + a_1)}{(y^3 + x)^2} - \left(\frac{1}{y^3 + x} - \frac{3y^3}{(y^3 + x)^2} \right) (xb_2 + yb_3 + b_1) = 0 \quad (5E)$$

Putting the above in normal form gives

$$\frac{y^6 b_2 + 4x y^3 b_2 - y^4 a_2 + 3y^4 b_3 + 2y^3 b_1 - xb_1 + ya_1}{(y^3 + x)^2} = 0$$

Setting the numerator to zero gives

$$y^6 b_2 + 4x y^3 b_2 - y^4 a_2 + 3y^4 b_3 + 2y^3 b_1 - xb_1 + ya_1 = 0 \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$b_2 v_2^6 - a_2 v_2^4 + 4b_2 v_1 v_2^3 + 3b_3 v_2^4 + 2b_1 v_2^3 + a_1 v_2 - b_1 v_1 = 0 \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$4b_2 v_1 v_2^3 - b_1 v_1 + b_2 v_2^6 + (-a_2 + 3b_3) v_2^4 + 2b_1 v_2^3 + a_1 v_2 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} a_1 &= 0 \\ b_2 &= 0 \\ -b_1 &= 0 \\ 2b_1 &= 0 \\ 4b_2 &= 0 \\ -a_2 + 3b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= 3b_3 \\ a_3 &= a_3 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the ansatz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= y \\ \eta &= 0 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= 0 - \left(\frac{y}{y^3 + x} \right) (y) \\ &= -\frac{y^2}{y^3 + x} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{-\frac{y^2}{y^3+x}} dy \end{aligned}$$

Which results in

$$S = -\frac{y^2}{2} + \frac{x}{y}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y}{y^3 + x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{1}{y} \\ S_y &= -y - \frac{x}{y^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{y^2}{2} + \frac{x}{y} = c_1$$

Which simplifies to

$$-\frac{y^2}{2} + \frac{x}{y} = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y}{y^3+x}$ 	$R = x$ $S = -\frac{y^2}{2} + \frac{x}{y}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$-\frac{y^2}{2} + \frac{x}{y} = c_1 \tag{1}$$




Figure 36: Slope field plot

Verification of solutions

$$-\frac{y^2}{2} + \frac{x}{y} = c_1$$

Verified OK.

1.17.2 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(y^3 + x) dy &= (y) dx \\ (-y) dx + (y^3 + x) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= -y \\ N(x, y) &= y^3 + x\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-y) \\ &= -1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(y^3 + x) \\ &= 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{y^3 + x} ((-1) - (1)) \\ &= -\frac{2}{y^3 + x} \end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned} B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= -\frac{1}{y} ((1) - (-1)) \\ &= -\frac{2}{y} \end{aligned}$$

Since B does not depend on x , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned} \mu &= e^{\int B \, dy} \\ &= e^{\int -\frac{2}{y} \, dy} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-2 \ln(y)} \\ &= \frac{1}{y^2} \end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \overline{M} and \overline{N} so not to confuse them with the original M and N .

$$\begin{aligned} \overline{M} &= \mu M \\ &= \frac{1}{y^2} (-y) \\ &= -\frac{1}{y} \end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{1}{y^2}(y^3 + x) \\ &= \frac{y^3 + x}{y^2}\end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(-\frac{1}{y}\right) + \left(\frac{y^3 + x}{y^2}\right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int -\frac{1}{y} dx \\ \phi &= -\frac{x}{y} + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \frac{x}{y^2} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{y^3+x}{y^2}$. Therefore equation (4) becomes

$$\frac{y^3 + x}{y^2} = \frac{x}{y^2} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = y$$

Integrating the above w.r.t y gives

$$\int f'(y) \, dy = \int (y) \, dy$$

$$f(y) = \frac{y^2}{2} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x}{y} + \frac{y^2}{2} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x}{y} + \frac{y^2}{2}$$

Summary

The solution(s) found are the following

$$\frac{y^2}{2} - \frac{x}{y} = c_1 \tag{1}$$




Figure 37: Slope field plot

Verification of solutions

$$\frac{y^2}{2} - \frac{x}{y} = c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
<- 1st order linear successful
<- inverse linear successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 224

```
dsolve(diff(y(x),x)=y(x)/(x+y(x)^3),y(x), singsol=all)
```

$$y(x) = \frac{\left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{2}{3}} - 6c_1}{3\left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{1}{3}}}$$

$$y(x) = -\frac{i\sqrt{3}\left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{2}{3}} + 6i\sqrt{3}c_1 + \left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{2}{3}} - 6c_1}{6\left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{1}{3}}}$$

$$y(x) = \frac{i\sqrt{3}\left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{2}{3}} + 6i\sqrt{3}c_1 - \left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{2}{3}} + 6c_1}{6\left(27x + 3\sqrt{24c_1^3 + 81x^2}\right)^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 2.895 (sec). Leaf size: 263

```
DSolve[y'[x]==y[x]/(x+y[x]^3),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2 \cdot 3^{2/3} c_1 - \sqrt[3]{3} (-9x + \sqrt{81x^2 + 24c_1^3})^{2/3}}{3 \sqrt[3]{-9x + \sqrt{81x^2 + 24c_1^3}}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{3} (1 - i\sqrt{3}) (-9x + \sqrt{81x^2 + 24c_1^3})^{2/3} - 2 \sqrt[6]{3} (\sqrt{3} + 3i) c_1}{6 \sqrt[3]{-9x + \sqrt{81x^2 + 24c_1^3}}}$$

$$y(x) \rightarrow \frac{\sqrt[3]{3} (1 + i\sqrt{3}) (-9x + \sqrt{81x^2 + 24c_1^3})^{2/3} - 2 \sqrt[6]{3} (\sqrt{3} - 3i) c_1}{6 \sqrt[3]{-9x + \sqrt{81x^2 + 24c_1^3}}}$$

$$y(x) \rightarrow 0$$

1.18 problem Problem 18

1.18.1 Solving as quadrature ode	182
1.18.2 Maple step by step solution	183

Internal problem ID [12129]

Internal file name [OUTPUT/10781_Tuesday_September_12_2023_08_51_50_AM_51096026/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 18.

ODE order: 1.

ODE degree: 4.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y - y'^4 + y'^3 = -2$$

1.18.1 Solving as quadrature ode

Integrating both sides gives

$$\int \frac{1}{\text{RootOf}(_Z^4 - _Z^3 - y - 2)} dy = \int dx$$
$$\int^y \frac{1}{\text{RootOf}(_Z^4 - _Z^3 - _a - 2)} d_a = x + c_1$$

Summary

The solution(s) found are the following

$$\int^y \frac{1}{\text{RootOf}(_Z^4 - _Z^3 - _a - 2)} d_a = x + c_1 \quad (1)$$

Verification of solutions

$$\int^y \frac{1}{\text{RootOf}(_Z^4 - _Z^3 - _a - 2)} d_a = x + c_1$$

Verified OK.

1.18.2 Maple step by step solution

Let's solve

$$y - y'^4 + y'^3 = -2$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'}{\text{RootOf}(-Z^4 - Z^3 - y - 2)} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\text{RootOf}(-Z^4 - Z^3 - y - 2)} dx = \int 1 dx + c_1$$

- Evaluate integral

$$-\frac{3\text{RootOf}(-Z^4 - Z^3 - y - 2)^2}{2} + \frac{4\text{RootOf}(-Z^4 - Z^3 - y - 2)^3}{3} = x + c_1$$

- Solve for y

$$y = -\frac{3c_1 \left(\frac{(27+192c_1+192x+24\sqrt{64c_1^2+128c_1x+64x^2+18c_1+18x})^{\frac{1}{3}}}{8} + \frac{9}{8(27+192c_1+192x+24\sqrt{64c_1^2+128c_1x+64x^2+18c_1+18x})^{\frac{1}{3}} + \frac{3}{8}} \right)}{4} + \dots$$

Maple trace

```
`Methods for first order ODEs:
```

```
*** Sublevel 2 ***
```

```
Methods for first order ODEs:
```

```
-> Solving 1st order ODE of high degree, 1st attempt
```

```
trying 1st order WeierstrassP solution for high degree ODE
```

```
trying 1st order WeierstrassPPrime solution for high degree ODE
```

```
trying 1st order JacobiSN solution for high degree ODE
```

```
trying 1st order ODE linearizable_by_differentiation
```

```
trying differential order: 1; missing variables
```

```
<- differential order: 1; missing x successful`
```


✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 247

```
dsolve(y(x)=diff(y(x),x)^4-diff(y(x),x)^3-2,y(x), singsol=all)
```

$$y(x) = -2$$

$$y(x)$$

$$= \frac{12 \left(\frac{243}{16384} + \frac{(\frac{9}{64} - c_1 + x)\sqrt{64}\sqrt{(x - c_1 + \frac{9}{32})(x - c_1)}}{16} + \frac{c_1^2}{2} + (-\frac{9}{64} - x)c_1 + \frac{x^2}{2} + \frac{9x}{64} \right) (27 - 192c_1 + 192x + 24\sqrt{64})}{\dots}$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y[x]==y'[x]^4-y'[x]^3-2,y[x],x,IncludeSingularSolutions -> True]
```

Timed out

1.19 problem Problem 26

1.19.1 Maple step by step solution 186

Internal problem ID [12130]

Internal file name [OUTPUT/10782_Tuesday_September_12_2023_08_51_52_AM_99008589/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 26.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y'^2 + y^2 = 4$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{4 - y^2} \tag{1}$$

$$y' = -\sqrt{4 - y^2} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{-y^2 + 4}} dy = x + c_1$$
$$\arcsin\left(\frac{y}{2}\right) = x + c_1$$

Solving for y gives these solutions

$$y_1 = 2 \sin(x + c_1)$$

Summary

The solution(s) found are the following

$$y = 2 \sin(x + c_1) \quad (1)$$

Verification of solutions

$$y = 2 \sin(x + c_1)$$

Verified OK.

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{-y^2 + 4}} dy = x + c_2$$
$$-\arcsin\left(\frac{y}{2}\right) = x + c_2$$

Solving for y gives these solutions

$$y_1 = -2 \sin(x + c_2)$$

Summary

The solution(s) found are the following

$$y = -2 \sin(x + c_2) \quad (1)$$

Verification of solutions

$$y = -2 \sin(x + c_2)$$

Verified OK.

1.19.1 Maple step by step solution

Let's solve

$$y'^2 + y^2 = 4$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Separate variables

$$\frac{y'}{\sqrt{4-y^2}} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\sqrt{4-y^2}} dx = \int 1 dx + c_1$$

- Evaluate integral
 $\arcsin\left(\frac{y}{2}\right) = x + c_1$
- Solve for y
 $y = 2 \sin(x + c_1)$

Maple trace

```

`Methods for first order ODEs:
*** Sublevel 2 ***
Methods for first order ODEs:
-> Solving 1st order ODE of high degree, 1st attempt
trying 1st order WeierstrassP solution for high degree ODE
trying 1st order WeierstrassPPrime solution for high degree ODE
trying 1st order JacobiSN solution for high degree ODE
trying 1st order ODE linearizable_by_differentiation
trying differential order: 1; missing variables
<- differential order: 1; missing x successful`

```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 31

```
dsolve(diff(y(x),x)^2+y(x)^2=4,y(x), singsol=all)
```

$$y(x) = -2$$

$$y(x) = 2$$

$$y(x) = -2 \sin(c_1 - x)$$

$$y(x) = 2 \sin(c_1 - x)$$

✓ Solution by Mathematica

Time used: 0.306 (sec). Leaf size: 43

```
DSolve[y'[x]^2+y[x]^2==4,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 2 \cos(x + c_1)$$

$$y(x) \rightarrow 2 \cos(x - c_1)$$

$$y(x) \rightarrow -2$$

$$y(x) \rightarrow 2$$

$$y(x) \rightarrow \text{Interval}[\{-2, 2\}]$$

1.20 problem Problem 28

1.20.1 Solving as homogeneousTypeMapleC ode 189

1.20.2 Solving as first order ode lie symmetry calculated ode 193

Internal problem ID [12131]

Internal file name [OUTPUT/10783_Tuesday_September_12_2023_08_51_52_AM_58373890/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 28.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"homogeneousTypeMapleC", "first_order_ode_lie_symmetry_calculated"**

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$y' - \frac{2y - x - 4}{2x - y + 5} = 0$$

1.20.1 Solving as homogeneousTypeMapleC ode

Let $Y = y + y_0$ and $X = x + x_0$ then the above is transformed to new ode in $Y(X)$

$$\frac{d}{dX}Y(X) = -\frac{2Y(X) + 2y_0 - X - x_0 - 4}{-2X - 2x_0 + Y(X) + y_0 - 5}$$

Solving for possible values of x_0 and y_0 which makes the above ode a homogeneous ode results in

$$x_0 = -2$$

$$y_0 = 1$$

Using these values now it is possible to easily solve for $Y(X)$. The above ode now becomes

$$\frac{d}{dX}Y(X) = -\frac{2Y(X) - X}{-2X + Y(X)}$$

In canonical form, the ODE is

$$\begin{aligned} Y' &= F(X, Y) \\ &= -\frac{2Y - X}{-2X + Y} \end{aligned} \quad (1)$$

An ode of the form $Y' = \frac{M(X, Y)}{N(X, Y)}$ is called homogeneous if the functions $M(X, Y)$ and $N(X, Y)$ are both homogeneous functions and of the same order. Recall that a function $f(X, Y)$ is homogeneous of order n if

$$f(t^n X, t^n Y) = t^n f(X, Y)$$

In this case, it can be seen that both $M = 2Y - X$ and $N = 2X - Y$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{Y}{X}$, or $Y = uX$. Hence

$$\frac{dY}{dX} = \frac{du}{dX}X + u$$

Applying the transformation $Y = uX$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dX}X + u &= \frac{-2u + 1}{u - 2} \\ \frac{du}{dX} &= \frac{\frac{-2u(X)+1}{u(X)-2} - u(X)}{X} \end{aligned}$$

Or

$$\frac{d}{dX}u(X) - \frac{\frac{-2u(X)+1}{u(X)-2} - u(X)}{X} = 0$$

Or

$$\left(\frac{d}{dX}u(X)\right)Xu(X) - 2\left(\frac{d}{dX}u(X)\right)X + u(X)^2 - 1 = 0$$

Or

$$X(u(X) - 2)\left(\frac{d}{dX}u(X)\right) + u(X)^2 - 1 = 0$$

Which is now solved as separable in $u(X)$. Which is now solved in $u(X)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(X, u) \\ &= f(X)g(u) \\ &= -\frac{u^2 - 1}{X(u - 2)} \end{aligned}$$

Where $f(X) = -\frac{1}{X}$ and $g(u) = \frac{u^2-1}{u-2}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{u^2-1}{u-2}} du &= -\frac{1}{X} dX \\ \int \frac{1}{\frac{u^2-1}{u-2}} du &= \int -\frac{1}{X} dX \\ -\frac{\ln(u-1)}{2} + \frac{3\ln(u+1)}{2} &= -\ln(X) + c_2\end{aligned}$$

The above can be written as

$$\begin{aligned}\frac{-\ln(u-1) + 3\ln(u+1)}{2} &= -\ln(X) + c_2 \\ -\ln(u-1) + 3\ln(u+1) &= (2)(-\ln(X) + c_2) \\ &= -2\ln(X) + 2c_2\end{aligned}$$

Raising both side to exponential gives

$$e^{-\ln(u-1)+3\ln(u+1)} = e^{-2\ln(X)+2c_2}$$

Which simplifies to

$$\begin{aligned}\frac{(u+1)^3}{u-1} &= \frac{2c_2}{X^2} \\ &= \frac{c_3}{X^2}\end{aligned}$$

Which simplifies to

$$\frac{(u(X)+1)^3}{u(X)-1} = \frac{c_3 e^{2c_2}}{X^2}$$

The solution is

$$\frac{(u(X)+1)^3}{u(X)-1} = \frac{c_3 e^{2c_2}}{X^2}$$

Now u in the above solution is replaced back by Y using $u = \frac{Y}{X}$ which results in the solution

$$\frac{\left(\frac{Y(X)}{X} + 1\right)^3}{\frac{Y(X)}{X} - 1} = \frac{c_3 e^{2c_2}}{X^2}$$

Which simplifies to

$$-\frac{(Y(X) + X)^3}{-Y(X) + X} = c_3 e^{2c_2}$$

Using the solution for $Y(X)$

$$-\frac{(Y(X) + X)^3}{-Y(X) + X} = c_3 e^{2c_2}$$

And replacing back terms in the above solution using

$$Y = y + y_0$$

$$X = x + x_0$$

Or

$$Y = y + 1$$

$$X = x - 2$$

Then the solution in y becomes

$$-\frac{(x + y + 1)^3}{-y + 3 + x} = c_3 e^{2c_2}$$

Summary

The solution(s) found are the following

$$-\frac{(x + y + 1)^3}{-y + 3 + x} = c_3 e^{2c_2} \quad (1)$$




Figure 38: Slope field plot

Verification of solutions

$$-\frac{(x+y+1)^3}{-y+3+x} = c_3 e^{2c_2}$$

Verified OK.

1.20.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = -\frac{2y-x-4}{-2x+y-5}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \tag{1E}$$

$$\eta = xb_2 + yb_3 + b_1 \tag{2E}$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 - \frac{(2y-x-4)(b_3-a_2)}{-2x+y-5} - \frac{(2y-x-4)^2 a_3}{(-2x+y-5)^2} \\ - \left(\frac{1}{-2x+y-5} - \frac{2(2y-x-4)}{(-2x+y-5)^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(-\frac{2}{-2x+y-5} + \frac{2y-x-4}{(-2x+y-5)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{2x^2a_2 - x^2a_3 + x^2b_2 - 2x^2b_3 - 2xya_2 + 4xya_3 - 4xyb_2 + 2xyb_3 + 2y^2a_2 - y^2a_3 + y^2b_2 - 2y^2b_3 + 10xa_2 - 8xa_3 - 3xb_1 + 14xb_2 - 13xb_3 + 3ya_1 - 14ya_2 + 13ya_3 - 10yb_2 + 8yb_3 - 3a_1 + 20a_2 - 16a_3 - 6b_1 + 25b_2 - 20b_3}{(2x+y-5)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} 2x^2a_2 - x^2a_3 + x^2b_2 - 2x^2b_3 - 2xya_2 + 4xya_3 - 4xyb_2 + 2xyb_3 + 2y^2a_2 \\ - y^2a_3 + y^2b_2 - 2y^2b_3 + 10xa_2 - 8xa_3 - 3xb_1 + 14xb_2 - 13xb_3 + 3ya_1 \\ - 14ya_2 + 13ya_3 - 10yb_2 + 8yb_3 - 3a_1 + 20a_2 - 16a_3 - 6b_1 + 25b_2 - 20b_3 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} 2a_2v_1^2 - 2a_2v_1v_2 + 2a_2v_2^2 - a_3v_1^2 + 4a_3v_1v_2 - a_3v_2^2 + b_2v_1^2 - 4b_2v_1v_2 + b_2v_2^2 \\ - 2b_3v_1^2 + 2b_3v_1v_2 - 2b_3v_2^2 + 3a_1v_2 + 10a_2v_1 - 14a_2v_2 - 8a_3v_1 + 13a_3v_2 - 3b_1v_1 \\ + 14b_2v_1 - 10b_2v_2 - 13b_3v_1 + 8b_3v_2 - 3a_1 + 20a_2 - 16a_3 - 6b_1 + 25b_2 - 20b_3 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned} & (2a_2 - a_3 + b_2 - 2b_3) v_1^2 + (-2a_2 + 4a_3 - 4b_2 + 2b_3) v_1 v_2 \\ & + (10a_2 - 8a_3 - 3b_1 + 14b_2 - 13b_3) v_1 + (2a_2 - a_3 + b_2 - 2b_3) v_2^2 \\ & + (3a_1 - 14a_2 + 13a_3 - 10b_2 + 8b_3) v_2 - 3a_1 + 20a_2 - 16a_3 - 6b_1 + 25b_2 - 20b_3 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} -2a_2 + 4a_3 - 4b_2 + 2b_3 &= 0 \\ 2a_2 - a_3 + b_2 - 2b_3 &= 0 \\ 3a_1 - 14a_2 + 13a_3 - 10b_2 + 8b_3 &= 0 \\ 10a_2 - 8a_3 - 3b_1 + 14b_2 - 13b_3 &= 0 \\ -3a_1 + 20a_2 - 16a_3 - 6b_1 + 25b_2 - 20b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= -b_2 + 2b_3 \\ a_2 &= b_3 \\ a_3 &= b_2 \\ b_1 &= 2b_2 - b_3 \\ b_2 &= b_2 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= x + 2 \\ \eta &= y - 1 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= y - 1 - \left(-\frac{2y - x - 4}{-2x + y - 5} \right) (x + 2) \\ &= \frac{x^2 - y^2 + 4x + 2y + 3}{2x - y + 5} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{x^2 - y^2 + 4x + 2y + 3}{2x - y + 5}} dy \end{aligned}$$

Which results in

$$S = \frac{3 \ln(x + y + 1)}{2} - \frac{\ln(y - 3 - x)}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{2y - x - 4}{-2x + y - 5}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{-2y + x + 4}{(x + y + 1)(x + 3 - y)} \\ S_y &= \frac{2x - y + 5}{(x + y + 1)(x + 3 - y)} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{3 \ln(x + y + 1)}{2} - \frac{\ln(y - 3 - x)}{2} = c_1$$

Which simplifies to

$$\frac{3 \ln(x + y + 1)}{2} - \frac{\ln(y - 3 - x)}{2} = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{2y-x-4}{-2x+y-5}$ 	$R = x$ $S = \frac{3 \ln(x + y + 1)}{2} - \frac{\ln(y - 3 - x)}{2}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$\frac{3 \ln(x + y + 1)}{2} - \frac{\ln(y - 3 - x)}{2} = c_1 \quad (1)$$




Figure 39: Slope field plot

Verification of solutions

$$\frac{3 \ln(x + y + 1)}{2} - \frac{\ln(y - 3 - x)}{2} = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous C  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.891 (sec). Leaf size: 117

```
dsolve(diff(y(x),x)=(2*y(x)-x-4)/(2*x-y(x)+5),y(x), singsol=all)
```

$y(x) =$

$$\frac{(i\sqrt{3} - 1) \left(3\sqrt{3} \sqrt{27c_1^2 (x+2)^2 - 1} + 27c_1(x+2) \right)^{\frac{2}{3}} - 3i\sqrt{3} - 3 + 6 \left(3\sqrt{3} \sqrt{27c_1^2 (x+2)^2 - 1} + 27c_1(x+2) \right)^{\frac{1}{3}}}{6 \left(3\sqrt{3} \sqrt{27c_1^2 (x+2)^2 - 1} + 27c_1(x+2) \right)^{\frac{1}{3}} c_1}$$

✓ Solution by Mathematica

Time used: 60.277 (sec). Leaf size: 1624

```
DSolve[y'[x]==(2*y[x]-x-4)/(2*x-y[x]+5),y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

1.21 problem Problem 29

1.21.1 Solving as first order ode lie symmetry lookup ode	200
1.21.2 Solving as bernoulli ode	204
1.21.3 Solving as exact ode	208
1.21.4 Solving as riccati ode	212

Internal problem ID [12132]

Internal file name [OUTPUT/10784_Tuesday_September_12_2023_08_51_55_AM_65763563/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 29.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**riccati**", "**bernoulli**", "**exactWithIntegrationFactor**", "**first_order_ode_lie_symmetry_lookup**"

Maple gives the following as the ode type

```
[[_1st_order, _with_linear_symmetries], _rational, _Bernoulli]
```

$$y' - \frac{y}{x+1} + y^2 = 0$$

1.21.1 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{y(xy + y - 1)}{x + 1}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is known. It is of type **Bernoulli**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 30: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{y^2}{x+1}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right)S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{y^2}{x+1}} dy \end{aligned}$$

Which results in

$$S = -\frac{x+1}{y}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{y(xy + y - 1)}{x + 1}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{1}{y} \\ S_y &= \frac{x+1}{y^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -x - 1 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -R - 1$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\frac{1}{2}R^2 - R + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{-x-1}{y} = -\frac{1}{2}x^2 - x + c_1$$



Which simplifies to

$$\frac{-x-1}{y} = -\frac{1}{2}x^2 - x + c_1$$

Which gives

$$y = -\frac{2(x+1)}{-x^2 + 2c_1 - 2x}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{y(xy+y-1)}{x+1}$ 	$R = x$ $S = \frac{-x-1}{y}$	$\frac{dS}{dR} = -R - 1$ 

Summary

The solution(s) found are the following

$$y = -\frac{2(x+1)}{-x^2 + 2c_1 - 2x} \quad (1)$$




Figure 40: Slope field plot

Verification of solutions

$$y = -\frac{2(x+1)}{-x^2 + 2c_1 - 2x}$$

Verified OK.

1.21.2 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= -\frac{y(xy + y - 1)}{x + 1} \end{aligned}$$

This is a Bernoulli ODE.

$$y' = \frac{1}{x+1}y - y^2 \quad (1)$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \quad (2)$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \quad (3)$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned} f_0(x) &= \frac{1}{x+1} \\ f_1(x) &= -1 \\ n &= 2 \end{aligned}$$

Dividing both sides of ODE (1) by $y^n = y^2$ gives

$$y' \frac{1}{y^2} = \frac{1}{(x+1)y} - 1 \quad (4)$$

Let

$$\begin{aligned} w &= y^{1-n} \\ &= \frac{1}{y} \end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = -\frac{1}{y^2}y' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} -w'(x) &= \frac{w(x)}{x+1} - 1 \\ w' &= -\frac{w}{x+1} + 1 \end{aligned} \quad (7)$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$p(x) = \frac{1}{x+1}$$
$$q(x) = 1$$

Hence the ode is

$$w'(x) + \frac{w(x)}{x+1} = 1$$

The integrating factor μ is

$$\mu = e^{\int \frac{1}{x+1} dx}$$
$$= x + 1$$

The ode becomes

$$\frac{d}{dx}(\mu w) = \mu$$
$$\frac{d}{dx}((x+1)w) = x+1$$
$$d((x+1)w) = (x+1)dx$$

Integrating gives

$$(x+1)w = \int (x+1) dx$$
$$(x+1)w = \frac{1}{2}x^2 + x + c_1$$

Dividing both sides by the integrating factor $\mu = x+1$ results in

$$w(x) = \frac{\frac{1}{2}x^2 + x}{x+1} + \frac{c_1}{x+1}$$

which simplifies to

$$w(x) = \frac{x^2 + 2c_1 + 2x}{2x + 2}$$

Replacing w in the above by $\frac{1}{y}$ using equation (5) gives the final solution.

$$\frac{1}{y} = \frac{x^2 + 2c_1 + 2x}{2x + 2}$$

Or

$$y = \frac{2x + 2}{x^2 + 2c_1 + 2x}$$

Summary

The solution(s) found are the following

$$y = \frac{2x + 2}{x^2 + 2c_1 + 2x} \tag{1}$$




Figure 41: Slope field plot

Verification of solutions

$$y = \frac{2x + 2}{x^2 + 2c_1 + 2x}$$

Verified OK.

1.21.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (x+1) dy &= (-y(xy+y-1)) dx \\ (y(xy+y-1)) dx + (x+1) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= y(xy+y-1) \\ N(x, y) &= x+1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(y(xy + y - 1)) \\ &= -1 + (2x + 2)y\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x + 1) \\ &= 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{x + 1} ((xy + y - 1 + y(x + 1)) - (1)) \\ &= \frac{2xy + 2y - 2}{x + 1}\end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned}B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= \frac{1}{y(xy + y - 1)} ((1) - (xy + y - 1 + y(x + 1))) \\ &= -\frac{2}{y}\end{aligned}$$

Since B does not depend on x , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned}\mu &= e^{\int B \, dy} \\ &= e^{\int -\frac{2}{y} \, dy}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{-2\ln(y)} \\ &= \frac{1}{y^2}\end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \overline{M} and \overline{N} so not to confuse them with the original M and N .

$$\begin{aligned}\overline{M} &= \mu M \\ &= \frac{1}{y^2}(y(xy + y - 1)) \\ &= \frac{xy + y - 1}{y}\end{aligned}$$

And

$$\begin{aligned}\overline{N} &= \mu N \\ &= \frac{1}{y^2}(x + 1) \\ &= \frac{x + 1}{y^2}\end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned}\overline{M} + \overline{N} \frac{dy}{dx} &= 0 \\ \left(\frac{xy + y - 1}{y} \right) + \left(\frac{x + 1}{y^2} \right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{xy + y - 1}{y} dx \\ \phi &= \frac{(-2 + (x + 2)y)x}{2y} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned}\frac{\partial\phi}{\partial y} &= \frac{(x+2)x}{2y} - \frac{(-2+(x+2)y)x}{2y^2} + f'(y) \\ &= \frac{x}{y^2} + f'(y)\end{aligned}\tag{4}$$

But equation (2) says that $\frac{\partial\phi}{\partial y} = \frac{x+1}{y^2}$. Therefore equation (4) becomes

$$\frac{x+1}{y^2} = \frac{x}{y^2} + f'(y)\tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = \frac{1}{y^2}$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int \left(\frac{1}{y^2}\right) dy \\ f(y) &= -\frac{1}{y} + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{(-2+(x+2)y)x}{2y} - \frac{1}{y} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{(-2+(x+2)y)x}{2y} - \frac{1}{y}$$

The solution becomes

$$y = -\frac{2(x+1)}{-x^2 + 2c_1 - 2x}$$

Summary

The solution(s) found are the following

$$y = -\frac{2(x+1)}{-x^2 + 2c_1 - 2x} \quad (1)$$




Figure 42: Slope field plot

Verification of solutions

$$y = -\frac{2(x+1)}{-x^2 + 2c_1 - 2x}$$

Verified OK.

1.21.4 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= -\frac{y(xy + y - 1)}{x + 1} \end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = -\frac{y^2 x}{x+1} - \frac{y^2}{x+1} + \frac{y}{x+1}$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = 0$, $f_1(x) = \frac{1}{x+1}$ and $f_2(x) = -1$. Let

$$\begin{aligned} y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{-u} \end{aligned} \tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \tag{2}$$

But

$$\begin{aligned} f_2' &= 0 \\ f_1 f_2 &= -\frac{1}{x+1} \\ f_2^2 f_0 &= 0 \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$-u''(x) + \frac{u'(x)}{x+1} = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = c_1 + c_2(x+1)^2$$

The above shows that

$$u'(x) = 2(x+1)c_2$$

Using the above in (1) gives the solution

$$y = \frac{2(x+1)c_2}{c_1 + c_2(x+1)^2}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = \frac{2x + 2}{x^2 + c_3 + 2x + 1}$$

Summary

The solution(s) found are the following

$$y = \frac{2x + 2}{x^2 + c_3 + 2x + 1} \tag{1}$$




Figure 43: Slope field plot

Verification of solutions

$$y = \frac{2x + 2}{x^2 + c_3 + 2x + 1}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 22

```
dsolve(diff(y(x),x)-y(x)/(1+x)+y(x)^2=0,y(x), singsol=all)
```

$$y(x) = \frac{2 + 2x}{x^2 + 2c_1 + 2x}$$

✓ Solution by Mathematica

Time used: 0.297 (sec). Leaf size: 28

```
DSolve[y'[x]-y[x]/(1+x)+y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{2(x+1)}{x^2 + 2x + 2c_1}$$
$$y(x) \rightarrow 0$$

1.22 problem Problem 30

1.22.1 Existence and uniqueness analysis	216
1.22.2 Solving as riccati ode	217

Internal problem ID [12133]

Internal file name [OUTPUT/10785_Tuesday_September_12_2023_08_51_56_AM_4946144/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 30.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**riccati**"

Maple gives the following as the ode type

```
[[_Riccati, _special]]
```

$$y' - y^2 = x$$

With initial conditions

$$[y(0) = 0]$$

1.22.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned}y' &= f(x, y) \\ &= y^2 + x\end{aligned}$$

The x domain of $f(x, y)$ when $y = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The y domain of $f(x, y)$ when $x = 0$ is

$$\{-\infty < y < \infty\}$$

And the point $y_0 = 0$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial y} &= \frac{\partial}{\partial y}(y^2 + x) \\ &= 2y\end{aligned}$$

The y domain of $\frac{\partial f}{\partial y}$ when $x = 0$ is

$$\{-\infty < y < \infty\}$$

And the point $y_0 = 0$ is inside this domain. Therefore solution exists and is unique.

1.22.2 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= y^2 + x\end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = y^2 + x$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = x$, $f_1(x) = 0$ and $f_2(x) = 1$. Let

$$\begin{aligned}y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{u}\end{aligned}\tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0\tag{2}$$

But

$$\begin{aligned}f_2' &= 0 \\ f_1 f_2 &= 0 \\ f_2^2 f_0 &= x\end{aligned}$$

Substituting the above terms back in equation (2) gives

$$u''(x) + xu(x) = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = c_1 \text{AiryAi}(-x) + c_2 \text{AiryBi}(-x)$$

The above shows that

$$u'(x) = -c_1 \text{AiryAi}(1, -x) - c_2 \text{AiryBi}(1, -x)$$

Using the above in (1) gives the solution

$$y = -\frac{-c_1 \text{AiryAi}(1, -x) - c_2 \text{AiryBi}(1, -x)}{c_1 \text{AiryAi}(-x) + c_2 \text{AiryBi}(-x)}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = \frac{c_3 \text{AiryAi}(1, -x) + \text{AiryBi}(1, -x)}{c_3 \text{AiryAi}(-x) + \text{AiryBi}(-x)}$$

Initial conditions are used to solve for c_3 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = \frac{3\Gamma\left(\frac{2}{3}\right)^2 3^{\frac{2}{3}} - 3\Gamma\left(\frac{2}{3}\right)^2 c_3 3^{\frac{1}{6}}}{2 \cdot 3^{\frac{5}{6}} \pi + 2\pi c_3 3^{\frac{1}{3}}}$$

$$c_3 = \sqrt{3}$$


Substituting c_3 found above in the general solution gives

$$y = \frac{\text{AiryAi}(1, -x) \sqrt{3} + \text{AiryBi}(1, -x)}{\text{AiryAi}(-x) \sqrt{3} + \text{AiryBi}(-x)}$$


Summary

The solution(s) found are the following

$$y = \frac{\text{AiryAi}(1, -x) \sqrt{3} + \text{AiryBi}(1, -x)}{\text{AiryAi}(-x) \sqrt{3} + \text{AiryBi}(-x)} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{\text{AiryAi}(1, -x) \sqrt{3} + \text{AiryBi}(1, -x)}{\text{AiryAi}(-x) \sqrt{3} + \text{AiryBi}(-x)}$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
differential order: 1; looking for linear symmetries
trying exact
Looking for potential symmetries
trying Riccati
trying Riccati Special
<- Riccati Special successful`

```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 35

```
dsolve([diff(y(x),x)=x+y(x)^2,y(0) = 0],y(x), singsol=all)
```

$$y(x) = \frac{\sqrt{3} \operatorname{AiryAi}(1, -x) + \operatorname{AiryBi}(1, -x)}{\sqrt{3} \operatorname{AiryAi}(-x) + \operatorname{AiryBi}(-x)}$$

✓ Solution by Mathematica

Time used: 1.869 (sec). Leaf size: 80

```
DSolve[{y'[x]==x+y[x]^2,{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x^{3/2} \operatorname{BesselJ}\left(-\frac{4}{3}, \frac{2x^{3/2}}{3}\right) - x^{3/2} \operatorname{BesselJ}\left(\frac{2}{3}, \frac{2x^{3/2}}{3}\right) + \operatorname{BesselJ}\left(-\frac{1}{3}, \frac{2x^{3/2}}{3}\right)}{2x \operatorname{BesselJ}\left(-\frac{1}{3}, \frac{2x^{3/2}}{3}\right)}$$

1.23 problem Problem 31

1.23.1 Existence and uniqueness analysis	221
1.23.2 Solving as <code>abelFirstKind</code> ode	222

Internal problem ID [12134]

Internal file name [OUTPUT/10786_Tuesday_September_12_2023_08_52_06_AM_59708118/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 31.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**abelFirstKind**"

Maple gives the following as the ode type

`[_Abe1]`

Unable to solve or complete the solution.

$$y' - y^3 x = x^2$$

With initial conditions

$$[y(0) = 0]$$

1.23.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned}y' &= f(x, y) \\ &= x y^3 + x^2\end{aligned}$$

The x domain of $f(x, y)$ when $y = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The y domain of $f(x, y)$ when $x = 0$ is

$$\{-\infty < y < \infty\}$$

And the point $y_0 = 0$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial y} &= \frac{\partial}{\partial y}(x y^3 + x^2) \\ &= 3x y^2\end{aligned}$$

The x domain of $\frac{\partial f}{\partial y}$ when $y = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The y domain of $\frac{\partial f}{\partial y}$ when $x = 0$ is

$$\{-\infty < y < \infty\}$$

And the point $y_0 = 0$ is inside this domain. Therefore solution exists and is unique.

1.23.2 Solving as AbelFirstKind ode

This is Abel first kind ODE, it has the form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2 + f_3(x)y^3$$

Comparing the above to given ODE which is

$$y' = y^3 x + x^2 \tag{1}$$

Therefore

$$\begin{aligned}f_0(x) &= x^2 \\ f_1(x) &= 0 \\ f_2(x) &= 0 \\ f_3(x) &= x\end{aligned}$$

Since $f_2(x) = 0$ then we check the Abel invariant to see if it depends on x or not. The Abel invariant is given by

$$-\frac{f_1^3}{f_0^2 f_3}$$

Which when evaluating gives

$$\frac{1}{27x^8}$$

Since the Abel invariant depends on x then unable to solve this ode at this time.

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
differential order: 1; looking for linear symmetries
trying exact
trying Abel
Looking for potential symmetries
Looking for potential symmetries
Looking for potential symmetries
trying inverse_Riccati
trying an equivalence to an Abel ODE
differential order: 1; trying a linearization to 2nd order
--- trying a change of variables {x -> y(x), y(x) -> x}
differential order: 1; trying a linearization to 2nd order
trying 1st order ODE linearizable_by_differentiation
--- Trying Lie symmetry methods, 1st order ---
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = 4
`, `-> Computing symmetries using: way = 2
trying symmetry patterns for 1st order ODEs
-> trying a symmetry pattern of the form [F(x)*G(y), 0]
-> trying a symmetry pattern of the form [0, F(x)*G(y)]
-> trying symmetry patterns of the forms [F(x),G(y)] and [G(y),F(x)]
-> trying a symmetry pattern of the form [F(x),G(x)]
-> trying a symmetry pattern of the form [F(y),G(y)]
-> trying a symmetry pattern of the form [F(x)+G(y), 0]
-> trying a symmetry pattern of the form [0, F(x)+G(y)]
-> trying a symmetry pattern of the form [F(x),G(x)*y+H(x)]
-> trying a symmetry pattern of conformal type`
```


X Solution by Maple

```
dsolve([diff(y(x),x)=x*y(x)^3+x^2,y(0) = 0],y(x), singsol=all)
```

No solution found

X Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[{y'[x]==x*y[x]^3+x^2,{y[0]==0}},y[x],x,IncludeSingularSolutions -> True]
```

Not solved

1.24 problem Problem 35

1.24.1 Solving as riccati ode 225

Internal problem ID [12135]

Internal file name [OUTPUT/10787_Tuesday_September_12_2023_08_52_06_AM_79625629/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 35.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**riccati**"

Maple gives the following as the ode type

[_Riccati]

$$y' + y^2 = x^2$$

1.24.1 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= x^2 - y^2\end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = x^2 - y^2$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = x^2$, $f_1(x) = 0$ and $f_2(x) = -1$. Let

$$\begin{aligned}y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{-u}\end{aligned}\tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \quad (2)$$

But

$$\begin{aligned} f_2' &= 0 \\ f_1 f_2 &= 0 \\ f_2^2 f_0 &= x^2 \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$-u''(x) + x^2 u(x) = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = \left(\text{BesselI} \left(\frac{1}{4}, \frac{x^2}{2} \right) c_1 + \text{BesselK} \left(\frac{1}{4}, \frac{x^2}{2} \right) c_2 \right) \sqrt{x}$$

The above shows that

$$u'(x) = x^{\frac{3}{2}} \left(\text{BesselI} \left(-\frac{3}{4}, \frac{x^2}{2} \right) c_1 - \text{BesselK} \left(\frac{3}{4}, \frac{x^2}{2} \right) c_2 \right)$$

Using the above in (1) gives the solution

$$y = \frac{x \left(\text{BesselI} \left(-\frac{3}{4}, \frac{x^2}{2} \right) c_1 - \text{BesselK} \left(\frac{3}{4}, \frac{x^2}{2} \right) c_2 \right)}{\text{BesselI} \left(\frac{1}{4}, \frac{x^2}{2} \right) c_1 + \text{BesselK} \left(\frac{1}{4}, \frac{x^2}{2} \right) c_2}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = \frac{x \left(\text{BesselI} \left(-\frac{3}{4}, \frac{x^2}{2} \right) c_3 - \text{BesselK} \left(\frac{3}{4}, \frac{x^2}{2} \right) \right)}{\text{BesselI} \left(\frac{1}{4}, \frac{x^2}{2} \right) c_3 + \text{BesselK} \left(\frac{1}{4}, \frac{x^2}{2} \right)}$$

Summary

The solution(s) found are the following

$$y = \frac{x \left(\text{BesselI} \left(-\frac{3}{4}, \frac{x^2}{2} \right) c_3 - \text{BesselK} \left(\frac{3}{4}, \frac{x^2}{2} \right) \right)}{\text{BesselI} \left(\frac{1}{4}, \frac{x^2}{2} \right) c_3 + \text{BesselK} \left(\frac{1}{4}, \frac{x^2}{2} \right)} \quad (1)$$




Figure 45: Slope field plot

Verification of solutions

$$y = \frac{x \left(\text{BesselI} \left(-\frac{3}{4}, \frac{x^2}{2} \right) c_3 - \text{BesselK} \left(\frac{3}{4}, \frac{x^2}{2} \right) \right)}{\text{BesselI} \left(\frac{1}{4}, \frac{x^2}{2} \right) c_3 + \text{BesselK} \left(\frac{1}{4}, \frac{x^2}{2} \right)}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying Chini  
differential order: 1; looking for linear symmetries  
trying exact  
Looking for potential symmetries  
trying Riccati  
trying Riccati Special  
<- Riccati Special successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 44

```
dsolve(diff(y(x),x)=x^2-y(x)^2,y(x), singsol=all)
```

$$y(x) = \frac{x \left(\text{BesselI} \left(-\frac{3}{4}, \frac{x^2}{2} \right) c_1 - \text{BesselK} \left(\frac{3}{4}, \frac{x^2}{2} \right) \right)}{c_1 \text{BesselI} \left(\frac{1}{4}, \frac{x^2}{2} \right) + \text{BesselK} \left(\frac{1}{4}, \frac{x^2}{2} \right)}$$

✓ Solution by Mathematica

Time used: 0.183 (sec). Leaf size: 197

```
DSolve[y'[x]==x^2-y[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$y(x) \rightarrow$

$$\frac{-ix^2 \left(2 \text{BesselJ} \left(-\frac{3}{4}, \frac{ix^2}{2} \right) + c_1 \left(\text{BesselJ} \left(-\frac{5}{4}, \frac{ix^2}{2} \right) - \text{BesselJ} \left(\frac{3}{4}, \frac{ix^2}{2} \right) \right) \right) - c_1 \text{BesselJ} \left(-\frac{1}{4}, \frac{ix^2}{2} \right)}{2x \left(\text{BesselJ} \left(\frac{1}{4}, \frac{ix^2}{2} \right) + c_1 \text{BesselJ} \left(-\frac{1}{4}, \frac{ix^2}{2} \right) \right)}$$
$$y(x) \rightarrow \frac{ix^2 \text{BesselJ} \left(-\frac{5}{4}, \frac{ix^2}{2} \right) - ix^2 \text{BesselJ} \left(\frac{3}{4}, \frac{ix^2}{2} \right) + \text{BesselJ} \left(-\frac{1}{4}, \frac{ix^2}{2} \right)}{2x \text{BesselJ} \left(-\frac{1}{4}, \frac{ix^2}{2} \right)}$$

1.25 problem Problem 36

1.25.1 Solving as first order ode lie symmetry calculated ode 229

Internal problem ID [12136]

Internal file name [OUTPUT/10788_Tuesday_September_12_2023_08_52_15_AM_63960132/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 36.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$2y + (x + y - 2)y' = -2x + 1$$

1.25.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = -\frac{2x + 2y - 1}{x + y - 2}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \tag{1E}$$

$$\eta = xb_2 + yb_3 + b_1 \tag{2E}$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 - \frac{(2x + 2y - 1)(b_3 - a_2)}{x + y - 2} - \frac{(2x + 2y - 1)^2 a_3}{(x + y - 2)^2} \\ - \left(-\frac{2}{x + y - 2} + \frac{2x + 2y - 1}{(x + y - 2)^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(-\frac{2}{x + y - 2} + \frac{2x + 2y - 1}{(x + y - 2)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{2x^2a_2 - 4x^2a_3 + x^2b_2 - 2x^2b_3 + 4xya_2 - 8xya_3 + 2xyb_2 - 4xyb_3 + 2y^2a_2 - 4y^2a_3 + y^2b_2 - 2y^2b_3 - 8xa_2 - 8ya_3 - 8xb_2 - 8yb_3 - 3a_1 + 2a_2 - a_3 - 3b_1 + 4b_2 - 2b_3}{(x + y - 2)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} 2x^2a_2 - 4x^2a_3 + x^2b_2 - 2x^2b_3 + 4xya_2 - 8xya_3 + 2xyb_2 - 4xyb_3 \\ + 2y^2a_2 - 4y^2a_3 + y^2b_2 - 2y^2b_3 - 8xa_2 + 4xa_3 - 7xb_2 + 5xb_3 \\ - 5ya_2 + ya_3 - 4yb_2 + 2yb_3 - 3a_1 + 2a_2 - a_3 - 3b_1 + 4b_2 - 2b_3 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} 2a_2v_1^2 + 4a_2v_1v_2 + 2a_2v_2^2 - 4a_3v_1^2 - 8a_3v_1v_2 - 4a_3v_2^2 + b_2v_1^2 + 2b_2v_1v_2 \\ + b_2v_2^2 - 2b_3v_1^2 - 4b_3v_1v_2 - 2b_3v_2^2 - 8a_2v_1 - 5a_2v_2 + 4a_3v_1 + a_3v_2 \\ - 7b_2v_1 - 4b_2v_2 + 5b_3v_1 + 2b_3v_2 - 3a_1 + 2a_2 - a_3 - 3b_1 + 4b_2 - 2b_3 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned} & (2a_2 - 4a_3 + b_2 - 2b_3)v_1^2 + (4a_2 - 8a_3 + 2b_2 - 4b_3)v_1v_2 \\ & + (-8a_2 + 4a_3 - 7b_2 + 5b_3)v_1 + (2a_2 - 4a_3 + b_2 - 2b_3)v_2^2 \\ & + (-5a_2 + a_3 - 4b_2 + 2b_3)v_2 - 3a_1 + 2a_2 - a_3 - 3b_1 + 4b_2 - 2b_3 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} -8a_2 + 4a_3 - 7b_2 + 5b_3 &= 0 \\ -5a_2 + a_3 - 4b_2 + 2b_3 &= 0 \\ 2a_2 - 4a_3 + b_2 - 2b_3 &= 0 \\ 4a_2 - 8a_3 + 2b_2 - 4b_3 &= 0 \\ -3a_1 + 2a_2 - a_3 - 3b_1 + 4b_2 - 2b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= -b_1 - a_2 \\ a_2 &= a_2 \\ a_3 &= a_2 \\ b_1 &= b_1 \\ b_2 &= -2a_2 \\ b_3 &= -2a_2 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= -1 \\ \eta &= 1 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= 1 - \left(-\frac{2x + 2y - 1}{x + y - 2} \right) (-1) \\ &= \frac{-x - y - 1}{x + y - 2} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{-x-y-1}{x+y-2}} dy \end{aligned}$$

Which results in

$$S = -y + 3 \ln(x + y + 1)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{2x + 2y - 1}{x + y - 2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{3}{x + y + 1} \\ S_y &= -1 + \frac{3}{x + y + 1} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 2 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 2$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = 2R + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-y + 3 \ln(x + y + 1) = 2x + c_1$$



Which simplifies to

$$-y + 3 \ln(x + y + 1) = 2x + c_1$$

Which gives

$$y = -3 \text{LambertW} \left(-\frac{e^{\frac{x}{3} + \frac{c_1}{3} - \frac{1}{3}}}{3} \right) - x - 1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{2x+2y-1}{x+y-2}$ 	$R = x$ $S = -y + 3 \ln(x + y + 1)$	$\frac{dS}{dR} = 2$ 

Summary

The solution(s) found are the following

$$y = -3 \text{LambertW} \left(-\frac{e^{\frac{x}{3} + \frac{c_1}{3} - \frac{1}{3}}}{3} \right) - x - 1 \tag{1}$$




Figure 46: Slope field plot

Verification of solutions

$$y = -3 \operatorname{LambertW} \left(-\frac{e^{\frac{x}{3} + \frac{c_1}{3} - \frac{1}{3}}}{3} \right) - x - 1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous C  
1st order, trying the canonical coordinates of the invariance group  
<- 1st order, canonical coordinates successful  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.032 (sec). Leaf size: 21

```
dsolve((2*x+2*y(x)-1)+(x+y(x)-2)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -x - 3 \operatorname{LambertW}\left(-\frac{c_1 e^{\frac{x}{3} - \frac{1}{3}}}{3}\right) - 1$$

✓ Solution by Mathematica

Time used: 5.15 (sec). Leaf size: 35

```
DSolve[(2*x+2*y[x]-1)+(x+y[x]-2)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned}y(x) &\rightarrow -3W\left(-e^{\frac{x}{3}-1+c_1}\right) - x - 1 \\y(x) &\rightarrow -x - 1\end{aligned}$$

1.26 problem Problem 37

1.26.1 Maple step by step solution 239

Internal problem ID [12137]

Internal file name [OUTPUT/10789_Tuesday_September_12_2023_08_52_16_AM_73897234/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 37.

ODE order: 1.

ODE degree: 3.

The type(s) of ODE detected by this program : "**quadrature**"

Maple gives the following as the ode type

[_quadrature]

$$y'^3 - y'e^{2x} = 0$$

Solving the given ode for y' results in 3 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = 0 \tag{1}$$

$$y' = e^x \tag{2}$$

$$y' = -e^x \tag{3}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned} y &= \int 0 \, dx \\ &= c_1 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \tag{1}$$

Verification of solutions

$$y = c_1$$

Verified OK.

Solving equation (2)

Integrating both sides gives

$$\begin{aligned}y &= \int e^x dx \\ &= e^x + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^x + c_2 \tag{1}$$

Verification of solutions

$$y = e^x + c_2$$

Verified OK.

Solving equation (3)

Integrating both sides gives

$$\begin{aligned}y &= \int -e^x dx \\ &= -e^x + c_3\end{aligned}$$

Summary

The solution(s) found are the following

$$y = -e^x + c_3 \tag{1}$$

Verification of solutions

$$y = -e^x + c_3$$

Verified OK.

1.26.1 Maple step by step solution

Let's solve

$$y'^3 - y'e^{2x} = 0$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Integrate both sides with respect to x

$$\int (y'^3 - y'e^{2x}) dx = \int 0 dx + c_1$$

- Cannot compute integral

$$\int (y'^3 - y'e^{2x}) dx = c_1$$

Maple trace

```
`Methods for first order ODEs:
-> Solving 1st order ODE of high degree, 1st attempt
trying 1st order WeierstrassP solution for high degree ODE
trying 1st order WeierstrassPPrime solution for high degree ODE
trying 1st order JacobiSN solution for high degree ODE
trying 1st order ODE linearizable_by_differentiation
trying differential order: 1; missing variables
<- differential order: 1; missing y(x) successful
Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 21

```
dsolve(diff(y(x),x)^3-diff(y(x),x)*exp(2*x)=0,y(x), singsol=all)
```

$$y(x) = -e^x + c_1$$

$$y(x) = e^x + c_1$$

$$y(x) = c_1$$

✓ Solution by Mathematica

Time used: 0.007 (sec). Leaf size: 29

```
DSolve[y'[x]^3-y'[x]*Exp[2*x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1$$

$$y(x) \rightarrow -e^x + c_1$$

$$y(x) \rightarrow e^x + c_1$$

1.27 problem Problem 39

1.27.1 Solving as dAlembert ode 241

Internal problem ID [12138]

Internal file name [OUTPUT/10790_Tuesday_September_12_2023_08_52_17_AM_10546755/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 39.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "dAlembert"

Maple gives the following as the ode type

```
[[_1st_order , _with_linear_symmetries] , _dAlembert]
```

$$y - 5y'x + y'^2 = 0$$

1.27.1 Solving as dAlembert ode

Let $p = y'$ the ode becomes

$$p^2 - 5px + y = 0$$

Solving for y from the above results in

$$y = -p^2 + 5px \tag{1A}$$

This has the form

$$y = xf(p) + g(p) \tag{*}$$

Where f, g are functions of $p = y'(x)$. The above ode is dAlembert ode which is now solved. Taking derivative of (*) w.r.t. x gives

$$\begin{aligned} p &= f + (xf' + g')\frac{dp}{dx} \\ p - f &= (xf' + g')\frac{dp}{dx} \end{aligned} \tag{2}$$

Comparing the form $y = xf + g$ to (1A) shows that

$$\begin{aligned}f &= 5p \\g &= -p^2\end{aligned}$$

Hence (2) becomes

$$-4p = (5x - 2p)p'(x) \tag{2A}$$

The singular solution is found by setting $\frac{dp}{dx} = 0$ in the above which gives

$$-4p = 0$$

Solving for p from the above gives

$$p = 0$$

Substituting these in (1A) gives

$$y = 0$$

The general solution is found when $\frac{dp}{dx} \neq 0$. From eq. (2A). This results in

$$p'(x) = -\frac{4p(x)}{5x - 2p(x)} \tag{3}$$

This ODE is now solved for $p(x)$.

Inverting the above ode gives

$$\frac{d}{dp}x(p) = -\frac{5x(p) - 2p}{4p} \tag{4}$$

This ODE is now solved for $x(p)$.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$\frac{d}{dp}x(p) + p(p)x(p) = q(p)$$

Where here

$$\begin{aligned}p(p) &= \frac{5}{4p} \\q(p) &= \frac{1}{2}\end{aligned}$$

Hence the ode is

$$\frac{d}{dp}x(p) + \frac{5x(p)}{4p} = \frac{1}{2}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{5}{4p} dp} \\ &= p^{\frac{5}{4}}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dp}(\mu x) &= (\mu) \left(\frac{1}{2}\right) \\ \frac{d}{dp}\left(p^{\frac{5}{4}}x\right) &= \left(p^{\frac{5}{4}}\right) \left(\frac{1}{2}\right) \\ d\left(p^{\frac{5}{4}}x\right) &= \left(\frac{p^{\frac{5}{4}}}{2}\right) dp\end{aligned}$$

Integrating gives

$$\begin{aligned}p^{\frac{5}{4}}x &= \int \frac{p^{\frac{5}{4}}}{2} dp \\ p^{\frac{5}{4}}x &= \frac{2p^{\frac{9}{4}}}{9} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = p^{\frac{5}{4}}$ results in

$$x(p) = \frac{2p}{9} + \frac{c_1}{p^{\frac{5}{4}}}$$

Now we need to eliminate p between the above and (1A). One way to do this is by solving (1) for p . This results in

$$\begin{aligned}p &= \frac{5x}{2} + \frac{\sqrt{25x^2 - 4y}}{2} \\ p &= \frac{5x}{2} - \frac{\sqrt{25x^2 - 4y}}{2}\end{aligned}$$

Substituting the above in the solution for x found above gives

$$\begin{aligned}x &= \frac{5x}{9} + \frac{\sqrt{25x^2 - 4y}}{9} + \frac{4c_1\sqrt{2}}{(10x + 2\sqrt{25x^2 - 4y})^{\frac{5}{4}}} \\ x &= \frac{5x}{9} - \frac{\sqrt{25x^2 - 4y}}{9} + \frac{4c_1\sqrt{2}}{(10x - 2\sqrt{25x^2 - 4y})^{\frac{5}{4}}}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = 0 \tag{1}$$

$$x = \frac{5x}{9} + \frac{\sqrt{25x^2 - 4y}}{9} + \frac{4c_1\sqrt{2}}{(10x + 2\sqrt{25x^2 - 4y})^{\frac{5}{4}}} \tag{2}$$

$$x = \frac{5x}{9} - \frac{\sqrt{25x^2 - 4y}}{9} + \frac{4c_1\sqrt{2}}{(10x - 2\sqrt{25x^2 - 4y})^{\frac{5}{4}}} \tag{3}$$

Verification of solutions

$$y = 0$$

Verified OK.

$$x = \frac{5x}{9} + \frac{\sqrt{25x^2 - 4y}}{9} + \frac{4c_1\sqrt{2}}{(10x + 2\sqrt{25x^2 - 4y})^{\frac{5}{4}}}$$

Verified OK.

$$x = \frac{5x}{9} - \frac{\sqrt{25x^2 - 4y}}{9} + \frac{4c_1\sqrt{2}}{(10x - 2\sqrt{25x^2 - 4y})^{\frac{5}{4}}}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
-> Solving 1st order ODE of high degree, 1st attempt  
trying 1st order WeierstrassP solution for high degree ODE  
trying 1st order WeierstrassPPrime solution for high degree ODE  
trying 1st order JacobiSN solution for high degree ODE  
trying 1st order ODE linearizable_by_differentiation  
trying differential order: 1; missing variables  
trying dAlembert  
<- dAlembert successful`
```

✓ Solution by Maple

Time used: 0.063 (sec). Leaf size: 93

```
dsolve(y(x)=5*x*diff(y(x),x)-diff(y(x),x)^2,y(x), singsol=all)
```

$$-\frac{4\sqrt{2}c_1}{\left(10x - 2\sqrt{25x^2 - 4y(x)}\right)^{\frac{5}{4}}} + \frac{4x}{9} + \frac{\sqrt{25x^2 - 4y(x)}}{9} = 0$$
$$-\frac{4\sqrt{2}c_1}{\left(10x + 2\sqrt{25x^2 - 4y(x)}\right)^{\frac{5}{4}}} + \frac{4x}{9} - \frac{\sqrt{25x^2 - 4y(x)}}{9} = 0$$

✓ Solution by Mathematica

Time used: 60.449 (sec). Leaf size: 2233

```
DSolve[y[x]==5*x*y'[x]-y'[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

Too large to display

1.28 problem Problem 40

1.28.1 Existence and uniqueness analysis	246
1.28.2 Solving as riccati ode	247

Internal problem ID [12139]

Internal file name [OUTPUT/10791_Tuesday_September_12_2023_08_53_19_AM_27305349/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 40.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**riccati**"

Maple gives the following as the ode type

```
[[_Riccati, _special]]
```

$$y' + y^2 = x$$

With initial conditions

$$[y(1) = 0]$$

1.28.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned}y' &= f(x, y) \\ &= -y^2 + x\end{aligned}$$

The x domain of $f(x, y)$ when $y = 0$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 1$ is inside this domain. The y domain of $f(x, y)$ when $x = 1$ is

$$\{-\infty < y < \infty\}$$

And the point $y_0 = 0$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial y} &= \frac{\partial}{\partial y}(-y^2 + x) \\ &= -2y\end{aligned}$$

The y domain of $\frac{\partial f}{\partial y}$ when $x = 1$ is

$$\{-\infty < y < \infty\}$$

And the point $y_0 = 0$ is inside this domain. Therefore solution exists and is unique.

1.28.2 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= -y^2 + x\end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = -y^2 + x$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = x$, $f_1(x) = 0$ and $f_2(x) = -1$. Let

$$\begin{aligned}y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{-u}\end{aligned}\tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0\tag{2}$$

But

$$\begin{aligned}f_2' &= 0 \\ f_1 f_2 &= 0 \\ f_2^2 f_0 &= x\end{aligned}$$

Substituting the above terms back in equation (2) gives

$$-u''(x) + xu(x) = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)$$

The above shows that

$$u'(x) = c_1 \text{AiryAi}(1, x) + c_2 \text{AiryBi}(1, x)$$

Using the above in (1) gives the solution

$$y = \frac{c_1 \text{AiryAi}(1, x) + c_2 \text{AiryBi}(1, x)}{c_1 \text{AiryAi}(x) + c_2 \text{AiryBi}(x)}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = \frac{c_3 \text{AiryAi}(1, x) + \text{AiryBi}(1, x)}{c_3 \text{AiryAi}(x) + \text{AiryBi}(x)}$$

Initial conditions are used to solve for c_3 . Substituting $x = 1$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = \frac{c_3 \text{AiryAi}(1, 1) + \text{AiryBi}(1, 1)}{c_3 \text{AiryAi}(1) + \text{AiryBi}(1)}$$

$$c_3 = -\frac{\text{AiryBi}(1, 1)}{\text{AiryAi}(1, 1)}$$


Substituting c_3 found above in the general solution gives

$$y = \frac{\text{AiryBi}(1, x) \text{AiryAi}(1, 1) - \text{AiryBi}(1, 1) \text{AiryAi}(1, x)}{\text{AiryBi}(x) \text{AiryAi}(1, 1) - \text{AiryBi}(1, 1) \text{AiryAi}(x)}$$


Summary

The solution(s) found are the following

$$y = \frac{\text{AiryBi}(1, x) \text{AiryAi}(1, 1) - \text{AiryBi}(1, 1) \text{AiryAi}(1, x)}{\text{AiryBi}(x) \text{AiryAi}(1, 1) - \text{AiryBi}(1, 1) \text{AiryAi}(x)} \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = \frac{\text{AiryBi}(1, x) \text{AiryAi}(1, 1) - \text{AiryBi}(1, 1) \text{AiryAi}(1, x)}{\text{AiryBi}(x) \text{AiryAi}(1, 1) - \text{AiryBi}(1, 1) \text{AiryAi}(x)}$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
differential order: 1; looking for linear symmetries
trying exact
Looking for potential symmetries
trying Riccati
trying Riccati Special
<- Riccati Special successful`

```

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 37

```
dsolve([diff(y(x),x)=x-y(x)^2,y(1) = 0],y(x), singsol=all)
```

$$y(x) = \frac{\text{AiryBi}(1, 1) \text{AiryAi}(1, x) - \text{AiryBi}(1, x) \text{AiryAi}(1, 1)}{\text{AiryBi}(1, 1) \text{AiryAi}(x) - \text{AiryBi}(x) \text{AiryAi}(1, 1)}$$

✓ Solution by Mathematica

Time used: 0.206 (sec). Leaf size: 229

```
DSolve[{y'[x]==x-y[x]^2,{y[1]==0}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{i(x^{3/2}(-\text{BesselJ}(-\frac{4}{3}, \frac{2i}{3}) + i\text{BesselJ}(-\frac{1}{3}, \frac{2i}{3}) + \text{BesselJ}(\frac{2}{3}, \frac{2i}{3})))\text{BesselJ}(-\frac{2}{3}, \frac{2}{3}ix^{3/2}) + x^{3/2}\text{BesselJ}(-\frac{2}{3}, \frac{2}{3}ix^{3/2})}{x(2\text{BesselJ}(-\frac{2}{3}, \frac{2i}{3})\text{BesselJ}(-\frac{1}{3}, \frac{2}{3}ix^{3/2}) + (-\text{BesselJ}(-\frac{4}{3}, \frac{2i}{3}) + i\text{BesselJ}(-\frac{1}{3}, \frac{2i}{3}) + \text{BesselJ}(\frac{2}{3}, \frac{2i}{3}))\text{BesselJ}(-\frac{2}{3}, \frac{2}{3}ix^{3/2}))}$$

1.29 problem Problem 42

1.29.1 Solving as first order ode lie symmetry calculated ode 251

Internal problem ID [12140]

Internal file name [OUTPUT/10792_Tuesday_September_12_2023_08_53_28_AM_81792238/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 42.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_homogeneous , `class C`], _dAlembert]
```

$$y' - (x - 5y)^{\frac{1}{3}} = 2$$

1.29.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = (x - 5y)^{\frac{1}{3}} + 2$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (\text{1E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 + \left((x - 5y)^{\frac{1}{3}} + 2 \right) (b_3 - a_2) - \left((x - 5y)^{\frac{1}{3}} + 2 \right)^2 a_3 \quad (5E)$$

$$- \frac{xa_2 + ya_3 + a_1}{3(x - 5y)^{\frac{2}{3}}} + \frac{\frac{5xb_2}{3} + \frac{5yb_3}{3} + \frac{5b_1}{3}}{(x - 5y)^{\frac{2}{3}}} = 0$$

Putting the above in normal form gives

$$\frac{3(x - 5y)^{\frac{4}{3}} a_3 + 6(x - 5y)^{\frac{2}{3}} a_2 + 12a_3(x - 5y)^{\frac{2}{3}} - 3b_2(x - 5y)^{\frac{2}{3}} - 6(x - 5y)^{\frac{2}{3}} b_3 + 4xa_2 + 12a_3x - 5xb_2}{3(x - 5y)^{\frac{2}{3}}}$$

$$= 0$$

Setting the numerator to zero gives

$$-3(x - 5y)^{\frac{4}{3}} a_3 - 6(x - 5y)^{\frac{2}{3}} a_2 - 12a_3(x - 5y)^{\frac{2}{3}} + 3b_2(x - 5y)^{\frac{2}{3}} + 6(x - 5y)^{\frac{2}{3}} b_3 \quad (6E)$$

$$- 4xa_2 - 12a_3x + 5xb_2 + 3b_3x + 15a_2y + 59ya_3 - 10yb_3 - a_1 + 5b_1 = 0$$

Simplifying the above gives

$$-3(x - 5y)^{\frac{4}{3}} a_3 - 3(x - 5y) a_2 - 12(x - 5y) a_3 + 3(x - 5y) b_3 \quad (6E)$$

$$- 6(x - 5y)^{\frac{2}{3}} a_2 - 12a_3(x - 5y)^{\frac{2}{3}} + 3b_2(x - 5y)^{\frac{2}{3}}$$

$$+ 6(x - 5y)^{\frac{2}{3}} b_3 - xa_2 + 5xb_2 - ya_3 + 5yb_3 - a_1 + 5b_1 = 0$$

Since the PDE has radicals, simplifying gives

$$-3(x - 5y)^{\frac{1}{3}} a_3x - 6(x - 5y)^{\frac{2}{3}} a_2 - 12a_3(x - 5y)^{\frac{2}{3}} + 3b_2(x - 5y)^{\frac{2}{3}} + 6(x - 5y)^{\frac{2}{3}} b_3$$

$$+ 15(x - 5y)^{\frac{1}{3}} a_3y - 4xa_2 - 12a_3x + 5xb_2 + 3b_3x + 15a_2y + 59ya_3 - 10yb_3 - a_1 + 5b_1 = 0$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\left\{ x, y, (x - 5y)^{\frac{1}{3}}, (x - 5y)^{\frac{2}{3}} \right\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\left\{ x = v_1, y = v_2, (x - 5y)^{\frac{1}{3}} = v_3, (x - 5y)^{\frac{2}{3}} = v_4 \right\}$$

The above PDE (6E) now becomes

$$\begin{aligned} & -3v_3a_3v_1 + 15v_3a_3v_2 - 4v_1a_2 + 15a_2v_2 - 6v_4a_2 - 12a_3v_1 + 59v_2a_3 \\ & - 12a_3v_4 + 5v_1b_2 + 3b_2v_4 + 3b_3v_1 - 10v_2b_3 + 6v_4b_3 - a_1 + 5b_1 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3, v_4\}$$

Equation (7E) now becomes

$$\begin{aligned} & -3v_3a_3v_1 + (-4a_2 - 12a_3 + 5b_2 + 3b_3)v_1 + 15v_3a_3v_2 \\ & + (15a_2 + 59a_3 - 10b_3)v_2 + (-6a_2 - 12a_3 + 3b_2 + 6b_3)v_4 - a_1 + 5b_1 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} -3a_3 &= 0 \\ 15a_3 &= 0 \\ -a_1 + 5b_1 &= 0 \\ 15a_2 + 59a_3 - 10b_3 &= 0 \\ -6a_2 - 12a_3 + 3b_2 + 6b_3 &= 0 \\ -4a_2 - 12a_3 + 5b_2 + 3b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 5b_1 \\ a_2 &= 0 \\ a_3 &= 0 \\ b_1 &= b_1 \\ b_2 &= 0 \\ b_3 &= 0 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= 5 \\ \eta &= 1 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}\eta &= \eta - \omega(x, y) \xi \\ &= 1 - \left((x - 5y)^{\frac{1}{3}} + 2 \right) \quad (5) \\ &= -9 - 5(x - 5y)^{\frac{1}{3}} \\ \xi &= 0\end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{-9 - 5(x - 5y)^{\frac{1}{3}}} dy\end{aligned}$$

Which results in

$$S = \frac{81 \ln(729 + 125x - 625y)}{625} - \frac{27(x - 5y)^{\frac{1}{3}}}{125} - \frac{81 \ln\left(25(x - 5y)^{\frac{2}{3}} - 45(x - 5y)^{\frac{1}{3}} + 81\right)}{625} + \frac{162 \ln\left(5(x - 5y)^{\frac{1}{3}} + 3\right)}{625}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = (x - 5y)^{\frac{1}{3}} + 2$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{1}{25(x-5y)^{\frac{1}{3}} + 45} \\ S_y &= \frac{1}{-9 - 5(x-5y)^{\frac{1}{3}}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -\frac{1}{5} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{1}{5}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\frac{R}{5} + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{81 \ln(729 + 125x - 625y)}{625} - \frac{27(x-5y)^{\frac{1}{3}}}{125} - \frac{81 \ln\left(25(x-5y)^{\frac{2}{3}} - 45(x-5y)^{\frac{1}{3}} + 81\right)}{625} + \frac{162 \ln\left(5(x-5y)\right)}{625}$$

Which simplifies to

$$\frac{81 \ln(729 + 125x - 625y)}{625} - \frac{27(x-5y)^{\frac{1}{3}}}{125} - \frac{81 \ln\left(25(x-5y)^{\frac{2}{3}} - 45(x-5y)^{\frac{1}{3}} + 81\right)}{625} + \frac{162 \ln\left(5(x-5y)\right)}{625}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = (x - 5y)^{\frac{1}{3}} + 2$ 	$R = x$ $S = \frac{81 \ln(729 + 125x - 625y) - 125(x - 5y)^{\frac{1}{3}}}{625}$	$\frac{dS}{dR} = -\frac{1}{5}$ 

Summary

The solution(s) found are the following

$$\begin{aligned}
 & \frac{81 \ln(729 + 125x - 625y) - 125(x - 5y)^{\frac{1}{3}}}{625} - \frac{27(x - 5y)^{\frac{1}{3}}}{125} \\
 & - \frac{81 \ln\left(25(x - 5y)^{\frac{2}{3}} - 45(x - 5y)^{\frac{1}{3}} + 81\right)}{625} \\
 & + \frac{162 \ln\left(5(x - 5y)^{\frac{1}{3}} + 9\right)}{625} + \frac{3(x - 5y)^{\frac{2}{3}}}{50} = -\frac{x}{5} + c_1
 \end{aligned} \tag{1}$$




Figure 48: Slope field plot

Verification of solutions

$$\begin{aligned}
 & \frac{81 \ln(729 + 125x - 625y)}{625} - \frac{27(x - 5y)^{\frac{1}{3}}}{125} \\
 & - \frac{81 \ln\left(25(x - 5y)^{\frac{2}{3}} - 45(x - 5y)^{\frac{1}{3}} + 81\right)}{625} \\
 & + \frac{162 \ln\left(5(x - 5y)^{\frac{1}{3}} + 9\right)}{625} + \frac{3(x - 5y)^{\frac{2}{3}}}{50} = -\frac{x}{5} + c_1
 \end{aligned}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying homogeneous types:
trying homogeneous C
1st order, trying the canonical coordinates of the invariance group
  -> Calling odsolve with the ODE`, diff(y(x), x) = 1/5, y(x)` *** Sublevel 2 ***
      Methods for first order ODEs:
      --- Trying classification methods ---
      trying a quadrature
      trying 1st order linear
      <- 1st order linear successful
<- 1st order, canonical coordinates successful
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 80

```
dsolve(diff(y(x),x)=(x-5*y(x))^(1/3)+2,y(x), singsol=all)
```

$$\begin{aligned} & x + \frac{81 \ln(729 - 625y(x) + 125x)}{125} - \frac{27(x - 5y(x))^{\frac{1}{3}}}{25} \\ & - \frac{81 \ln\left(25(x - 5y(x))^{\frac{2}{3}} - 45(x - 5y(x))^{\frac{1}{3}} + 81\right)}{125} \\ & + \frac{162 \ln\left(9 + 5(x - 5y(x))^{\frac{1}{3}}\right)}{125} + \frac{3(x - 5y(x))^{\frac{2}{3}}}{10} - c_1 = 0 \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.347 (sec). Leaf size: 70

```
DSolve[y'[x]==(x-5*y[x])^(1/3)+2,y[x],x,IncludeSingularSolutions -> True]
```

$$\begin{aligned} & \text{Solve}\left[5y(x) + 5\left(-y(x) + \frac{3}{50}(x - 5y(x))^{2/3} - \frac{27}{125}\sqrt[3]{x - 5y(x)}\right) \right. \\ & \left. + \frac{243}{625} \log\left(5\sqrt[3]{x - 5y(x)} + 9\right) + \frac{x}{5}\right] = c_1, y(x) \end{aligned}$$

1.30 problem Problem 43

1.30.1 Solving as homogeneousTypeD2 ode	259
1.30.2 Solving as first order ode lie symmetry lookup ode	261
1.30.3 Solving as bernoulli ode	265
1.30.4 Solving as exact ode	269
1.30.5 Solving as riccati ode	274

Internal problem ID [12141]

Internal file name [OUTPUT/10793_Tuesday_September_12_2023_08_53_29_AM_30062239/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 43.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "riccati", "bernoulli", "exactByInspection", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _Bernoulli]
```

$$y(-y + x) - x^2y' = 0$$

1.30.1 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u(x)x(-u(x)x + x) - x^2(u'(x)x + u(x)) = 0$$

In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u^2}{x}\end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = u^2$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u^2} du &= -\frac{1}{x} dx \\ \int \frac{1}{u^2} du &= \int -\frac{1}{x} dx \\ -\frac{1}{u} &= -\ln(x) + c_2\end{aligned}$$

The solution is

$$-\frac{1}{u(x)} + \ln(x) - c_2 = 0$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$\begin{aligned}-\frac{x}{y} + \ln(x) - c_2 &= 0 \\ -\frac{x}{y} + \ln(x) - c_2 &= 0\end{aligned}$$

Summary

The solution(s) found are the following

$$-\frac{x}{y} + \ln(x) - c_2 = 0 \tag{1}$$




Figure 49: Slope field plot

Verification of solutions

$$-\frac{x}{y} + \ln(x) - c_2 = 0$$

Verified OK.

1.30.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{y(y-x)}{x^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **Bernoulli**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 33: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{y^2}{x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right)S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{y^2}{x}} dy \end{aligned}$$

Which results in

$$S = -\frac{x}{y}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{y(y-x)}{x^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{1}{y} \\ S_y &= \frac{x}{y^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -\frac{1}{x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{1}{R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\ln(R) + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{x}{y} = -\ln(x) + c_1$$



Which simplifies to

$$-\frac{x}{y} = -\ln(x) + c_1$$

Which gives

$$y = \frac{x}{\ln(x) - c_1}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{y(y-x)}{x^2}$ 	$R = x$ $S = -\frac{x}{y}$	$\frac{dS}{dR} = -\frac{1}{R}$ 

Summary

The solution(s) found are the following

$$y = \frac{x}{\ln(x) - c_1} \quad (1)$$




Figure 50: Slope field plot

Verification of solutions

$$y = \frac{x}{\ln(x) - c_1}$$

Verified OK.

1.30.3 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= -\frac{y(y-x)}{x^2} \end{aligned}$$

This is a Bernoulli ODE.

$$y' = \frac{1}{x}y - \frac{1}{x^2}y^2 \quad (1)$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \quad (2)$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \quad (3)$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned} f_0(x) &= \frac{1}{x} \\ f_1(x) &= -\frac{1}{x^2} \\ n &= 2 \end{aligned}$$

Dividing both sides of ODE (1) by $y^n = y^2$ gives

$$y' \frac{1}{y^2} = \frac{1}{xy} - \frac{1}{x^2} \quad (4)$$

Let

$$\begin{aligned} w &= y^{1-n} \\ &= \frac{1}{y} \end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = -\frac{1}{y^2}y' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} -w'(x) &= \frac{w(x)}{x} - \frac{1}{x^2} \\ w' &= -\frac{w}{x} + \frac{1}{x^2} \end{aligned} \quad (7)$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{1}{x^2}$$

Hence the ode is

$$w'(x) + \frac{w(x)}{x} = \frac{1}{x^2}$$

The integrating factor μ is

$$\mu = e^{\int \frac{1}{x} dx}$$
$$= x$$

The ode becomes

$$\frac{d}{dx}(\mu w) = (\mu) \left(\frac{1}{x^2} \right)$$
$$\frac{d}{dx}(wx) = (x) \left(\frac{1}{x^2} \right)$$
$$d(wx) = \frac{1}{x} dx$$

Integrating gives

$$wx = \int \frac{1}{x} dx$$
$$wx = \ln(x) + c_1$$

Dividing both sides by the integrating factor $\mu = x$ results in

$$w(x) = \frac{\ln(x)}{x} + \frac{c_1}{x}$$

which simplifies to

$$w(x) = \frac{\ln(x) + c_1}{x}$$

Replacing w in the above by $\frac{1}{y}$ using equation (5) gives the final solution.

$$\frac{1}{y} = \frac{\ln(x) + c_1}{x}$$

Or

$$y = \frac{x}{\ln(x) + c_1}$$

Summary

The solution(s) found are the following

$$y = \frac{x}{\ln(x) + c_1} \tag{1}$$




Figure 51: Slope field plot

Verification of solutions

$$y = \frac{x}{\ln(x) + c_1}$$

Verified OK.

1.30.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (-x^2) dy &= (-y(-y + x)) dx \\ (y(-y + x)) dx + (-x^2) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= y(-y + x) \\ N(x, y) &= -x^2 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(y(-y+x)) \\ &= x - 2y\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(-x^2) \\ &= -2x\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. By inspection $\frac{1}{xy^2}$ is an integrating factor. Therefore by multiplying $M = y(-y+x)$ and $N = -x^2$ by this integrating factor the ode becomes exact. The new M, N are

$$\begin{aligned}M &= \frac{-y+x}{xy} \\ N &= -\frac{x}{y^2}\end{aligned}$$

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} \left(-\frac{x}{y^2}\right) dy &= \left(-\frac{-y+x}{xy}\right) dx \\ \left(\frac{-y+x}{xy}\right) dx + \left(-\frac{x}{y^2}\right) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= \frac{-y+x}{xy} \\ N(x, y) &= -\frac{x}{y^2} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{-y+x}{xy} \right) \\ &= -\frac{1}{y^2} \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(-\frac{x}{y^2} \right) \\ &= -\frac{1}{y^2} \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int \frac{-y + x}{xy} dx$$

$$\phi = -\ln(x) + \frac{x}{y} + f(y) \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = -\frac{x}{y^2} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -\frac{x}{y^2}$. Therefore equation (4) becomes

$$-\frac{x}{y^2} = -\frac{x}{y^2} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x) + \frac{x}{y} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x) + \frac{x}{y}$$

The solution becomes

$$y = \frac{x}{\ln(x) + c_1}$$

Summary

The solution(s) found are the following

$$y = \frac{x}{\ln(x) + c_1} \tag{1}$$




Figure 52: Slope field plot

Verification of solutions

$$y = \frac{x}{\ln(x) + c_1}$$

Verified OK.

1.30.5 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= -\frac{y(y-x)}{x^2}\end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = -\frac{y^2}{x^2} + \frac{y}{x}$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = 0$, $f_1(x) = \frac{1}{x}$ and $f_2(x) = -\frac{1}{x^2}$. Let

$$\begin{aligned}y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{-\frac{u}{x^2}}\end{aligned}\tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0\tag{2}$$

But

$$\begin{aligned}f_2' &= \frac{2}{x^3} \\ f_1 f_2 &= -\frac{1}{x^3} \\ f_2^2 f_0 &= 0\end{aligned}$$

Substituting the above terms back in equation (2) gives

$$-\frac{u''(x)}{x^2} - \frac{u'(x)}{x^3} = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = c_1 + \ln(x) c_2$$

The above shows that

$$u'(x) = \frac{c_2}{x}$$

Using the above in (1) gives the solution

$$y = \frac{c_2 x}{c_1 + \ln(x) c_2}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = \frac{x}{c_3 + \ln(x)}$$

Summary

The solution(s) found are the following

$$y = \frac{x}{c_3 + \ln(x)} \tag{1}$$




Figure 53: Slope field plot

Verification of solutions

$$y = \frac{x}{c_3 + \ln(x)}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve((x-y(x))*y(x)-x^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x}{\ln(x) + c_1}$$

✓ Solution by Mathematica

Time used: 0.231 (sec). Leaf size: 19

```
DSolve[(x-y[x])*y[x]-x^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x}{\log(x) + c_1}$$
$$y(x) \rightarrow 0$$

1.31 problem Problem 45

1.31.1 Existence and uniqueness analysis	277
1.31.2 Solving as linear ode	278
1.31.3 Solving as homogeneousTypeD2 ode	280
1.31.4 Solving as first order ode lie symmetry lookup ode	281
1.31.5 Solving as exact ode	285
1.31.6 Maple step by step solution	290

Internal problem ID [12142]

Internal file name [OUTPUT/10794_Tuesday_September_12_2023_08_53_30_AM_12814526/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 45.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear"**, **"homogeneousTypeD2"**, **"exactWithIntegrationFactor"**, **"first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

```
[[_linear, `class A`]]
```

$$x' + 5x = 10t + 2$$

With initial conditions

$$[x(1) = 2]$$

1.31.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$x' + p(t)x = q(t)$$

Where here

$$p(t) = 5$$

$$q(t) = 10t + 2$$

Hence the ode is

$$x' + 5x = 10t + 2$$

The domain of $p(t) = 5$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 1$ is inside this domain. The domain of $q(t) = 10t + 2$ is

$$\{-\infty < t < \infty\}$$

And the point $t_0 = 1$ is also inside this domain. Hence solution exists and is unique.

1.31.2 Solving as linear ode

Entering Linear first order ODE solver. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int 5dt} \\ &= e^{5t}\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dt}(\mu x) &= (\mu)(10t + 2) \\ \frac{d}{dt}(e^{5t}x) &= (e^{5t})(10t + 2) \\ d(e^{5t}x) &= ((10t + 2)e^{5t}) dt\end{aligned}$$

Integrating gives

$$\begin{aligned}e^{5t}x &= \int (10t + 2)e^{5t} dt \\ e^{5t}x &= 2te^{5t} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = e^{5t}$ results in

$$x = 2e^{-5t}te^{5t} + c_1e^{-5t}$$

which simplifies to

$$x = 2t + c_1e^{-5t}$$

Initial conditions are used to solve for c_1 . Substituting $t = 1$ and $x = 2$ in the above solution gives an equation to solve for the constant of integration.

$$2 = 2 + c_1 e^{-5}$$

$$c_1 = 0$$


Substituting c_1 found above in the general solution gives

$$x = 2t$$


Summary

The solution(s) found are the following

$$x = 2t \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$x = 2t$$

Verified OK.

1.31.3 Solving as homogeneous Type D2 ode

Using the change of variables $x = u(t)t$ on the above ode results in new ode in $u(t)$

$$u'(t)t + u(t) + 5u(t)t = 10t + 2$$

In canonical form the ODE is

$$\begin{aligned}u' &= F(t, u) \\ &= f(t)g(u) \\ &= \frac{(5t + 1)(-u + 2)}{t}\end{aligned}$$

Where $f(t) = \frac{5t+1}{t}$ and $g(u) = -u + 2$. Integrating both sides gives

$$\begin{aligned}\frac{1}{-u + 2} du &= \frac{5t + 1}{t} dt \\ \int \frac{1}{-u + 2} du &= \int \frac{5t + 1}{t} dt \\ -\ln(u - 2) &= 5t + \ln(t) + c_2\end{aligned}$$

Raising both side to exponential gives

$$\frac{1}{u - 2} = e^{5t + \ln(t) + c_2}$$

Which simplifies to

$$\frac{1}{u - 2} = c_3 e^{5t + \ln(t)}$$

Which simplifies to

$$u(t) = \frac{(2c_3 e^{5t} t e^{c_2} + 1) e^{-5t} e^{-c_2}}{c_3 t}$$

Therefore the solution x is

$$\begin{aligned}x &= ut \\ &= \frac{(2c_3 e^{5t} t e^{c_2} + 1) e^{-5t} e^{-c_2}}{c_3}\end{aligned}$$

Initial conditions are used to solve for c_2 . Substituting $t = 1$ and $x = 2$ in the above solution gives an equation to solve for the constant of integration.


$$2 = \frac{2e^{5+c_2} e^{-5-c_2} c_3 + e^{-5-c_2}}{c_3}$$

Unable to solve for constant of integration. Since $\lim_{c_1 \rightarrow \infty}$ gives $x = \frac{(2c_3 e^{5t} t e^{c_2} + 1) e^{-5t} e^{-c_2}}{c_3} =$


Summary

$x = 2t$ and this result satisfies the given initial condition. The solution(s) found are the following

$$x = 2t$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$x = 2t$$

Verified OK.

1.31.4 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$x' = -5x + 10t + 2$$

$$x' = \omega(t, x)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_t + \omega(\eta_x - \xi_t) - \omega^2 \xi_x - \omega_t \xi - \omega_x \eta = 0 \tag{A}$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 35: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int(n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(t, x) &= 0 \\ \eta(t, x) &= e^{-5t}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(t, x) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dt}{\xi} = \frac{dx}{\eta} = dS\tag{1}$$

The above comes from the requirements that $(\xi \frac{\partial}{\partial t} + \eta \frac{\partial}{\partial x}) S(t, x) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = t$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-5t}} dy \end{aligned}$$

Which results in

$$S = e^{5t} x$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_t + \omega(t, x)S_x}{R_t + \omega(t, x)R_x} \quad (2)$$

Where in the above R_t, R_x, S_t, S_x are all partial derivatives and $\omega(t, x)$ is the right hand side of the original ode given by

$$\omega(t, x) = -5x + 10t + 2$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_t &= 1 \\ R_x &= 0 \\ S_t &= 5 e^{5t} x \\ S_x &= e^{5t} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = (10t + 2) e^{5t} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for t, x in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = (10R + 2) e^{5R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = 2 e^{5R} R + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to t, x coordinates. This results in

$$e^{5t} x = 2t e^{5t} + c_1$$



Which simplifies to

$$e^{5t} x = 2t e^{5t} + c_1$$

Which gives

$$x = (2t e^{5t} + c_1) e^{-5t}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in t, x coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dx}{dt} = -5x + 10t + 2$ 	$R = t$ $S = e^{5t} x$	$\frac{dS}{dR} = (10R + 2) e^{5R}$ 

Initial conditions are used to solve for c_1 . Substituting $t = 1$ and $x = 2$ in the above solution gives an equation to solve for the constant of integration.

$$2 = 2 + c_1 e^{-5}$$

$$c_1 = 0$$


Substituting c_1 found above in the general solution gives

$$x = 2t$$


Summary

The solution(s) found are the following

$$x = 2t \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$x = 2t$$

Verified OK.

1.31.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(t, x) dt + N(t, x) dx = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} dx &= (-5x + 10t + 2) dt \\ (5x - 10t - 2) dt + dx &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(t, x) &= 5x - 10t - 2 \\ N(t, x) &= 1 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial x} = \frac{\partial N}{\partial t}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial x} &= \frac{\partial}{\partial x} (5x - 10t - 2) \\ &= 5 \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial t} &= \frac{\partial}{\partial t}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial x} \neq \frac{\partial N}{\partial t}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial x} - \frac{\partial N}{\partial t} \right) \\ &= 1((5) - (0)) \\ &= 5\end{aligned}$$

Since A does not depend on x , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A dt} \\ &= e^{\int 5 dt}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{5t} \\ &= e^{5t}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= e^{5t}(5x - 10t - 2) \\ &= (5x - 10t - 2)e^{5t}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= e^{5t}(1) \\ &= e^{5t}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dx}{dt} &= 0 \\ ((5x - 10t - 2)e^{5t}) + (e^{5t}) \frac{dx}{dt} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(t, x)$

$$\frac{\partial \phi}{\partial t} = \bar{M} \quad (1)$$

$$\frac{\partial \phi}{\partial x} = \bar{N} \quad (2)$$

Integrating (1) w.r.t. t gives

$$\int \frac{\partial \phi}{\partial t} dt = \int \bar{M} dt$$

$$\int \frac{\partial \phi}{\partial t} dt = \int (5x - 10t - 2) e^{5t} dt$$

$$\phi = (-2t + x) e^{5t} + f(x) \quad (3)$$

Where $f(x)$ is used for the constant of integration since ϕ is a function of both t and x . Taking derivative of equation (3) w.r.t x gives

$$\frac{\partial \phi}{\partial x} = e^{5t} + f'(x) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial x} = e^{5t}$. Therefore equation (4) becomes

$$e^{5t} = e^{5t} + f'(x) \quad (5)$$

Solving equation (5) for $f'(x)$ gives

$$f'(x) = 0$$

Therefore

$$f(x) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(x)$ into equation (3) gives ϕ

$$\phi = (-2t + x) e^{5t} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = (-2t + x) e^{5t}$$

The solution becomes

$$x = (2te^{5t} + c_1) e^{-5t}$$

Initial conditions are used to solve for c_1 . Substituting $t = 1$ and $x = 2$ in the above solution gives an equation to solve for the constant of integration.

$$2 = 2 + c_1 e^{-5}$$

$$c_1 = 0$$


Substituting c_1 found above in the general solution gives

$$x = 2t$$


Summary

The solution(s) found are the following

$$x = 2t \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$x = 2t$$

Verified OK.

1.31.6 Maple step by step solution

Let's solve

$$[x' + 5x = 10t + 2, x(1) = 2]$$

- Highest derivative means the order of the ODE is 1

$$x'$$

- Isolate the derivative

$$x' = -5x + 10t + 2$$

- Group terms with x on the lhs of the ODE and the rest on the rhs of the ODE

$$x' + 5x = 10t + 2$$

- The ODE is linear; multiply by an integrating factor $\mu(t)$

$$\mu(t)(x' + 5x) = \mu(t)(10t + 2)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dt}(\mu(t)x)$

$$\mu(t)(x' + 5x) = \mu'(t)x + \mu(t)x'$$

- Isolate $\mu'(t)$

$$\mu'(t) = 5\mu(t)$$

- Solve to find the integrating factor

$$\mu(t) = e^{5t}$$

- Integrate both sides with respect to t

$$\int \left(\frac{d}{dt}(\mu(t)x) \right) dt = \int \mu(t)(10t + 2) dt + c_1$$

- Evaluate the integral on the lhs

$$\mu(t)x = \int \mu(t)(10t + 2) dt + c_1$$

- Solve for x

$$x = \frac{\int \mu(t)(10t+2)dt+c_1}{\mu(t)}$$

- Substitute $\mu(t) = e^{5t}$

$$x = \frac{\int (10t+2)e^{5t}dt+c_1}{e^{5t}}$$

- Evaluate the integrals on the rhs

$$x = \frac{2te^{5t}+c_1}{e^{5t}}$$

- Simplify

$$x = 2t + c_1e^{-5t}$$

- Use initial condition $x(1) = 2$
 $2 = 2 + c_1 e^{-5}$
- Solve for c_1
 $c_1 = 0$
- Substitute $c_1 = 0$ into general solution and simplify
 $x = 2t$
- Solution to the IVP
 $x = 2t$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 7

```
dsolve([diff(x(t),t)+5*x(t)=10*t+2,x(1) = 2],x(t), singsol=all)
```

$$x(t) = 2t$$

✓ Solution by Mathematica

Time used: 0.082 (sec). Leaf size: 8

```
DSolve[{x'[t]+5*x[t]==10*t+2,{x[1]==2}},x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow 2t$$

1.32 problem Problem 46

1.32.1 Existence and uniqueness analysis	293
1.32.2 Solving as homogeneousTypeD2 ode	293
1.32.3 Solving as first order ode lie symmetry lookup ode	295
1.32.4 Solving as bernoulli ode	300
1.32.5 Solving as exact ode	303
1.32.6 Solving as riccati ode	308

Internal problem ID [12143]

Internal file name [OUTPUT/10795_Tuesday_September_12_2023_08_53_31_AM_95298020/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 46.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "riccati", "bernoulli", "homogeneousTypeD2", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_homogeneous, `class D`], _rational, _Bernoulli]
```

$$x' - \frac{x}{t} - \frac{x^2}{t^3} = 0$$

With initial conditions

$$[x(2) = 4]$$

1.32.1 Existence and uniqueness analysis

This is non linear first order ODE. In canonical form it is written as

$$\begin{aligned}x' &= f(t, x) \\ &= \frac{x(t^2 + x)}{t^3}\end{aligned}$$

The t domain of $f(t, x)$ when $x = 4$ is

$$\{t < 0 \vee 0 < t\}$$

And the point $t_0 = 2$ is inside this domain. The x domain of $f(t, x)$ when $t = 2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 4$ is inside this domain. Now we will look at the continuity of

$$\begin{aligned}\frac{\partial f}{\partial x} &= \frac{\partial}{\partial x} \left(\frac{x(t^2 + x)}{t^3} \right) \\ &= \frac{t^2 + x}{t^3} + \frac{x}{t^3}\end{aligned}$$

The t domain of $\frac{\partial f}{\partial x}$ when $x = 4$ is

$$\{t < 0 \vee 0 < t\}$$

And the point $t_0 = 2$ is inside this domain. The x domain of $\frac{\partial f}{\partial x}$ when $t = 2$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 4$ is inside this domain. Therefore solution exists and is unique.

1.32.2 Solving as homogeneous TypeD2 ode

Using the change of variables $x = u(t)t$ on the above ode results in new ode in $u(t)$

$$u'(t)t - \frac{u(t)^2}{t} = 0$$

In canonical form the ODE is

$$\begin{aligned}u' &= F(t, u) \\ &= f(t)g(u) \\ &= \frac{u^2}{t^2}\end{aligned}$$

Where $f(t) = \frac{1}{t^2}$ and $g(u) = u^2$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u^2} du &= \frac{1}{t^2} dt \\ \int \frac{1}{u^2} du &= \int \frac{1}{t^2} dt \\ -\frac{1}{u} &= -\frac{1}{t} + c_2\end{aligned}$$

The solution is

$$-\frac{1}{u(t)} + \frac{1}{t} - c_2 = 0$$

Replacing $u(t)$ in the above solution by $\frac{x}{t}$ results in the solution for x in implicit form

$$\begin{aligned}-\frac{t}{x} + \frac{1}{t} - c_2 &= 0 \\ -\frac{t}{x} + \frac{1}{t} - c_2 &= 0\end{aligned}$$


Substituting initial conditions and solving for c_2 gives $c_2 = 0$. Hence the solution becomes Solving for x from the above gives

$$x = t^2$$


Summary

The solution(s) found are the following

$$x = t^2 \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$x = t^2$$

Verified OK.

1.32.3 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$x' = \frac{x(t^2 + x)}{t^3}$$

$$x' = \omega(t, x)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_t + \omega(\eta_x - \xi_t) - \omega^2 \xi_x - \omega_t \xi - \omega_x \eta = 0 \tag{A}$$

The type of this ode is known. It is of type **Bernoulli**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 38: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(t, x) &= 0 \\ \eta(t, x) &= \frac{x^2}{t}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(t, x) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dt}{\xi} = \frac{dx}{\eta} = dS\tag{1}$$

The above comes from the requirements that $(\xi \frac{\partial}{\partial t} + \eta \frac{\partial}{\partial x}) S(t, x) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = t$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{x^2}{t}} dy \end{aligned}$$

Which results in

$$S = -\frac{t}{x}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_t + \omega(t, x)S_x}{R_t + \omega(t, x)R_x} \quad (2)$$

Where in the above R_t, R_x, S_t, S_x are all partial derivatives and $\omega(t, x)$ is the right hand side of the original ode given by

$$\omega(t, x) = \frac{x(t^2 + x)}{t^3}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_t &= 1 \\ R_x &= 0 \\ S_t &= -\frac{1}{x} \\ S_x &= \frac{t}{x^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{1}{t^2} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for t, x in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{R^2}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\frac{1}{R} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to t, x coordinates. This results in

$$-\frac{t}{x} = -\frac{1}{t} + c_1$$



Which simplifies to

$$-\frac{t}{x} = -\frac{1}{t} + c_1$$

Which gives

$$x = -\frac{t^2}{c_1 t - 1}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in t, x coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dx}{dt} = \frac{x(t^2+x)}{t^3}$ 	$R = t$ $S = -\frac{t}{x}$	$\frac{dS}{dR} = \frac{1}{R^2}$ 

Initial conditions are used to solve for c_1 . Substituting $t = 2$ and $x = 4$ in the above solution gives an equation to solve for the constant of integration.

$$4 = -\frac{4}{2c_1 - 1}$$

$$c_1 = 0$$


Substituting c_1 found above in the general solution gives

$$x = t^2$$


Summary

The solution(s) found are the following

$$x = t^2 \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$x = t^2$$

Verified OK.

1.32.4 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned}x' &= F(t, x) \\ &= \frac{x(t^2 + x)}{t^3}\end{aligned}$$

This is a Bernoulli ODE.

$$x' = \frac{1}{t}x + \frac{1}{t^3}x^2 \quad (1)$$

The standard Bernoulli ODE has the form

$$x' = f_0(t)x + f_1(t)x^n \quad (2)$$

The first step is to divide the above equation by x^n which gives

$$\frac{x'}{x^n} = f_0(t)x^{1-n} + f_1(t) \quad (3)$$

The next step is use the substitution $w = x^{1-n}$ in equation (3) which generates a new ODE in $w(t)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $x(t)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned}f_0(t) &= \frac{1}{t} \\ f_1(t) &= \frac{1}{t^3} \\ n &= 2\end{aligned}$$

Dividing both sides of ODE (1) by $x^n = x^2$ gives

$$x' \frac{1}{x^2} = \frac{1}{tx} + \frac{1}{t^3} \quad (4)$$

Let

$$\begin{aligned}w &= x^{1-n} \\ &= \frac{1}{x}\end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t t gives

$$w' = -\frac{1}{x^2}x' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} -w'(t) &= \frac{w(t)}{t} + \frac{1}{t^3} \\ w' &= -\frac{w}{t} - \frac{1}{t^3} \end{aligned} \tag{7}$$

The above now is a linear ODE in $w(t)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(t) + p(t)w(t) = q(t)$$

Where here

$$\begin{aligned} p(t) &= \frac{1}{t} \\ q(t) &= -\frac{1}{t^3} \end{aligned}$$

Hence the ode is

$$w'(t) + \frac{w(t)}{t} = -\frac{1}{t^3}$$

The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int \frac{1}{t} dt} \\ &= t \end{aligned}$$

The ode becomes

$$\begin{aligned} \frac{d}{dt}(\mu w) &= (\mu) \left(-\frac{1}{t^3} \right) \\ \frac{d}{dt}(wt) &= (t) \left(-\frac{1}{t^3} \right) \\ d(wt) &= \left(-\frac{1}{t^2} \right) dt \end{aligned}$$

Integrating gives

$$\begin{aligned} wt &= \int -\frac{1}{t^2} dt \\ wt &= \frac{1}{t} + c_1 \end{aligned}$$

Dividing both sides by the integrating factor $\mu = t$ results in

$$w(t) = \frac{1}{t^2} + \frac{c_1}{t}$$

Replacing w in the above by $\frac{1}{x}$ using equation (5) gives the final solution.

$$\frac{1}{x} = \frac{1}{t^2} + \frac{c_1}{t}$$

Or

$$x = \frac{1}{\frac{1}{t^2} + \frac{c_1}{t}}$$

Initial conditions are used to solve for c_1 . Substituting $t = 2$ and $x = 4$ in the above solution gives an equation to solve for the constant of integration.

$$4 = \frac{4}{2c_1 + 1}$$

$$c_1 = 0$$


Substituting c_1 found above in the general solution gives

$$x = t^2$$


Summary

The solution(s) found are the following

$$x = t^2 \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$x = t^2$$

Verified OK.

1.32.5 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(t, x) dt + N(t, x) dx = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (t^3) dx &= (x(t^2 + x)) dt \\ (-x(t^2 + x)) dt + (t^3) dx &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(t, x) &= -x(t^2 + x) \\ N(t, x) &= t^3 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial x} = \frac{\partial N}{\partial t}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial x} &= \frac{\partial}{\partial x}(-x(t^2 + x)) \\ &= -t^2 - 2x\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial t} &= \frac{\partial}{\partial t}(t^3) \\ &= 3t^2\end{aligned}$$

Since $\frac{\partial M}{\partial x} \neq \frac{\partial N}{\partial t}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial x} - \frac{\partial N}{\partial t} \right) \\ &= \frac{1}{t^3} ((-t^2 - 2x) - (3t^2)) \\ &= \frac{-4t^2 - 2x}{t^3}\end{aligned}$$

Since A depends on x , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned}B &= \frac{1}{M} \left(\frac{\partial N}{\partial t} - \frac{\partial M}{\partial x} \right) \\ &= -\frac{1}{x(t^2 + x)} ((3t^2) - (-t^2 - 2x)) \\ &= \frac{-4t^2 - 2x}{x(t^2 + x)}\end{aligned}$$

Since B depends on t , it can not be used to obtain an integrating factor. We will now try a third method to find an integrating factor. Let

$$R = \frac{\frac{\partial N}{\partial t} - \frac{\partial M}{\partial x}}{xM - yN}$$

R is now checked to see if it is a function of only $t = tx$. Therefore

$$\begin{aligned} R &= \frac{\frac{\partial N}{\partial t} - \frac{\partial M}{\partial x}}{xM - yN} \\ &= \frac{(3t^2) - (-t^2 - 2x)}{t(-x(t^2 + x)) - x(t^3)} \\ &= -\frac{2}{tx} \end{aligned}$$

Replacing all powers of terms tx by t gives

$$R = -\frac{2}{t}$$

Since R depends on t only, then it can be used to find an integrating factor. Let the integrating factor be μ then

$$\begin{aligned} \mu &= e^{\int R dt} \\ &= e^{\int (-\frac{2}{t}) dt} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-2 \ln(t)} \\ &= \frac{1}{t^2} \end{aligned}$$

Now t is replaced back with tx giving

$$\mu = \frac{1}{t^2 x^2}$$

Multiplying M and N by this integrating factor gives new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N

$$\begin{aligned} \bar{M} &= \mu M \\ &= \frac{1}{t^2 x^2} (-x(t^2 + x)) \\ &= \frac{-t^2 - x}{t^2 x} \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= \frac{1}{t^2 x^2} (t^3) \\ &= \frac{t}{x^2} \end{aligned}$$

A modified ODE is now obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \overline{M} + \overline{N} \frac{dx}{dt} &= 0 \\ \left(\frac{-t^2 - x}{t^2 x} \right) + \left(\frac{t}{x^2} \right) \frac{dx}{dt} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(t, x)$

$$\frac{\partial \phi}{\partial t} = \overline{M} \quad (1)$$

$$\frac{\partial \phi}{\partial x} = \overline{N} \quad (2)$$

Integrating (1) w.r.t. t gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial t} dt &= \int \overline{M} dt \\ \int \frac{\partial \phi}{\partial t} dt &= \int \frac{-t^2 - x}{t^2 x} dt \\ \phi &= \frac{-t^2 + x}{tx} + f(x) \end{aligned} \quad (3)$$

Where $f(x)$ is used for the constant of integration since ϕ is a function of both t and x . Taking derivative of equation (3) w.r.t x gives

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= \frac{1}{tx} - \frac{-t^2 + x}{tx^2} + f'(x) \\ &= \frac{t}{x^2} + f'(x) \end{aligned} \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial x} = \frac{t}{x^2}$. Therefore equation (4) becomes

$$\frac{t}{x^2} = \frac{t}{x^2} + f'(x) \quad (5)$$

Solving equation (5) for $f'(x)$ gives

$$f'(x) = 0$$

Therefore

$$f(x) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(x)$ into equation (3) gives ϕ

$$\phi = \frac{-t^2 + x}{tx} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{-t^2 + x}{tx}$$

The solution becomes

$$x = -\frac{t^2}{c_1 t - 1}$$

Initial conditions are used to solve for c_1 . Substituting $t = 2$ and $x = 4$ in the above solution gives an equation to solve for the constant of integration.

$$4 = -\frac{4}{2c_1 - 1}$$

$$c_1 = 0$$


Substituting c_1 found above in the general solution gives

$$x = t^2$$


Summary

The solution(s) found are the following

$$x = t^2 \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$x = t^2$$

Verified OK.

1.32.6 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned} x' &= F(t, x) \\ &= \frac{x(t^2 + x)}{t^3} \end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$x' = \frac{x}{t} + \frac{x^2}{t^3}$$

With Riccati ODE standard form

$$x' = f_0(t) + f_1(t)x + f_2(t)x^2$$

Shows that $f_0(t) = 0$, $f_1(t) = \frac{1}{t}$ and $f_2(t) = \frac{1}{t^3}$. Let

$$\begin{aligned} x &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{\frac{u}{t^3}} \end{aligned} \tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(t) - (f_2' + f_1 f_2) u'(t) + f_2^2 f_0 u(t) = 0 \quad (2)$$

But

$$\begin{aligned} f_2' &= -\frac{3}{t^4} \\ f_1 f_2 &= \frac{1}{t^4} \\ f_2^2 f_0 &= 0 \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$\frac{u''(t)}{t^3} + \frac{2u'(t)}{t^4} = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(t) = c_1 + \frac{c_2}{t}$$

The above shows that

$$u'(t) = -\frac{c_2}{t^2}$$

Using the above in (1) gives the solution

$$x = \frac{t c_2}{c_1 + \frac{c_2}{t}}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$x = \frac{t}{c_3 + \frac{1}{t}}$$

Initial conditions are used to solve for c_3 . Substituting $t = 2$ and $x = 4$ in the above solution gives an equation to solve for the constant of integration.

$$4 = \frac{4}{2c_3 + 1}$$

$$c_3 = 0$$


Substituting c_3 found above in the general solution gives

$$x = t^2$$


Summary

The solution(s) found are the following

$$x = t^2 \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$x = t^2$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 7

```
dsolve([diff(x(t),t)=x(t)/t+x(t)^2/t^3,x(2) = 4],x(t), singsol=all)
```

$$x(t) = t^2$$

✓ Solution by Mathematica

Time used: 0.264 (sec). Leaf size: 8

```
DSolve[{x'[t]==x[t]/t+x[t]^2/t^3,{x[2]==4}],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow t^2$$

1.33 problem Problem 47

1.33.1 Solving as clairaut ode 312

Internal problem ID [12144]

Internal file name [OUTPUT/10796_Tuesday_September_12_2023_08_53_33_AM_23258082/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 47.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "**clairaut**"

Maple gives the following as the ode type

```
[[_1st_order , _with_linear_symmetries], _Clairaut]
```

$$y - y'x - y'^2 = 0$$

With initial conditions

$$[y(2) = -1]$$

1.33.1 Solving as clairaut ode

This is Clairaut ODE. It has the form

$$y = y'x + g(y')$$

Where g is function of $y'(x)$. Let $p = y'$ the ode becomes

$$-p^2 - px + y = 0$$

Solving for y from the above results in

$$y = p^2 + px \tag{1A}$$

The above ode is a Clairaut ode which is now solved. We start by replacing y' by p which gives

$$\begin{aligned}y &= p^2 + px \\ &= p^2 + px\end{aligned}$$

Writing the ode as

$$y = px + g(p)$$

We now write $g \equiv g(p)$ to make notation simpler but we should always remember that g is function of p which in turn is function of x . Hence the above becomes

$$y = px + g \tag{1}$$

Then we see that

$$g = p^2$$

Taking derivative of (1) w.r.t. x gives

$$\begin{aligned}p &= \frac{d}{dx}(xp + g) \\ p &= \left(p + x \frac{dp}{dx}\right) + \left(g' \frac{dp}{dx}\right) \\ p &= p + (x + g') \frac{dp}{dx} \\ 0 &= (x + g') \frac{dp}{dx}\end{aligned}$$

Where g' is derivative of $g(p)$ w.r.t. p . The general solution is given by

$$\begin{aligned}\frac{dp}{dx} &= 0 \\ p &= c_1\end{aligned}$$

Substituting this in (1) gives the general solution as

$$y = c_1^2 + c_1x$$

The singular solution is found from solving for p from

$$x + g'(p) = 0$$

And substituting the result back in (1). Since we found above that $g = p^2$, then the above equation becomes

$$\begin{aligned}x + g'(p) &= x + 2p \\ &= 0\end{aligned}$$

Solving the above for p results in

$$p_1 = -\frac{x}{2}$$

Substituting the above back in (1) results in

$$y_1 = -\frac{x^2}{4}$$

Initial conditions are used to solve for c_1 . Substituting $x = 2$ and $y = -1$ in the above solution gives an equation to solve for the constant of integration.

$$-1 = c_1^2 + 2c_1$$

$$c_1 = -1$$

Substituting c_1 found above in the general solution gives

$$y = 1 - x$$

Summary

The solution(s) found are the following

$$y = 1 - x \tag{1}$$

$$y = -\frac{x^2}{4} \tag{2}$$




Figure 63: Solution plot

Verification of solutions

$$y = 1 - x$$

Verified OK.

$$y = -\frac{x^2}{4}$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
  *** Sublevel 2 ***
  Methods for first order ODEs:
  -> Solving 1st order ODE of high degree, 1st attempt
  trying 1st order WeierstrassP solution for high degree ODE
  trying 1st order WeierstrassPPrime solution for high degree ODE
  trying 1st order JacobiSN solution for high degree ODE
  trying 1st order ODE linearizable_by_differentiation
  trying differential order: 1; missing variables
  trying dAlembert
  <- dAlembert successful`

```

✓ Solution by Maple

Time used: 1.422 (sec). Leaf size: 17

```
dsolve([y(x)=x*diff(y(x),x)+diff(y(x),x)^2,y(2) = -1],y(x), singsol=all)
```

$$y(x) = 1 - x$$
$$y(x) = -\frac{x^2}{4}$$

✓ Solution by Mathematica

Time used: 0.014 (sec). Leaf size: 21

```
DSolve[{y[x]==x*y'[x]+y'[x]^2,{y[2]==-1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow 1 - x$$
$$y(x) \rightarrow -\frac{x^2}{4}$$

1.34 problem Problem 48

1.34.1 Solving as clairaut ode 317

Internal problem ID [12145]

Internal file name [OUTPUT/10797_Thursday_September_21_2023_05_46_02_AM_75656588/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 48.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "**clairaut**"

Maple gives the following as the ode type

```
[[_1st_order , _with_linear_symmetries], _Clairaut]
```

$$y - y'x - y'^2 = 0$$

With initial conditions

$$[y(1) = -1]$$

1.34.1 Solving as clairaut ode

This is Clairaut ODE. It has the form

$$y = y'x + g(y')$$

Where g is function of $y'(x)$. Let $p = y'$ the ode becomes

$$-p^2 - px + y = 0$$

Solving for y from the above results in

$$y = p^2 + px \tag{1A}$$

The above ode is a Clairaut ode which is now solved. We start by replacing y' by p which gives

$$\begin{aligned}y &= p^2 + px \\ &= p^2 + px\end{aligned}$$

Writing the ode as

$$y = px + g(p)$$

We now write $g \equiv g(p)$ to make notation simpler but we should always remember that g is function of p which in turn is function of x . Hence the above becomes

$$y = px + g \tag{1}$$

Then we see that

$$g = p^2$$

Taking derivative of (1) w.r.t. x gives

$$\begin{aligned}p &= \frac{d}{dx}(xp + g) \\ p &= \left(p + x \frac{dp}{dx}\right) + \left(g' \frac{dp}{dx}\right) \\ p &= p + (x + g') \frac{dp}{dx} \\ 0 &= (x + g') \frac{dp}{dx}\end{aligned}$$

Where g' is derivative of $g(p)$ w.r.t. p . The general solution is given by

$$\begin{aligned}\frac{dp}{dx} &= 0 \\ p &= c_1\end{aligned}$$

Substituting this in (1) gives the general solution as

$$y = c_1^2 + c_1x$$

The singular solution is found from solving for p from

$$x + g'(p) = 0$$

And substituting the result back in (1). Since we found above that $g = p^2$, then the above equation becomes

$$\begin{aligned}x + g'(p) &= x + 2p \\ &= 0\end{aligned}$$

Solving the above for p results in

$$p_1 = -\frac{x}{2}$$

Substituting the above back in (1) results in

$$y_1 = -\frac{x^2}{4}$$

Initial conditions are used to solve for c_1 . Substituting $x = 1$ and $y = -1$ in the above solution gives an equation to solve for the constant of integration.

$$-1 = c_1^2 + c_1$$

$$c_1 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

Substituting c_1 found above in the general solution gives

$$y = -\frac{1}{2} + \frac{i\sqrt{3}}{2} - \frac{x}{2} - \frac{i\sqrt{3}x}{2}$$

Summary

The solution(s) found are the following

$$y = -\frac{1}{2} + \frac{i\sqrt{3}}{2} - \frac{x}{2} - \frac{i\sqrt{3}x}{2} \quad (1)$$

$$y = -\frac{x^2}{4} \quad (2)$$




Figure 64: Solution plot

Verification of solutions

$$y = -\frac{1}{2} + \frac{i\sqrt{3}}{2} - \frac{x}{2} - \frac{i\sqrt{3}x}{2}$$

Verified OK.

$$y = -\frac{x^2}{4}$$

Warning, solution could not be verified

Maple trace

```

`Methods for first order ODEs:
  *** Sublevel 2 ***
  Methods for first order ODEs:
  -> Solving 1st order ODE of high degree, 1st attempt
  trying 1st order WeierstrassP solution for high degree ODE
  trying 1st order WeierstrassPPrime solution for high degree ODE
  trying 1st order JacobiSN solution for high degree ODE
  trying 1st order ODE linearizable_by_differentiation
  trying differential order: 1; missing variables
  trying dAlembert
  <- dAlembert successful`

```

✓ Solution by Maple

Time used: 0.391 (sec). Leaf size: 66

```
dsolve([y(x)=x*diff(y(x),x)+diff(y(x),x)^2,y(1) = -1],y(x), singsol=all)
```

$$y(x) = -\frac{1}{2} + \frac{i(-1+x)\sqrt{3}}{2} - \frac{x}{2}$$
$$y(x) = \frac{(1+i\sqrt{3})(i\sqrt{3}-2x+1)}{4}$$
$$y(x) = \frac{(i\sqrt{3}-1)(i\sqrt{3}+2x-1)}{4}$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 38

```
DSolve[{y[x]==x*y'[x]+y'[x]^2,{y[1]==-1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (-1)^{2/3} - \sqrt[3]{-1}x$$
$$y(x) \rightarrow \sqrt[3]{-1}(\sqrt[3]{-1}x - 1)$$

1.35 problem Problem 49

1.35.1 Solving as first order ode lie symmetry calculated ode 322

Internal problem ID [12146]

Internal file name [OUTPUT/10798_Thursday_September_21_2023_05_46_04_AM_77594040/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 49.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"first_order_ode_lie_symmetry_calculated"**

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$y' - \frac{3x - 4y - 2}{3x - 4y - 3} = 0$$

1.35.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{-3x + 4y + 2}{-3x + 4y + 3}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (\text{1E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 + \frac{(-3x + 4y + 2)(b_3 - a_2)}{-3x + 4y + 3} - \frac{(-3x + 4y + 2)^2 a_3}{(-3x + 4y + 3)^2} \\ - \left(-\frac{3}{-3x + 4y + 3} + \frac{-9x + 12y + 6}{(-3x + 4y + 3)^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(\frac{4}{-3x + 4y + 3} - \frac{4(-3x + 4y + 2)}{(-3x + 4y + 3)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{9x^2a_2 + 9x^2a_3 - 9x^2b_2 - 9x^2b_3 - 24xya_2 - 24xya_3 + 24xyb_2 + 24xyb_3 + 16y^2a_2 + 16y^2a_3 - 16y^2b_2 - 16y^2b_3 - 18xa_2 - 12xa_3 + 22xb_2 + 15xb_3 - 20ya_2 - 13ya_3 + 24yb_2 + 16yb_3 + 3a_1 - 6a_2 - 4a_3 - 4b_1 + 9b_2 + 6b_3}{(3x + 4y + 3)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} -9x^2a_2 - 9x^2a_3 + 9x^2b_2 + 9x^2b_3 + 24xya_2 + 24xya_3 - 24xyb_2 - 24xyb_3 \\ - 16y^2a_2 - 16y^2a_3 + 16y^2b_2 + 16y^2b_3 + 18xa_2 + 12xa_3 - 22xb_2 - 15xb_3 \\ - 20ya_2 - 13ya_3 + 24yb_2 + 16yb_3 + 3a_1 - 6a_2 - 4a_3 - 4b_1 + 9b_2 + 6b_3 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} -9a_2v_1^2 + 24a_2v_1v_2 - 16a_2v_2^2 - 9a_3v_1^2 + 24a_3v_1v_2 - 16a_3v_2^2 + 9b_2v_1^2 - 24b_2v_1v_2 \\ + 16b_2v_2^2 + 9b_3v_1^2 - 24b_3v_1v_2 + 16b_3v_2^2 + 18a_2v_1 - 20a_2v_2 + 12a_3v_1 - 13a_3v_2 \\ - 22b_2v_1 + 24b_2v_2 - 15b_3v_1 + 16b_3v_2 + 3a_1 - 6a_2 - 4a_3 - 4b_1 + 9b_2 + 6b_3 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned} &(-9a_2 - 9a_3 + 9b_2 + 9b_3) v_1^2 + (24a_2 + 24a_3 - 24b_2 - 24b_3) v_1 v_2 \\ &+ (18a_2 + 12a_3 - 22b_2 - 15b_3) v_1 + (-16a_2 - 16a_3 + 16b_2 + 16b_3) v_2^2 \\ &+ (-20a_2 - 13a_3 + 24b_2 + 16b_3) v_2 + 3a_1 - 6a_2 - 4a_3 - 4b_1 + 9b_2 + 6b_3 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} -20a_2 - 13a_3 + 24b_2 + 16b_3 &= 0 \\ -16a_2 - 16a_3 + 16b_2 + 16b_3 &= 0 \\ -9a_2 - 9a_3 + 9b_2 + 9b_3 &= 0 \\ 18a_2 + 12a_3 - 22b_2 - 15b_3 &= 0 \\ 24a_2 + 24a_3 - 24b_2 - 24b_3 &= 0 \\ 3a_1 - 6a_2 - 4a_3 - 4b_1 + 9b_2 + 6b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= a_1 \\ a_2 &= -9a_1 + 12b_1 \\ a_3 &= 12a_1 - 16b_1 \\ b_1 &= b_1 \\ b_2 &= -9a_1 + 12b_1 \\ b_3 &= 12a_1 - 16b_1 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= 12x - 16y \\ \eta &= 12x - 16y + 1 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}\eta &= \eta - \omega(x, y) \xi \\ &= 12x - 16y + 1 - \left(\frac{-3x + 4y + 2}{-3x + 4y + 3} \right) (12x - 16y) \\ &= \frac{-9x + 12y - 3}{3x - 4y - 3} \\ \xi &= 0\end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{-9x+12y-3}{3x-4y-3}} dy\end{aligned}$$

Which results in

$$S = -\frac{y}{3} - \frac{\ln(-1 - 3x + 4y)}{3}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y) S_y}{R_x + \omega(x, y) R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{-3x + 4y + 2}{-3x + 4y + 3}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{1}{-1 - 3x + 4y} \\ S_y &= -\frac{1}{3} + \frac{4}{9x - 12y + 3} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -\frac{1}{3} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{1}{3}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\frac{R}{3} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{y}{3} - \frac{\ln(-1 - 3x + 4y)}{3} = -\frac{x}{3} + c_1$$



Which simplifies to

$$-\frac{y}{3} - \frac{\ln(-1 - 3x + 4y)}{3} = -\frac{x}{3} + c_1$$

Which gives

$$y = \frac{3x}{4} + \text{LambertW}\left(\frac{e^{\frac{x}{4} - 3c_1 - \frac{1}{4}}}{4}\right) + \frac{1}{4}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{-3x+4y+2}{-3x+4y+3}$ 	$R = x$ $S = -\frac{y}{3} - \frac{\ln(-1 - 3x + 4y + 3)}{3}$	$\frac{dS}{dR} = -\frac{1}{3}$ 

Summary

The solution(s) found are the following

$$y = \frac{3x}{4} + \text{LambertW}\left(\frac{e^{\frac{x}{4}-3c_1-\frac{1}{4}}}{4}\right) + \frac{1}{4} \quad (1)$$




Figure 65: Slope field plot

Verification of solutions

$$y = \frac{3x}{4} + \text{LambertW}\left(\frac{e^{\frac{x}{4}-3c_1-\frac{1}{4}}}{4}\right) + \frac{1}{4}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous C
1st order, trying the canonical coordinates of the invariance group
  -> Calling odsolve with the ODE`, diff(y(x), x) = 3/4, y(x)` *** Sublevel 2 ***
      Methods for first order ODEs:
      --- Trying classification methods ---
      trying a quadrature
      trying 1st order linear
      <- 1st order linear successful
<- 1st order, canonical coordinates successful
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 19

```
dsolve(diff(y(x),x)=(3*x-4*y(x)-2)/(3*x-4*y(x)-3),y(x), singsol=all)
```

$$y(x) = \frac{3x}{4} + \text{LambertW}\left(\frac{c_1 e^{-\frac{1}{4} + \frac{x}{4}}}{4}\right) + \frac{1}{4}$$

✓ Solution by Mathematica

Time used: 5.353 (sec). Leaf size: 41

```
DSolve[y'[x]==(3*x-4*y[x]-2)/(3*x-4*y[x]-3),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow W(-e^{\frac{x}{4}-1+c_1}) + \frac{3x}{4} + \frac{1}{4}$$
$$y(x) \rightarrow \frac{1}{4}(3x + 1)$$

1.36 problem Problem 50

1.36.1 Solving as linear ode	330
1.36.2 Solving as first order ode lie symmetry lookup ode	332
1.36.3 Solving as exact ode	336
1.36.4 Maple step by step solution	340

Internal problem ID [12147]

Internal file name [OUTPUT/10799_Thursday_September_21_2023_05_46_05_AM_44492199/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 50.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"linear", "exactWithIntegrationFactor", "first_order_ode_lie_symmetry_lookup"**

Maple gives the following as the ode type

`[_linear]`

$$x' - x \cot(t) = 4 \sin(t)$$

1.36.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$x' + p(t)x = q(t)$$

Where here

$$p(t) = -\cot(t)$$

$$q(t) = 4 \sin(t)$$

Hence the ode is

$$x' - x \cot(t) = 4 \sin(t)$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int -\cot(t)dt} \\ &= \frac{1}{\sin(t)}\end{aligned}$$

Which simplifies to

$$\mu = \csc(t)$$

The ode becomes

$$\begin{aligned}\frac{d}{dt}(\mu x) &= (\mu)(4 \sin(t)) \\ \frac{d}{dt}(\csc(t) x) &= (\csc(t))(4 \sin(t)) \\ d(\csc(t) x) &= 4 dt\end{aligned}$$

Integrating gives

$$\begin{aligned}\csc(t) x &= \int 4 dt \\ \csc(t) x &= 4t + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = \csc(t)$ results in

$$x = 4 \sin(t) t + c_1 \sin(t)$$

which simplifies to

$$x = \sin(t) (4t + c_1)$$

Summary

The solution(s) found are the following

$$x = \sin(t) (4t + c_1) \tag{1}$$




Figure 66: Slope field plot

Verification of solutions

$$x = \sin(t) (4t + c_1)$$

Verified OK.

1.36.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$x' = x \cot(t) + 4 \sin(t)$$

$$x' = \omega(t, x)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_t + \omega(\eta_x - \xi_t) - \omega^2 \xi_x - \omega_t \xi - \omega_x \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 40: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(t, x) &= 0 \\ \eta(t, x) &= \sin(t)\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(t, x) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dt}{\xi} = \frac{dx}{\eta} = dS\tag{1}$$

The above comes from the requirements that $(\xi \frac{\partial}{\partial t} + \eta \frac{\partial}{\partial x})S(t, x) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = t$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\sin(t)} dy \end{aligned}$$

Which results in

$$S = \frac{x}{\sin(t)}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_t + \omega(t, x)S_x}{R_t + \omega(t, x)R_x} \quad (2)$$

Where in the above R_t, R_x, S_t, S_x are all partial derivatives and $\omega(t, x)$ is the right hand side of the original ode given by

$$\omega(t, x) = x \cot(t) + 4 \sin(t)$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_t &= 1 \\ R_x &= 0 \\ S_t &= -\cot(t) \csc(t) x \\ S_x &= \csc(t) \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 4 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for t, x in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 4$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = 4R + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to t, x coordinates. This results in

$$\csc(t) x = 4t + c_1$$



Which simplifies to

$$\csc(t) x = 4t + c_1$$

Which gives

$$x = \frac{4t + c_1}{\csc(t)}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in t, x coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dx}{dt} = x \cot(t) + 4 \sin(t)$ 	$R = t$ $S = \csc(t) x$	$\frac{dS}{dR} = 4$ 

Summary

The solution(s) found are the following

$$x = \frac{4t + c_1}{\csc(t)} \quad (1)$$




Figure 67: Slope field plot

Verification of solutions

$$x = \frac{4t + c_1}{\csc(t)}$$

Verified OK.

1.36.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(t, x) dt + N(t, x) dx = 0 \tag{1A}$$

Therefore

$$\begin{aligned}dx &= (x \cot(t) + 4 \sin(t)) dt \\ (-x \cot(t) - 4 \sin(t)) dt + dx &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(t, x) &= -x \cot(t) - 4 \sin(t) \\ N(t, x) &= 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial x} = \frac{\partial N}{\partial t}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial x} &= \frac{\partial}{\partial x}(-x \cot(t) - 4 \sin(t)) \\ &= -\cot(t)\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial t} &= \frac{\partial}{\partial t}(1) \\ &= 0\end{aligned}$$

Since $\frac{\partial M}{\partial x} \neq \frac{\partial N}{\partial t}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial x} - \frac{\partial N}{\partial t} \right) \\ &= 1((- \cot(t)) - (0)) \\ &= - \cot(t) \end{aligned}$$

Since A does not depend on x , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned} \mu &= e^{\int A dt} \\ &= e^{\int - \cot(t) dt} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-\ln(\sin(t))} \\ &= \csc(t) \end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \overline{M} and \overline{N} for now so not to confuse them with the original M and N .

$$\begin{aligned} \overline{M} &= \mu M \\ &= \csc(t) (-x \cot(t) - 4 \sin(t)) \\ &= -4 - \cot(t) \csc(t) x \end{aligned}$$

And

$$\begin{aligned} \overline{N} &= \mu N \\ &= \csc(t) (1) \\ &= \csc(t) \end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \overline{M} + \overline{N} \frac{dx}{dt} &= 0 \\ (-4 - \cot(t) \csc(t) x) + (\csc(t)) \frac{dx}{dt} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(t, x)$

$$\frac{\partial \phi}{\partial t} = \overline{M} \tag{1}$$

$$\frac{\partial \phi}{\partial x} = \overline{N} \tag{2}$$

Integrating (1) w.r.t. t gives

$$\int \frac{\partial \phi}{\partial t} dt = \int \overline{M} dt$$

$$\int \frac{\partial \phi}{\partial t} dt = \int -4 - \cot(t) \csc(t) x dt$$

$$\phi = -4t + \csc(t) x + f(x) \quad (3)$$

Where $f(x)$ is used for the constant of integration since ϕ is a function of both t and x . Taking derivative of equation (3) w.r.t x gives

$$\frac{\partial \phi}{\partial x} = \csc(t) + f'(x) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial x} = \csc(t)$. Therefore equation (4) becomes

$$\csc(t) = \csc(t) + f'(x) \quad (5)$$

Solving equation (5) for $f'(x)$ gives

$$f'(x) = 0$$

Therefore

$$f(x) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(x)$ into equation (3) gives ϕ

$$\phi = -4t + \csc(t) x + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -4t + \csc(t) x$$

The solution becomes

$$x = \frac{4t + c_1}{\csc(t)}$$

Summary

The solution(s) found are the following

$$x = \frac{4t + c_1}{\csc(t)} \quad (1)$$




Figure 68: Slope field plot

Verification of solutions

$$x = \frac{4t + c_1}{\csc(t)}$$

Verified OK.

1.36.4 Maple step by step solution

Let's solve

$$x' - x \cot(t) = 4 \sin(t)$$

- Highest derivative means the order of the ODE is 1

x'

- Isolate the derivative

$$x' = x \cot(t) + 4 \sin(t)$$

- Group terms with x on the lhs of the ODE and the rest on the rhs of the ODE

$$x' - x \cot(t) = 4 \sin(t)$$

- The ODE is linear; multiply by an integrating factor $\mu(t)$

$$\mu(t) (x' - x \cot(t)) = 4\mu(t) \sin(t)$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dt}(\mu(t) x)$

$$\mu(t) (x' - x \cot(t)) = \mu'(t) x + \mu(t) x'$$

- Isolate $\mu'(t)$

$$\mu'(t) = -\mu(t) \cot(t)$$

- Solve to find the integrating factor

$$\mu(t) = \frac{1}{\sin(t)}$$

- Integrate both sides with respect to t

$$\int \left(\frac{d}{dt}(\mu(t) x) \right) dt = \int 4\mu(t) \sin(t) dt + c_1$$

- Evaluate the integral on the lhs

$$\mu(t) x = \int 4\mu(t) \sin(t) dt + c_1$$

- Solve for x

$$x = \frac{\int 4\mu(t) \sin(t) dt + c_1}{\mu(t)}$$

- Substitute $\mu(t) = \frac{1}{\sin(t)}$

$$x = \sin(t) \left(\int 4 dt + c_1 \right)$$

- Evaluate the integrals on the rhs

$$x = \sin(t) (4t + c_1)$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
<- 1st order linear successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve(diff(x(t),t)-x(t)*cot(t)=4*sin(t),x(t), singsol=all)
```

$$x(t) = (4t + c_1) \sin(t)$$

✓ Solution by Mathematica

Time used: 0.073 (sec). Leaf size: 14

```
DSolve[x'[t]-x[t]*Cot[t]==4*Sin[t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow (4t + c_1) \sin(t)$$

1.37 problem Problem 51

Internal problem ID [12148]

Internal file name [OUTPUT/10800_Thursday_September_21_2023_05_46_06_AM_18357711/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 51.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_homogeneous, `class G`]]
```

$$y - 2y'x - \frac{y'^2}{2} = x^2$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = -2x + \sqrt{2x^2 + 2y} \quad (1)$$

$$y' = -2x - \sqrt{2x^2 + 2y} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Writing the ode as

$$y' = -2x + \sqrt{2x^2 + 2y}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (A)$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (1E)$$

$$\eta = xb_2 + yb_3 + b_1 \quad (2E)$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} & b_2 + \left(-2x + \sqrt{2x^2 + 2y}\right) (b_3 - a_2) - \left(-2x + \sqrt{2x^2 + 2y}\right)^2 a_3 \\ & - \left(-2 + \frac{2x}{\sqrt{2x^2 + 2y}}\right) (xa_2 + ya_3 + a_1) - \frac{xb_2 + yb_3 + b_1}{\sqrt{2x^2 + 2y}} = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{(2x^2 + 2y)^{\frac{3}{2}} a_3 + 4\sqrt{2x^2 + 2y} x^2 a_3 - 8x^3 a_3 - 4\sqrt{2x^2 + 2y} xa_2 + 2\sqrt{2x^2 + 2y} xb_3 - 2\sqrt{2x^2 + 2y} ya_3 + 4a_1 + 2\sqrt{2x^2 + 2y} b_1}{\sqrt{2x^2 + 2y}} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} & -(2x^2 + 2y)^{\frac{3}{2}} a_3 - 4\sqrt{2x^2 + 2y} x^2 a_3 + 8x^3 a_3 + 4\sqrt{2x^2 + 2y} xa_2 \\ & - 2\sqrt{2x^2 + 2y} xb_3 + 2\sqrt{2x^2 + 2y} ya_3 - 4x^2 a_2 + 2x^2 b_3 + 6xya_3 \\ & + 2\sqrt{2x^2 + 2y} a_1 + b_2\sqrt{2x^2 + 2y} - 2xa_1 - xb_2 - 2ya_2 + yb_3 - b_1 = 0 \end{aligned} \quad (6E)$$

Simplifying the above gives

$$\begin{aligned} & -(2x^2 + 2y)^{\frac{3}{2}} a_3 + 4(2x^2 + 2y) xa_3 - 4\sqrt{2x^2 + 2y} x^2 a_3 - (2x^2 + 2y) a_2 \\ & + (2x^2 + 2y) b_3 + 4\sqrt{2x^2 + 2y} xa_2 - 2\sqrt{2x^2 + 2y} xb_3 + 2\sqrt{2x^2 + 2y} ya_3 \\ & - 2x^2 a_2 - 2xya_3 + 2\sqrt{2x^2 + 2y} a_1 + b_2\sqrt{2x^2 + 2y} - 2xa_1 - xb_2 - yb_3 - b_1 = 0 \end{aligned} \quad (6E)$$

Since the PDE has radicals, simplifying gives

$$8x^3a_3 - 6\sqrt{2x^2 + 2y}x^2a_3 - 4x^2a_2 + 2x^2b_3 + 4\sqrt{2x^2 + 2y}xa_2 - 2\sqrt{2x^2 + 2y}xb_3 \\ + 6xya_3 - 2xa_1 - xb_2 + 2\sqrt{2x^2 + 2y}a_1 + b_2\sqrt{2x^2 + 2y} - 2ya_2 + yb_3 - b_1 = 0$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y, \sqrt{2x^2 + 2y}\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2, \sqrt{2x^2 + 2y} = v_3\}$$

The above PDE (6E) now becomes

$$8v_1^3a_3 - 6v_3v_1^2a_3 - 4v_1^2a_2 + 4v_3v_1a_2 + 6v_1v_2a_3 + 2v_1^2b_3 - 2v_3v_1b_3 \\ - 2v_1a_1 + 2v_3a_1 - 2v_2a_2 - v_1b_2 + b_2v_3 + v_2b_3 - b_1 = 0 \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3\}$$

Equation (7E) now becomes

$$8v_1^3a_3 - 6v_3v_1^2a_3 + (-4a_2 + 2b_3)v_1^2 + 6v_1v_2a_3 + (4a_2 - 2b_3)v_1v_3 \\ + (-2a_1 - b_2)v_1 + (-2a_2 + b_3)v_2 + (2a_1 + b_2)v_3 - b_1 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} -6a_3 &= 0 \\ 6a_3 &= 0 \\ 8a_3 &= 0 \\ -b_1 &= 0 \\ -2a_1 - b_2 &= 0 \\ 2a_1 + b_2 &= 0 \\ -4a_2 + 2b_3 &= 0 \\ -2a_2 + b_3 &= 0 \\ 4a_2 - 2b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned}
 a_1 &= a_1 \\
 a_2 &= a_2 \\
 a_3 &= 0 \\
 b_1 &= 0 \\
 b_2 &= -2a_1 \\
 b_3 &= 2a_2
 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned}
 \xi &= 1 \\
 \eta &= -2x
 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}
 \eta &= \eta - \omega(x, y) \xi \\
 &= -2x - \left(-2x + \sqrt{2x^2 + 2y}\right) (1) \\
 &= -\sqrt{2x^2 + 2y} \\
 \xi &= 0
 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}
 S &= \int \frac{1}{\eta} dy \\
 &= \int \frac{1}{-\sqrt{2x^2 + 2y}} dy
 \end{aligned}$$

Which results in

$$S = -\sqrt{2x^2 + 2y}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -2x + \sqrt{2x^2 + 2y}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{2x}{\sqrt{2x^2 + 2y}} \\ S_y &= -\frac{1}{\sqrt{2x^2 + 2y}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -1 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -1$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -R + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\sqrt{2x^2 + 2y} = -x + c_1$$

Which simplifies to

$$-\sqrt{2x^2 + 2y} = -x + c_1$$

Which gives

$$y = \frac{1}{2}c_1^2 - c_1x - \frac{1}{2}x^2$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2}c_1^2 - c_1x - \frac{1}{2}x^2 \quad (1)$$

Verification of solutions

$$y = \frac{1}{2}c_1^2 - c_1x - \frac{1}{2}x^2$$

Verified OK.

Solving equation (2)

Writing the ode as

$$y' = -2x - \sqrt{2x^2 + 2y}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2\xi_y - \omega_x\xi - \omega_y\eta = 0 \quad (A)$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (1E)$$

$$\eta = xb_2 + yb_3 + b_1 \quad (2E)$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$b_2 + \left(-2x - \sqrt{2x^2 + 2y}\right) (b_3 - a_2) - \left(-2x - \sqrt{2x^2 + 2y}\right)^2 a_3$$

$$- \left(-2 - \frac{2x}{\sqrt{2x^2 + 2y}}\right) (xa_2 + ya_3 + a_1) + \frac{xb_2 + yb_3 + b_1}{\sqrt{2x^2 + 2y}} = 0 \quad (5E)$$

Putting the above in normal form gives

$$\frac{(2x^2 + 2y)^{\frac{3}{2}} a_3 + 4\sqrt{2x^2 + 2y} x^2 a_3 + 8x^3 a_3 - 4\sqrt{2x^2 + 2y} x a_2 + 2\sqrt{2x^2 + 2y} x b_3 - 2\sqrt{2x^2 + 2y} y a_3 - 4x^2 a_2 - 2x^2 b_3 - 6xy a_3}{\sqrt{2x^2 + 2y}} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} & -(2x^2 + 2y)^{\frac{3}{2}} a_3 - 4\sqrt{2x^2 + 2y} x^2 a_3 - 8x^3 a_3 + 4\sqrt{2x^2 + 2y} x a_2 \\ & - 2\sqrt{2x^2 + 2y} x b_3 + 2\sqrt{2x^2 + 2y} y a_3 + 4x^2 a_2 - 2x^2 b_3 - 6xy a_3 \\ & + 2\sqrt{2x^2 + 2y} a_1 + b_2 \sqrt{2x^2 + 2y} + 2xa_1 + xb_2 + 2ya_2 - yb_3 + b_1 = 0 \end{aligned} \quad (6E)$$

Simplifying the above gives

$$\begin{aligned} & -(2x^2 + 2y)^{\frac{3}{2}} a_3 - 4(2x^2 + 2y) x a_3 - 4\sqrt{2x^2 + 2y} x^2 a_3 + (2x^2 + 2y) a_2 \\ & - (2x^2 + 2y) b_3 + 4\sqrt{2x^2 + 2y} x a_2 - 2\sqrt{2x^2 + 2y} x b_3 + 2\sqrt{2x^2 + 2y} y a_3 \\ & + 2x^2 a_2 + 2xy a_3 + 2\sqrt{2x^2 + 2y} a_1 + b_2 \sqrt{2x^2 + 2y} + 2xa_1 + xb_2 + yb_3 + b_1 = 0 \end{aligned} \quad (6E)$$

Since the PDE has radicals, simplifying gives

$$\begin{aligned} & -8x^3 a_3 - 6\sqrt{2x^2 + 2y} x^2 a_3 + 4x^2 a_2 - 2x^2 b_3 + 4\sqrt{2x^2 + 2y} x a_2 - 2\sqrt{2x^2 + 2y} x b_3 \\ & - 6xy a_3 + 2xa_1 + xb_2 + 2\sqrt{2x^2 + 2y} a_1 + b_2 \sqrt{2x^2 + 2y} + 2ya_2 - yb_3 + b_1 = 0 \end{aligned}$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\left\{ x, y, \sqrt{2x^2 + 2y} \right\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\left\{ x = v_1, y = v_2, \sqrt{2x^2 + 2y} = v_3 \right\}$$

The above PDE (6E) now becomes

$$\begin{aligned} & -8v_1^3 a_3 - 6v_3 v_1^2 a_3 + 4v_1^2 a_2 + 4v_3 v_1 a_2 - 6v_1 v_2 a_3 - 2v_1^2 b_3 - 2v_3 v_1 b_3 \\ & + 2v_1 a_1 + 2v_3 a_1 + 2v_2 a_2 + v_1 b_2 + b_2 v_3 - v_2 b_3 + b_1 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3\}$$

Equation (7E) now becomes

$$\begin{aligned} -8v_1^3 a_3 - 6v_3 v_1^2 a_3 + (4a_2 - 2b_3) v_1^2 - 6v_1 v_2 a_3 + (4a_2 - 2b_3) v_1 v_3 \\ + (2a_1 + b_2) v_1 + (2a_2 - b_3) v_2 + (2a_1 + b_2) v_3 + b_1 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} b_1 &= 0 \\ -8a_3 &= 0 \\ -6a_3 &= 0 \\ 2a_1 + b_2 &= 0 \\ 2a_2 - b_3 &= 0 \\ 4a_2 - 2b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= a_1 \\ a_2 &= a_2 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= -2a_1 \\ b_3 &= 2a_2 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= 1 \\ \eta &= -2x \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= -2x - \left(-2x - \sqrt{2x^2 + 2y}\right) (1) \\ &= \sqrt{2x^2 + 2y} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\sqrt{2x^2 + 2y}} dy \end{aligned}$$

Which results in

$$S = \sqrt{2x^2 + 2y}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -2x - \sqrt{2x^2 + 2y}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{2x}{\sqrt{2x^2 + 2y}} \\ S_y &= \frac{1}{\sqrt{2x^2 + 2y}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -1 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -1$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -R + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\sqrt{2x^2 + 2y} = -x + c_1$$

Which simplifies to

$$\sqrt{2x^2 + 2y} = -x + c_1$$

Which gives

$$y = \frac{1}{2}c_1^2 - c_1x - \frac{1}{2}x^2$$

Summary

The solution(s) found are the following

$$y = \frac{1}{2}c_1^2 - c_1x - \frac{1}{2}x^2 \quad (1)$$

Verification of solutions

$$y = \frac{1}{2}c_1^2 - c_1x - \frac{1}{2}x^2$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:
  *** Sublevel 2 ***
  Methods for first order ODEs:
  -> Solving 1st order ODE of high degree, 1st attempt
  trying 1st order WeierstrassP solution for high degree ODE
  trying 1st order WeierstrassPPrime solution for high degree ODE
  trying 1st order JacobiSN solution for high degree ODE
  trying 1st order ODE linearizable_by_differentiation
  trying differential order: 1; missing variables
  trying dAlembert
  trying simple symmetries for implicit equations
  Successful isolation of dy/dx: 2 solutions were found. Trying to solve each resulting ODE
    *** Sublevel 3 ***
    Methods for first order ODEs:
    --- Trying classification methods ---
    trying homogeneous types:
    trying homogeneous G
    1st order, trying the canonical coordinates of the invariance group
    <- 1st order, canonical coordinates successful
    <- homogeneous successful
    -----
  * Tackling next ODE.
    *** Sublevel 3 ***
    Methods for first order ODEs:
    --- Trying classification methods ---
    trying homogeneous types:
    trying homogeneous G
    1st order, trying the canonical coordinates of the invariance group
    <- 1st order, canonical coordinates successful
    <- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 79

```
dsolve(y(x)=x^2+2*diff(y(x),x)*x+(diff(y(x),x)^2)/2,y(x), singsol=all)
```

$$y(x) = -x^2$$

$$y(x) = -\frac{1}{2}x^2 + c_1x + \frac{1}{2}c_1^2$$

$$y(x) = -\frac{1}{2}x^2 - c_1x + \frac{1}{2}c_1^2$$

$$y(x) = -\frac{1}{2}x^2 - c_1x + \frac{1}{2}c_1^2$$

$$y(x) = -\frac{1}{2}x^2 + c_1x + \frac{1}{2}c_1^2$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y[x]==x^2+2*y'[x]*x+(y'[x]^2)/2,y[x],x,IncludeSingularSolutions -> True]
```

Timed out

1.38 problem Problem 52

1.38.1 Solving as first order ode lie symmetry lookup ode	355
1.38.2 Solving as bernoulli ode	359
1.38.3 Solving as riccati ode	363

Internal problem ID [12149]

Internal file name [OUTPUT/10801_Thursday_September_21_2023_05_46_07_AM_93917829/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 52.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "riccati", "bernoulli", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _rational, _Bernoulli]
```

$$y' - \frac{3y}{x} + y^2 x^3 = 0$$

1.38.1 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{y(x^4 y - 3)}{x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is known. It is of type **Bernoulli**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 43: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{y^2}{x^3}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{y^2}{x^3}} dy \end{aligned}$$

Which results in

$$S = -\frac{x^3}{y}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{y(x^4y - 3)}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{3x^2}{y} \\ S_y &= \frac{x^3}{y^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -x^6 \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -R^6$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\frac{R^7}{7} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{x^3}{y} = -\frac{x^7}{7} + c_1$$



Which simplifies to

$$-\frac{x^3}{y} = -\frac{x^7}{7} + c_1$$

Which gives

$$y = -\frac{7x^3}{-x^7 + 7c_1}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{y(x^4y-3)}{x}$ 	$R = x$ $S = -\frac{x^3}{y}$	$\frac{dS}{dR} = -R^6$ 

Summary

The solution(s) found are the following

$$y = -\frac{7x^3}{-x^7 + 7c_1} \quad (1)$$




Figure 69: Slope field plot

Verification of solutions

$$y = -\frac{7x^3}{-x^7 + 7c_1}$$

Verified OK.

1.38.2 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= -\frac{y(x^4y - 3)}{x} \end{aligned}$$

This is a Bernoulli ODE.

$$y' = \frac{3}{x}y - x^3y^2 \quad (1)$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \quad (2)$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \quad (3)$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned} f_0(x) &= \frac{3}{x} \\ f_1(x) &= -x^3 \\ n &= 2 \end{aligned}$$

Dividing both sides of ODE (1) by $y^n = y^2$ gives

$$y' \frac{1}{y^2} = \frac{3}{xy} - x^3 \quad (4)$$

Let

$$\begin{aligned} w &= y^{1-n} \\ &= \frac{1}{y} \end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = -\frac{1}{y^2}y' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} -w'(x) &= \frac{3w(x)}{x} - x^3 \\ w' &= -\frac{3w}{x} + x^3 \end{aligned} \quad (7)$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$p(x) = \frac{3}{x}$$
$$q(x) = x^3$$

Hence the ode is

$$w'(x) + \frac{3w(x)}{x} = x^3$$

The integrating factor μ is

$$\mu = e^{\int \frac{3}{x} dx}$$
$$= x^3$$

The ode becomes

$$\frac{d}{dx}(\mu w) = (\mu) (x^3)$$
$$\frac{d}{dx}(x^3 w) = (x^3) (x^3)$$
$$d(x^3 w) = x^6 dx$$

Integrating gives

$$x^3 w = \int x^6 dx$$
$$x^3 w = \frac{x^7}{7} + c_1$$

Dividing both sides by the integrating factor $\mu = x^3$ results in

$$w(x) = \frac{x^4}{7} + \frac{c_1}{x^3}$$

Replacing w in the above by $\frac{1}{y}$ using equation (5) gives the final solution.

$$\frac{1}{y} = \frac{x^4}{7} + \frac{c_1}{x^3}$$

Or

$$y = \frac{1}{\frac{x^4}{7} + \frac{c_1}{x^3}}$$

Which is simplified to

$$y = \frac{7x^3}{x^7 + 7c_1}$$

Summary

The solution(s) found are the following

$$y = \frac{7x^3}{x^7 + 7c_1} \tag{1}$$




Figure 70: Slope field plot

Verification of solutions

$$y = \frac{7x^3}{x^7 + 7c_1}$$

Verified OK.

1.38.3 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= -\frac{y(x^4y - 3)}{x}\end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = \frac{3y}{x} - y^2x^3$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = 0$, $f_1(x) = \frac{3}{x}$ and $f_2(x) = -x^3$. Let

$$\begin{aligned}y &= \frac{-u'}{f_2u} \\ &= \frac{-u'}{-x^3u}\end{aligned}\tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2u''(x) - (f_2' + f_1f_2)u'(x) + f_2^2f_0u(x) = 0\tag{2}$$

But

$$\begin{aligned}f_2' &= -3x^2 \\ f_1f_2 &= -3x^2 \\ f_2^2f_0 &= 0\end{aligned}$$

Substituting the above terms back in equation (2) gives

$$-x^3u''(x) + 6x^2u'(x) = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = c_2x^7 + c_1$$

The above shows that

$$u'(x) = 7c_2x^6$$

Using the above in (1) gives the solution

$$y = \frac{7c_2x^3}{c_2x^7 + c_1}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = \frac{7x^3}{x^7 + c_3}$$

Summary

The solution(s) found are the following

$$y = \frac{7x^3}{x^7 + c_3} \tag{1}$$




Figure 71: Slope field plot

Verification of solutions

$$y = \frac{7x^3}{x^7 + c_3}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 18

```
dsolve(diff(y(x),x)-3*y(x)/x+x^3*y(x)^2=0,y(x), singsol=all)
```

$$y(x) = \frac{7x^3}{x^7 + 7c_1}$$

✓ Solution by Mathematica

Time used: 0.238 (sec). Leaf size: 25

```
DSolve[y'[x]-3*y[x]/x+x^3*y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{7x^3}{x^7 + 7c_1}$$
$$y(x) \rightarrow 0$$

1.39 problem Problem 53

1.39.1 Maple step by step solution 367

Internal problem ID [12150]

Internal file name [OUTPUT/10802_Thursday_September_21_2023_05_46_08_AM_55045757/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 53.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : **"quadrature"**

Maple gives the following as the ode type

[_quadrature]

$$y(y'^2 + 1) = a$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{\sqrt{-y(y-a)}}{y} \tag{1}$$

$$y' = -\frac{\sqrt{-y(y-a)}}{y} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{y}{\sqrt{-y(-a+y)}} dy = \int dx$$
$$-\sqrt{ay-y^2} + \frac{a \arctan\left(\frac{y-\frac{a}{2}}{\sqrt{ay-y^2}}\right)}{2} = x + c_1$$

Summary

The solution(s) found are the following

$$-\sqrt{ay - y^2} + \frac{a \arctan\left(\frac{y - \frac{a}{2}}{\sqrt{ay - y^2}}\right)}{2} = x + c_1 \quad (1)$$

Verification of solutions

$$-\sqrt{ay - y^2} + \frac{a \arctan\left(\frac{y - \frac{a}{2}}{\sqrt{ay - y^2}}\right)}{2} = x + c_1$$

Verified OK.

Solving equation (2)

Integrating both sides gives

$$\int -\frac{y}{\sqrt{-y(-a+y)}} dy = \int dx$$
$$\sqrt{ay - y^2} - \frac{a \arctan\left(\frac{y - \frac{a}{2}}{\sqrt{ay - y^2}}\right)}{2} = x + c_2$$

Summary

The solution(s) found are the following

$$\sqrt{ay - y^2} - \frac{a \arctan\left(\frac{y - \frac{a}{2}}{\sqrt{ay - y^2}}\right)}{2} = x + c_2 \quad (1)$$

Verification of solutions

$$\sqrt{ay - y^2} - \frac{a \arctan\left(\frac{y - \frac{a}{2}}{\sqrt{ay - y^2}}\right)}{2} = x + c_2$$

Verified OK.

1.39.1 Maple step by step solution

Let's solve

$$y(y'^2 + 1) = a$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'y}{\sqrt{-y(y-a)}} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'y}{\sqrt{-y(y-a)}} dx = \int 1 dx + c_1$$

- Evaluate integral

$$-\sqrt{ay - y^2} + \frac{a \arctan\left(\frac{y - \frac{a}{2}}{\sqrt{ay - y^2}}\right)}{2} = x + c_1$$

Maple trace

```

`Methods for first order ODEs:
*** Sublevel 2 ***
Methods for first order ODEs:
-> Solving 1st order ODE of high degree, 1st attempt
trying 1st order WeierstrassP solution for high degree ODE
trying 1st order WeierstrassPPrime solution for high degree ODE
trying 1st order JacobiSN solution for high degree ODE
trying 1st order ODE linearizable_by_differentiation
trying differential order: 1; missing variables
<- differential order: 1; missing x successful`

```

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 339

```
dsolve(y(x)*(1+diff(y(x),x)^2)=a,y(x), singsol=all)
```

$$y(x) = a$$

$$y(x) = \frac{(\text{RootOf}((\cos(_Z)a + _Za + 2c_1 - 2x)(-\cos(_Z)a + _Za + 2c_1 - 2x))a - 2x + 2c_1) \tan(\text{RootOf}(\dots))}{2} + \frac{a}{2}$$

$$y(x) = \frac{(-\text{RootOf}((\cos(_Z)a + _Za + 2c_1 - 2x)(-\cos(_Z)a + _Za + 2c_1 - 2x))a + 2x - 2c_1) \tan(\text{RootOf}(\dots))}{2} + \frac{a}{2}$$

$$y(x) = \frac{(\text{RootOf}((\cos(_Z)a - _Za + 2c_1 - 2x)(-\cos(_Z)a - _Za + 2c_1 - 2x))a + 2x - 2c_1) \tan(\text{RootOf}(\dots))}{2} + \frac{a}{2}$$

$$y(x) = \frac{(-\text{RootOf}((\cos(_Z)a - _Za + 2c_1 - 2x)(-\cos(_Z)a - _Za + 2c_1 - 2x))a - 2x + 2c_1) \tan(\text{RootOf}(\dots))}{2} + \frac{a}{2}$$

✓ Solution by Mathematica

Time used: 0.661 (sec). Leaf size: 106

```
DSolve[y[x]*(1+y'[x]^2)==a,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \text{InverseFunction} \left[a \arctan \left(\frac{\sqrt{\#1}}{\sqrt{a - \#1}} \right) - \sqrt{\#1} \sqrt{a - \#1} \& \right] [-x + c_1]$$

$$y(x) \rightarrow \text{InverseFunction} \left[a \arctan \left(\frac{\sqrt{\#1}}{\sqrt{a - \#1}} \right) - \sqrt{\#1} \sqrt{a - \#1} \& \right] [x + c_1]$$

$$y(x) \rightarrow a$$

1.40 problem Problem 54

1.40.1 Solving as exact ode 370

Internal problem ID [12151]

Internal file name [OUTPUT/10803_Thursday_September_21_2023_05_46_08_AM_83126452/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 54.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"exactWithIntegrationFactor"**

Maple gives the following as the ode type

`[_rational]`

$$-y + (x^2y^2 + x) y' = -x^2$$

1.40.1 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (x^2 y^2 + x) dy &= (-x^2 + y) dx \\ (x^2 - y) dx + (x^2 y^2 + x) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= x^2 - y \\ N(x, y) &= x^2 y^2 + x \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (x^2 - y) \\ &= -1 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (x^2 y^2 + x) \\ &= 2x y^2 + 1 \end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned} A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{x^2 y^2 + x} ((-1) - (2x y^2 + 1)) \\ &= -\frac{2}{x} \end{aligned}$$

Since A does not depend on y , then it can be used to find an integrating factor. The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int A \, dx} \\ &= e^{\int -\frac{2}{x} \, dx}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{-2\ln(x)} \\ &= \frac{1}{x^2}\end{aligned}$$

M and N are multiplied by this integrating factor, giving new M and new N which are called \overline{M} and \overline{N} for now so not to confuse them with the original M and N .

$$\begin{aligned}\overline{M} &= \mu M \\ &= \frac{1}{x^2}(x^2 - y) \\ &= \frac{x^2 - y}{x^2}\end{aligned}$$

And

$$\begin{aligned}\overline{N} &= \mu N \\ &= \frac{1}{x^2}(x^2 y^2 + x) \\ &= \frac{x y^2 + 1}{x}\end{aligned}$$

Now a modified ODE is obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned}\overline{M} + \overline{N} \frac{dy}{dx} &= 0 \\ \left(\frac{x^2 - y}{x^2}\right) + \left(\frac{x y^2 + 1}{x}\right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{x^2 - y}{x^2} dx \\ \phi &= x + \frac{y}{x} + f(y)\end{aligned}\quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = \frac{1}{x} + f'(y)\quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{xy^2+1}{x}$. Therefore equation (4) becomes

$$\frac{xy^2 + 1}{x} = \frac{1}{x} + f'(y)\quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = y^2$$

Integrating the above w.r.t y gives

$$\begin{aligned}\int f'(y) dy &= \int (y^2) dy \\ f(y) &= \frac{y^3}{3} + c_1\end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = x + \frac{y}{x} + \frac{y^3}{3} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = x + \frac{y}{x} + \frac{y^3}{3}$$

Summary

The solution(s) found are the following

$$x + \frac{y}{x} + \frac{y^3}{3} = c_1 \quad (1)$$




Figure 72: Slope field plot

Verification of solutions

$$x + \frac{y}{x} + \frac{y^3}{3} = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying Chini  
differential order: 1; looking for linear symmetries  
trying exact  
<- exact successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 346

```
dsolve((x^2-y(x))+x^2*y(x)^2+x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{2^{\frac{1}{3}} \left(-\frac{2^{\frac{1}{3}} \left(\left(-3c_1x - 3x^2 + \sqrt{\frac{9c_1^2x^3 + 18x^4c_1 + 9x^5 + 4}{x}} \right) x^2 \right)^{\frac{2}{3}}}{2} + x \right)}{\left(\left(-3c_1x - 3x^2 + \sqrt{\frac{9c_1^2x^3 + 18x^4c_1 + 9x^5 + 4}{x}} \right) x^2 \right)^{\frac{1}{3}} x}$$
$$y(x) = \frac{\left((1 + i\sqrt{3}) 2^{\frac{1}{3}} \left(\left(-3c_1x - 3x^2 + \sqrt{\frac{9c_1^2x^3 + 18x^4c_1 + 9x^5 + 4}{x}} \right) x^2 \right)^{\frac{2}{3}} + 2i\sqrt{3}x - 2x \right) 2^{\frac{1}{3}}}{4 \left(\left(-3c_1x - 3x^2 + \sqrt{\frac{9c_1^2x^3 + 18x^4c_1 + 9x^5 + 4}{x}} \right) x^2 \right)^{\frac{1}{3}} x}$$
$$y(x) = \frac{(i\sqrt{3} - 1) 2^{\frac{2}{3}} \left(\left(-3c_1x - 3x^2 + \sqrt{\frac{9c_1^2x^3 + 18x^4c_1 + 9x^5 + 4}{x}} \right) x^2 \right)^{\frac{2}{3}} + 2(1 + i\sqrt{3}) 2^{\frac{1}{3}} x}{4 \left(\left(-3c_1x - 3x^2 + \sqrt{\frac{9c_1^2x^3 + 18x^4c_1 + 9x^5 + 4}{x}} \right) x^2 \right)^{\frac{1}{3}} x}$$

✓ Solution by Mathematica

Time used: 56.22 (sec). Leaf size: 400

`DSolve[(x^2-y[x])+(x^2*y[x]^2+x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{-2\sqrt[3]{2}x + \left(-6x^4 + 6c_1x^3 + 2\sqrt{x^3(9x^5 - 18c_1x^4 + 9c_1^2x^3 + 4)}\right)^{2/3}}{2x\sqrt[3]{-3x^4 + 3c_1x^3 + \sqrt{x^3(9x^5 - 18c_1x^4 + 9c_1^2x^3 + 4)}}$$

$$y(x) \rightarrow \frac{i(\sqrt{3} + i) \left(-6x^4 + 6c_1x^3 + 2\sqrt{x^3(9x^5 - 18c_1x^4 + 9c_1^2x^3 + 4)}\right)^{2/3} + \sqrt[3]{2}(2 + 2i\sqrt{3})x}{4x\sqrt[3]{-3x^4 + 3c_1x^3 + \sqrt{x^3(9x^5 - 18c_1x^4 + 9c_1^2x^3 + 4)}}$$

$$y(x) \rightarrow \frac{(-1 - i\sqrt{3}) \left(-6x^4 + 6c_1x^3 + 2\sqrt{x^3(9x^5 - 18c_1x^4 + 9c_1^2x^3 + 4)}\right)^{2/3} + \sqrt[3]{2}(2 - 2i\sqrt{3})x}{4x\sqrt[3]{-3x^4 + 3c_1x^3 + \sqrt{x^3(9x^5 - 18c_1x^4 + 9c_1^2x^3 + 4)}}$$

1.41 problem Problem 55

1.41.1 Solving as first order ode lie symmetry calculated ode 377

Internal problem ID [12152]

Internal file name [OUTPUT/10804_Thursday_September_21_2023_05_46_09_AM_79245405/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 55.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : **"first_order_ode_lie_symmetry_calculated"**

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _rational]
```

$$3y^2 + 2y(y^2 - 3x)y' = x$$

1.41.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = -\frac{3y^2 - x}{2y(y^2 - 3x)}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (1\text{E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (2\text{E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 - \frac{(3y^2 - x)(b_3 - a_2)}{2y(y^2 - 3x)} - \frac{(3y^2 - x)^2 a_3}{4y^2(y^2 - 3x)^2} \\ - \left(\frac{1}{2y(y^2 - 3x)} - \frac{3(3y^2 - x)}{2y(y^2 - 3x)^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(-\frac{3}{y^2 - 3x} + \frac{3y^2 - x}{2y^2(y^2 - 3x)} + \frac{3y^2 - x}{(y^2 - 3x)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{-4y^6b_2 + 30xy^4b_2 - 6y^5a_2 + 12y^5b_3 - 24x^2y^2b_2 + 4xy^3a_2 - 8xy^3b_3 - 7y^4a_3 + 6y^4b_1 + 6x^3b_2 - 6x^2ya_2}{4y^2(-y^2 + 3x)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} 4y^6b_2 - 30xy^4b_2 + 6y^5a_2 - 12y^5b_3 + 24x^2y^2b_2 - 4xy^3a_2 + 8xy^3b_3 + 7y^4a_3 - 6y^4b_1 \\ - 6x^3b_2 + 6x^2ya_2 - 12x^2yb_3 + 6xy^2a_3 - 12xy^2b_1 + 16y^3a_1 - x^2a_3 - 6x^2b_1 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} 4b_2v_2^6 + 6a_2v_2^5 - 30b_2v_1v_2^4 - 12b_3v_2^5 - 4a_2v_1v_2^3 + 7a_3v_2^4 - 6b_1v_2^4 + 24b_2v_1^2v_2^2 \\ + 8b_3v_1v_2^3 + 16a_1v_2^3 + 6a_2v_1^2v_2 + 6a_3v_1v_2^2 - 12b_1v_1v_2^2 - 6b_2v_1^3 - 12b_3v_1^2v_2 - a_3v_1^2 \\ - 6b_1v_1^2 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned} & -6b_2v_1^3 + 24b_2v_1^2v_2^2 + (6a_2 - 12b_3)v_1^2v_2 + (-a_3 - 6b_1)v_1^2 \\ & - 30b_2v_1v_2^4 + (-4a_2 + 8b_3)v_1v_2^3 + (6a_3 - 12b_1)v_1v_2^2 \\ & + 4b_2v_2^6 + (6a_2 - 12b_3)v_2^5 + (7a_3 - 6b_1)v_2^4 + 16a_1v_2^3 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} 16a_1 &= 0 \\ -30b_2 &= 0 \\ -6b_2 &= 0 \\ 4b_2 &= 0 \\ 24b_2 &= 0 \\ -4a_2 + 8b_3 &= 0 \\ 6a_2 - 12b_3 &= 0 \\ -a_3 - 6b_1 &= 0 \\ 6a_3 - 12b_1 &= 0 \\ 7a_3 - 6b_1 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= 2b_3 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= 2x \\ \eta &= y \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned}\eta &= \eta - \omega(x, y) \xi \\ &= y - \left(-\frac{3y^2 - x}{2y(y^2 - 3x)} \right) (2x) \\ &= \frac{-y^4 + x^2}{-y^3 + 3xy} \\ \xi &= 0\end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y} \right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned}S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{-y^4 + x^2}{-y^3 + 3xy}} dy\end{aligned}$$

Which results in

$$S = \ln(y^2 + x) - \frac{\ln(y^2 - x)}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{3y^2 - x}{2y(y^2 - 3x)}$$

Evaluating all the partial derivatives gives

$$\begin{aligned}R_x &= 1 \\R_y &= 0 \\S_x &= \frac{-3y^2 + x}{-2y^4 + 2x^2} \\S_y &= \frac{-y^3 + 3xy}{-y^4 + x^2}\end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\ln(x + y^2) - \frac{\ln(-x + y^2)}{2} = c_1$$

Which simplifies to

$$\ln(x + y^2) - \frac{\ln(-x + y^2)}{2} = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{3y^2 - x}{2y(y^2 - 3x)}$ 	$R = x$ $S = \ln(y^2 + x) - \frac{\ln(y^2 - 3x)}{2}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$\ln(x + y^2) - \frac{\ln(-x + y^2)}{2} = c_1 \tag{1}$$




Figure 73: Slope field plot

Verification of solutions

$$\ln(x + y^2) - \frac{\ln(-x + y^2)}{2} = c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous G
<- homogeneous successful`

```


✓ Solution by Maple

Time used: 0.156 (sec). Leaf size: 101

```
dsolve((3*y(x)^2-x)+(2*y(x))*(y(x)^2-3*x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{\sqrt{2c_1 - 2\sqrt{c_1(c_1 - 8x)} - 4x}}{2}$$

$$y(x) = \frac{\sqrt{2c_1 - 2\sqrt{c_1(c_1 - 8x)} - 4x}}{2}$$

$$y(x) = -\frac{\sqrt{2c_1 + 2\sqrt{c_1(c_1 - 8x)} - 4x}}{2}$$

$$y(x) = \frac{\sqrt{2c_1 + 2\sqrt{c_1(c_1 - 8x)} - 4x}}{2}$$

✓ Solution by Mathematica

Time used: 15.503 (sec). Leaf size: 185

```
DSolve[(3*y[x]^2-x)+(2*y[x]*(y[x]^2-3*x)*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\sqrt{-2x - e^{\frac{c_1}{2}}\sqrt{8x + e^{c_1}} - e^{c_1}}}{\sqrt{2}}$$

$$y(x) \rightarrow \frac{\sqrt{-2x - e^{\frac{c_1}{2}}\sqrt{8x + e^{c_1}} - e^{c_1}}}{\sqrt{2}}$$

$$y(x) \rightarrow -\frac{\sqrt{-2x + e^{\frac{c_1}{2}}\sqrt{8x + e^{c_1}} - e^{c_1}}}{\sqrt{2}}$$

$$y(x) \rightarrow \frac{\sqrt{-2x + e^{\frac{c_1}{2}}\sqrt{8x + e^{c_1}} - e^{c_1}}}{\sqrt{2}}$$

1.42 problem Problem 56

1.42.1 Solving as homogeneousTypeD2 ode	385
1.42.2 Solving as first order ode lie symmetry lookup ode	387
1.42.3 Solving as bernoulli ode	391
1.42.4 Solving as exact ode	395
1.42.5 Solving as riccati ode	400

Internal problem ID [12153]

Internal file name [OUTPUT/10805_Thursday_September_21_2023_05_46_10_AM_44523640/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 56.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "riccati", "bernoulli", "exactByInspection", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _Bernoulli]
```

$$y(-y + x) - x^2y' = 0$$

1.42.1 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$u(x)x(-u(x)x + x) - x^2(u'(x)x + u(x)) = 0$$

In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u^2}{x}\end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = u^2$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u^2} du &= -\frac{1}{x} dx \\ \int \frac{1}{u^2} du &= \int -\frac{1}{x} dx \\ -\frac{1}{u} &= -\ln(x) + c_2\end{aligned}$$

The solution is

$$-\frac{1}{u(x)} + \ln(x) - c_2 = 0$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$\begin{aligned}-\frac{x}{y} + \ln(x) - c_2 &= 0 \\ -\frac{x}{y} + \ln(x) - c_2 &= 0\end{aligned}$$

Summary

The solution(s) found are the following

$$-\frac{x}{y} + \ln(x) - c_2 = 0 \tag{1}$$




Figure 74: Slope field plot

Verification of solutions

$$-\frac{x}{y} + \ln(x) - c_2 = 0$$

Verified OK.

1.42.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{y(y-x)}{x^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **Bernoulli**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 46: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{y^2}{x}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{y^2}{x}} dy \end{aligned}$$

Which results in

$$S = -\frac{x}{y}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{y(y-x)}{x^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= -\frac{1}{y} \\ S_y &= \frac{x}{y^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -\frac{1}{x} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{1}{R}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\ln(R) + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{x}{y} = -\ln(x) + c_1$$



Which simplifies to

$$-\frac{x}{y} = -\ln(x) + c_1$$

Which gives

$$y = \frac{x}{\ln(x) - c_1}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{y(y-x)}{x^2}$ 	$R = x$ $S = -\frac{x}{y}$	$\frac{dS}{dR} = -\frac{1}{R}$ 

Summary

The solution(s) found are the following

$$y = \frac{x}{\ln(x) - c_1} \quad (1)$$




Figure 75: Slope field plot

Verification of solutions

$$y = \frac{x}{\ln(x) - c_1}$$

Verified OK.

1.42.3 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= -\frac{y(y-x)}{x^2} \end{aligned}$$

This is a Bernoulli ODE.

$$y' = \frac{1}{x}y - \frac{1}{x^2}y^2 \quad (1)$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \quad (2)$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \quad (3)$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned} f_0(x) &= \frac{1}{x} \\ f_1(x) &= -\frac{1}{x^2} \\ n &= 2 \end{aligned}$$

Dividing both sides of ODE (1) by $y^n = y^2$ gives

$$y' \frac{1}{y^2} = \frac{1}{xy} - \frac{1}{x^2} \quad (4)$$

Let

$$\begin{aligned} w &= y^{1-n} \\ &= \frac{1}{y} \end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = -\frac{1}{y^2}y' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} -w'(x) &= \frac{w(x)}{x} - \frac{1}{x^2} \\ w' &= -\frac{w}{x} + \frac{1}{x^2} \end{aligned} \quad (7)$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$p(x) = \frac{1}{x}$$
$$q(x) = \frac{1}{x^2}$$

Hence the ode is

$$w'(x) + \frac{w(x)}{x} = \frac{1}{x^2}$$

The integrating factor μ is

$$\mu = e^{\int \frac{1}{x} dx}$$
$$= x$$

The ode becomes

$$\frac{d}{dx}(\mu w) = (\mu) \left(\frac{1}{x^2} \right)$$
$$\frac{d}{dx}(wx) = (x) \left(\frac{1}{x^2} \right)$$
$$d(wx) = \frac{1}{x} dx$$

Integrating gives

$$wx = \int \frac{1}{x} dx$$
$$wx = \ln(x) + c_1$$

Dividing both sides by the integrating factor $\mu = x$ results in

$$w(x) = \frac{\ln(x)}{x} + \frac{c_1}{x}$$

which simplifies to

$$w(x) = \frac{\ln(x) + c_1}{x}$$

Replacing w in the above by $\frac{1}{y}$ using equation (5) gives the final solution.

$$\frac{1}{y} = \frac{\ln(x) + c_1}{x}$$

Or

$$y = \frac{x}{\ln(x) + c_1}$$

Summary

The solution(s) found are the following

$$y = \frac{x}{\ln(x) + c_1} \tag{1}$$




Figure 76: Slope field plot

Verification of solutions

$$y = \frac{x}{\ln(x) + c_1}$$

Verified OK.

1.42.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (-x^2) dy &= (-y(-y + x)) dx \\ (y(-y + x)) dx + (-x^2) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= y(-y + x) \\ N(x, y) &= -x^2 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(y(-y+x)) \\ &= x - 2y\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(-x^2) \\ &= -2x\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. By inspection $\frac{1}{xy^2}$ is an integrating factor. Therefore by multiplying $M = y(-y+x)$ and $N = -x^2$ by this integrating factor the ode becomes exact. The new M, N are

$$\begin{aligned}M &= \frac{-y+x}{xy} \\ N &= -\frac{x}{y^2}\end{aligned}$$

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} \left(-\frac{x}{y^2}\right) dy &= \left(-\frac{-y+x}{xy}\right) dx \\ \left(\frac{-y+x}{xy}\right) dx + \left(-\frac{x}{y^2}\right) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= \frac{-y+x}{xy} \\ N(x, y) &= -\frac{x}{y^2} \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} \left(\frac{-y+x}{xy} \right) \\ &= -\frac{1}{y^2} \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} \left(-\frac{x}{y^2} \right) \\ &= -\frac{1}{y^2} \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{-y+x}{xy} dx \\ \phi &= -\ln(x) + \frac{x}{y} + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = -\frac{x}{y^2} + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = -\frac{x}{y^2}$. Therefore equation (4) becomes

$$-\frac{x}{y^2} = -\frac{x}{y^2} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = -\ln(x) + \frac{x}{y} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\ln(x) + \frac{x}{y}$$

The solution becomes

$$y = \frac{x}{\ln(x) + c_1}$$

Summary

The solution(s) found are the following

$$y = \frac{x}{\ln(x) + c_1} \tag{1}$$




Figure 77: Slope field plot

Verification of solutions

$$y = \frac{x}{\ln(x) + c_1}$$

Verified OK.

1.42.5 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned}y' &= F(x, y) \\ &= -\frac{y(y-x)}{x^2}\end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = -\frac{y^2}{x^2} + \frac{y}{x}$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = 0$, $f_1(x) = \frac{1}{x}$ and $f_2(x) = -\frac{1}{x^2}$. Let

$$\begin{aligned}y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{-\frac{u}{x^2}}\end{aligned}\tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0\tag{2}$$

But

$$\begin{aligned}f_2' &= \frac{2}{x^3} \\ f_1 f_2 &= -\frac{1}{x^3} \\ f_2^2 f_0 &= 0\end{aligned}$$

Substituting the above terms back in equation (2) gives

$$-\frac{u''(x)}{x^2} - \frac{u'(x)}{x^3} = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = c_1 + \ln(x) c_2$$

The above shows that

$$u'(x) = \frac{c_2}{x}$$

Using the above in (1) gives the solution

$$y = \frac{c_2 x}{c_1 + \ln(x) c_2}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = \frac{x}{c_3 + \ln(x)}$$

Summary

The solution(s) found are the following

$$y = \frac{x}{c_3 + \ln(x)} \tag{1}$$




Figure 78: Slope field plot

Verification of solutions

$$y = \frac{x}{c_3 + \ln(x)}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve((x-y(x))*y(x)- x^2*diff(y(x),x)=0,y(x), singsol=all)
```

$$y(x) = \frac{x}{\ln(x) + c_1}$$

✓ Solution by Mathematica

Time used: 0.242 (sec). Leaf size: 19

```
DSolve[(x-y[x])*y[x]- x^2*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x}{\log(x) + c_1}$$
$$y(x) \rightarrow 0$$

1.43 problem Problem 57

1.43.1 Solving as differentialType ode	403
1.43.2 Solving as homogeneousTypeMapleC ode	405
1.43.3 Solving as first order ode lie symmetry calculated ode	408
1.43.4 Solving as exact ode	413

Internal problem ID [12154]

Internal file name [OUTPUT/10806_Thursday_September_21_2023_05_46_11_AM_52519551/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 57.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "differentialType", "homogeneousTypeMapleC", "first_order_ode_lie_symmetry_calculated"

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$y' - \frac{x + y - 3}{1 - x + y} = 0$$

1.43.1 Solving as differentialType ode

Writing the ode as

$$y' = \frac{x + y - 3}{1 - x + y} \tag{1}$$

Which becomes

$$(-y - 1) dy = (-x) dy + (-x - y + 3) dx \tag{2}$$

But the RHS is complete differential because

$$(-x) dy + (-x - y + 3) dx = d\left(-\frac{1}{2}x^2 - xy + 3x\right)$$

Hence (2) becomes

$$(-y - 1) dy = d\left(-\frac{1}{2}x^2 - xy + 3x\right)$$

Integrating both sides gives gives these solutions

$$y = x - 1 + \sqrt{2x^2 - 2c_1 - 8x + 1} + c_1$$

$$y = x - 1 - \sqrt{2x^2 - 2c_1 - 8x + 1} + c_1$$

Summary

The solution(s) found are the following

$$y = x - 1 + \sqrt{2x^2 - 2c_1 - 8x + 1} + c_1 \quad (1)$$

$$y = x - 1 - \sqrt{2x^2 - 2c_1 - 8x + 1} + c_1 \quad (2)$$




Figure 79: Slope field plot

Verification of solutions

$$y = x - 1 + \sqrt{2x^2 - 2c_1 - 8x + 1} + c_1$$

Verified OK.

$$y = x - 1 - \sqrt{2x^2 - 2c_1 - 8x + 1} + c_1$$

Verified OK.

1.43.2 Solving as homogeneous Type MapleC ode

Let $Y = y + y_0$ and $X = x + x_0$ then the above is transformed to new ode in $Y(X)$

$$\frac{d}{dX}Y(X) = \frac{X + x_0 + Y(X) + y_0 - 3}{1 - X - x_0 + Y(X) + y_0}$$

Solving for possible values of x_0 and y_0 which makes the above ode a homogeneous ode results in

$$x_0 = 2$$

$$y_0 = 1$$

Using these values now it is possible to easily solve for $Y(X)$. The above ode now becomes

$$\frac{d}{dX}Y(X) = \frac{X + Y(X)}{-X + Y(X)}$$

In canonical form, the ODE is

$$\begin{aligned} Y' &= F(X, Y) \\ &= \frac{X + Y}{-X + Y} \end{aligned} \tag{1}$$

An ode of the form $Y' = \frac{M(X, Y)}{N(X, Y)}$ is called homogeneous if the functions $M(X, Y)$ and $N(X, Y)$ are both homogeneous functions and of the same order. Recall that a function $f(X, Y)$ is homogeneous of order n if

$$f(t^n X, t^n Y) = t^n f(X, Y)$$

In this case, it can be seen that both $M = -X - Y$ and $N = X - Y$ are both homogeneous and of the same order $n = 1$. Therefore this is a homogeneous ode. Since this ode is homogeneous, it is converted to separable ODE using the substitution $u = \frac{Y}{X}$, or $Y = uX$. Hence

$$\frac{dY}{dX} = \frac{du}{dX}X + u$$

Applying the transformation $Y = uX$ to the above ODE in (1) gives

$$\begin{aligned} \frac{du}{dX}X + u &= \frac{u + 1}{u - 1} \\ \frac{du}{dX} &= \frac{\frac{u(X)+1}{u(X)-1} - u(X)}{X} \end{aligned}$$

Or

$$\frac{d}{dX}u(X) - \frac{\frac{u(X)+1}{u(X)-1} - u(X)}{X} = 0$$

Or

$$\left(\frac{d}{dX}u(X)\right)Xu(X) - \left(\frac{d}{dX}u(X)\right)X + u(X)^2 - 2u(X) - 1 = 0$$

Or

$$X(u(X) - 1)\left(\frac{d}{dX}u(X)\right) + u(X)^2 - 2u(X) - 1 = 0$$

Which is now solved as separable in $u(X)$. Which is now solved in $u(X)$. In canonical form the ODE is

$$\begin{aligned}u' &= F(X, u) \\ &= f(X)g(u) \\ &= -\frac{u^2 - 2u - 1}{X(u - 1)}\end{aligned}$$

Where $f(X) = -\frac{1}{X}$ and $g(u) = \frac{u^2 - 2u - 1}{u - 1}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{u^2 - 2u - 1}{u - 1}} du &= -\frac{1}{X} dX \\ \int \frac{1}{\frac{u^2 - 2u - 1}{u - 1}} du &= \int -\frac{1}{X} dX \\ \frac{\ln(u^2 - 2u - 1)}{2} &= -\ln(X) + c_2\end{aligned}$$

Raising both side to exponential gives

$$\sqrt{u^2 - 2u - 1} = e^{-\ln(X) + c_2}$$

Which simplifies to

$$\sqrt{u^2 - 2u - 1} = \frac{c_3}{X}$$

Which simplifies to

$$\sqrt{u(X)^2 - 2u(X) - 1} = \frac{c_3 e^{c_2}}{X}$$

The solution is

$$\sqrt{u(X)^2 - 2u(X) - 1} = \frac{c_3 e^{c_2}}{X}$$

Now u in the above solution is replaced back by Y using $u = \frac{Y}{X}$ which results in the solution

$$\sqrt{\frac{Y(X)^2}{X^2} - \frac{2Y(X)}{X} - 1} = \frac{c_3 e^{c_2}}{X}$$

Using the solution for $Y(X)$

$$\sqrt{\frac{Y(X)^2 - 2Y(X)X - X^2}{X^2}} = \frac{c_3 e^{c_2}}{X}$$

And replacing back terms in the above solution using

$$Y = y + y_0$$

$$X = x + x_0$$

Or

$$Y = y + 1$$

$$X = x + 2$$

Then the solution in y becomes

$$\sqrt{\frac{(y-1)^2 - 2(y-1)(x-2) - (x-2)^2}{(x-2)^2}} = \frac{c_3 e^{c_2}}{x-2}$$

Summary

The solution(s) found are the following

$$\sqrt{\frac{(y-1)^2 - 2(y-1)(x-2) - (x-2)^2}{(x-2)^2}} = \frac{c_3 e^{c_2}}{x-2} \quad (1)$$




Figure 80: Slope field plot

Verification of solutions

$$\sqrt{\frac{(y-1)^2 - 2(y-1)(x-2) - (x-2)^2}{(x-2)^2}} = \frac{c_3 e^{c_2}}{x-2}$$

Verified OK.

1.43.3 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{x+y-3}{1-x+y}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (\text{1E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 + \frac{(x+y-3)(b_3-a_2)}{1-x+y} - \frac{(x+y-3)^2 a_3}{(1-x+y)^2} \\ - \left(\frac{1}{1-x+y} + \frac{x+y-3}{(1-x+y)^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(\frac{1}{1-x+y} - \frac{x+y-3}{(1-x+y)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{x^2 a_2 - x^2 a_3 + 3x^2 b_2 - x^2 b_3 - 2xy a_2 - 2xy a_3 - 2xy b_2 + 2xy b_3 - y^2 a_2 - 3y^2 a_3 + y^2 b_2 + y^2 b_3 - 2xa_2 + 6xa_3 - 2xb_2 + 4xb_3 - 2ya_1 + 2ya_2 + 8ya_3 + 2yb_2 - 6yb_3 + 2a_1 + 3a_2 - 9a_3 - 4b_1 + b_2 - 3b_3}{(-1+x-y)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} x^2 a_2 - x^2 a_3 + 3x^2 b_2 - x^2 b_3 - 2xy a_2 - 2xy a_3 - 2xy b_2 + 2xy b_3 - y^2 a_2 \\ - 3y^2 a_3 + y^2 b_2 + y^2 b_3 - 2xa_2 + 6xa_3 + 2xb_2 - 6xb_3 + 4xb_3 - 2ya_1 \\ + 2ya_2 + 8ya_3 + 2yb_2 - 6yb_3 + 2a_1 + 3a_2 - 9a_3 - 4b_1 + b_2 - 3b_3 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} a_2 v_1^2 - 2a_2 v_1 v_2 - a_2 v_2^2 - a_3 v_1^2 - 2a_3 v_1 v_2 - 3a_3 v_2^2 + 3b_2 v_1^2 - 2b_2 v_1 v_2 + b_2 v_2^2 \\ - b_3 v_1^2 + 2b_3 v_1 v_2 + b_3 v_2^2 - 2a_1 v_2 - 2a_2 v_1 + 2a_2 v_2 + 6a_3 v_1 + 8a_3 v_2 + 2b_1 v_1 \\ - 6b_2 v_1 + 2b_2 v_2 + 4b_3 v_1 - 6b_3 v_2 + 2a_1 + 3a_2 - 9a_3 - 4b_1 + b_2 - 3b_3 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned} & (a_2 - a_3 + 3b_2 - b_3) v_1^2 + (-2a_2 - 2a_3 - 2b_2 + 2b_3) v_1 v_2 \\ & + (-2a_2 + 6a_3 + 2b_1 - 6b_2 + 4b_3) v_1 + (-a_2 - 3a_3 + b_2 + b_3) v_2^2 \\ & + (-2a_1 + 2a_2 + 8a_3 + 2b_2 - 6b_3) v_2 + 2a_1 + 3a_2 - 9a_3 - 4b_1 + b_2 - 3b_3 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} -2a_2 - 2a_3 - 2b_2 + 2b_3 &= 0 \\ -a_2 - 3a_3 + b_2 + b_3 &= 0 \\ a_2 - a_3 + 3b_2 - b_3 &= 0 \\ -2a_1 + 2a_2 + 8a_3 + 2b_2 - 6b_3 &= 0 \\ -2a_2 + 6a_3 + 2b_1 - 6b_2 + 4b_3 &= 0 \\ 2a_1 + 3a_2 - 9a_3 - 4b_1 + b_2 - 3b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 3b_2 - 2b_3 \\ a_2 &= -2b_2 + b_3 \\ a_3 &= b_2 \\ b_1 &= -2b_2 - b_3 \\ b_2 &= b_2 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= x - 2 \\ \eta &= y - 1 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= y - 1 - \left(\frac{x + y - 3}{1 - x + y} \right) (x - 2) \\ &= \frac{x^2 + 2xy - y^2 - 6x - 2y + 7}{-1 + x - y} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{x^2 + 2xy - y^2 - 6x - 2y + 7}{-1 + x - y}} dy \end{aligned}$$

Which results in

$$S = \frac{\ln(-x^2 - 2xy + y^2 + 6x + 2y - 7)}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{x + y - 3}{1 - x + y}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{x + y - 3}{x^2 + (2y - 6)x - y^2 - 2y + 7} \\ S_y &= \frac{-1 + x - y}{x^2 + (2y - 6)x - y^2 - 2y + 7} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{\ln(-x^2 + (-2y + 6)x + y^2 + 2y - 7)}{2} = c_1$$

Which simplifies to

$$\frac{\ln(-x^2 + (-2y + 6)x + y^2 + 2y - 7)}{2} = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{x+y-3}{1-x+y}$ 	$R = x$ $S = \frac{\ln(-x^2 + (-2y + 6)x + y^2 + 2y - 7)}{2}$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$\frac{\ln(-x^2 + (-2y + 6)x + y^2 + 2y - 7)}{2} = c_1 \quad (1)$$




Figure 81: Slope field plot

Verification of solutions

$$\frac{\ln(-x^2 + (-2y + 6)x + y^2 + 2y - 7)}{2} = c_1$$

Verified OK.

1.43.4 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (A)$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (1 - x + y) dy &= (x + y - 3) dx \\ (-x - y + 3) dx + (1 - x + y) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -x - y + 3 \\ N(x, y) &= 1 - x + y \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-x - y + 3) \\ &= -1 \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(1 - x + y) \\ &= -1\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \tag{1}$$

$$\frac{\partial \phi}{\partial y} = N \tag{2}$$

Integrating (1) w.r.t. x gives

$$\int \frac{\partial \phi}{\partial x} dx = \int M dx$$

$$\int \frac{\partial \phi}{\partial x} dx = \int -x - y + 3 dx$$

$$\phi = -\frac{x(x + 2y - 6)}{2} + f(y) \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = -x + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = 1 - x + y$. Therefore equation (4) becomes

$$1 - x + y = -x + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = y + 1$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int (y + 1) dy$$

$$f(y) = \frac{1}{2}y^2 + y + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = -\frac{x(x + 2y - 6)}{2} + \frac{y^2}{2} + y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = -\frac{x(x + 2y - 6)}{2} + \frac{y^2}{2} + y$$

Summary

The solution(s) found are the following

$$-\frac{x(x + 2y - 6)}{2} + \frac{y^2}{2} + y = c_1 \quad (1)$$




Figure 82: Slope field plot

Verification of solutions

$$-\frac{x(x + 2y - 6)}{2} + \frac{y^2}{2} + y = c_1$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
trying separable  
trying inverse linear  
trying homogeneous types:  
trying homogeneous C  
trying homogeneous types:  
trying homogeneous D  
<- homogeneous successful  
<- homogeneous successful`
```

✓ Solution by Maple

Time used: 0.578 (sec). Leaf size: 30

```
dsolve(diff(y(x),x)= (x+y(x)-3)/(1-x+y(x)),y(x), singsol=all)
```

$$y(x) = \frac{-\sqrt{2(x-2)^2 c_1^2 + 1} + (-1+x) c_1}{c_1}$$

✓ Solution by Mathematica

Time used: 0.202 (sec). Leaf size: 59

```
DSolve[y'[x]== (x+y[x]-3)/(1-x+y[x]),y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -i\sqrt{-2x^2 + 8x - 1 - c_1} + x - 1$$
$$y(x) \rightarrow i\sqrt{-2x^2 + 8x - 1 - c_1} + x - 1$$

1.44 problem Problem 58

1.44.1 Solving as first order ode lie symmetry lookup ode	418
1.44.2 Solving as bernoulli ode	422
1.44.3 Solving as exact ode	426
1.44.4 Solving as riccati ode	431

Internal problem ID [12155]

Internal file name [OUTPUT/10807_Thursday_September_21_2023_05_46_13_AM_75412810/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 58.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**riccati**", "**bernoulli**", "**exactWithIntegrationFactor**", "**first_order_ode_lie_symmetry_lookup**"

Maple gives the following as the ode type

`[_Bernoulli]`

$$y'x - y^2 \ln(x) + y = 0$$

1.44.1 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{y(y \ln(x) - 1)}{x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is known. It is of type **Bernoulli**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 48: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= xy^2\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{x y^2} dy \end{aligned}$$

Which results in

$$S = -\frac{1}{xy}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{y(y \ln(x) - 1)}{x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{1}{x^2 y} \\ S_y &= \frac{1}{x y^2} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{\ln(x)}{x^2} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{\ln(R)}{R^2}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\frac{\ln(R)}{R} - \frac{1}{R} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$-\frac{1}{yx} = -\frac{\ln(x)}{x} - \frac{1}{x} + c_1$$



Which simplifies to

$$\frac{-c_1xy + y \ln(x) + y - 1}{xy} = 0$$

Which gives

$$y = \frac{1}{-c_1x + \ln(x) + 1}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{y(y \ln(x) - 1)}{x}$ 	$R = x$ $S = -\frac{1}{xy}$	$\frac{dS}{dR} = \frac{\ln(R)}{R^2}$ 

Summary

The solution(s) found are the following

$$y = \frac{1}{-c_1x + \ln(x) + 1} \quad (1)$$




Figure 83: Slope field plot

Verification of solutions

$$y = \frac{1}{-c_1x + \ln(x) + 1}$$

Verified OK.

1.44.2 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{y(y \ln(x) - 1)}{x} \end{aligned}$$

This is a Bernoulli ODE.

$$y' = -\frac{1}{x}y + \frac{\ln(x)}{x}y^2 \quad (1)$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \quad (2)$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \quad (3)$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned} f_0(x) &= -\frac{1}{x} \\ f_1(x) &= \frac{\ln(x)}{x} \\ n &= 2 \end{aligned}$$

Dividing both sides of ODE (1) by $y^n = y^2$ gives

$$y' \frac{1}{y^2} = -\frac{1}{xy} + \frac{\ln(x)}{x} \quad (4)$$

Let

$$\begin{aligned} w &= y^{1-n} \\ &= \frac{1}{y} \end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = -\frac{1}{y^2}y' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned} -w'(x) &= -\frac{w(x)}{x} + \frac{\ln(x)}{x} \\ w' &= \frac{w}{x} - \frac{\ln(x)}{x} \end{aligned} \quad (7)$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$p(x) = -\frac{1}{x}$$
$$q(x) = -\frac{\ln(x)}{x}$$

Hence the ode is

$$w'(x) - \frac{w(x)}{x} = -\frac{\ln(x)}{x}$$

The integrating factor μ is

$$\mu = e^{\int -\frac{1}{x} dx}$$
$$= \frac{1}{x}$$

The ode becomes

$$\frac{d}{dx}(\mu w) = (\mu) \left(-\frac{\ln(x)}{x} \right)$$
$$\frac{d}{dx} \left(\frac{w}{x} \right) = \left(\frac{1}{x} \right) \left(-\frac{\ln(x)}{x} \right)$$
$$d \left(\frac{w}{x} \right) = \left(-\frac{\ln(x)}{x^2} \right) dx$$

Integrating gives

$$\frac{w}{x} = \int -\frac{\ln(x)}{x^2} dx$$
$$\frac{w}{x} = \frac{\ln(x)}{x} + \frac{1}{x} + c_1$$

Dividing both sides by the integrating factor $\mu = \frac{1}{x}$ results in

$$w(x) = x \left(\frac{\ln(x)}{x} + \frac{1}{x} \right) + c_1 x$$

which simplifies to

$$w(x) = c_1 x + \ln(x) + 1$$

Replacing w in the above by $\frac{1}{y}$ using equation (5) gives the final solution.

$$\frac{1}{y} = c_1x + \ln(x) + 1$$

Or

$$y = \frac{1}{c_1x + \ln(x) + 1}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{c_1x + \ln(x) + 1} \tag{1}$$




Figure 84: Slope field plot

Verification of solutions

$$y = \frac{1}{c_1x + \ln(x) + 1}$$

Verified OK.

1.44.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (x) dy &= (-y + y^2 \ln(x)) dx \\ (-y^2 \ln(x) + y) dx + (x) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= -y^2 \ln(x) + y \\ N(x, y) &= x \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(-y^2 \ln(x) + y) \\ &= -2y \ln(x) + 1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x) \\ &= 1\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{x} ((-2y \ln(x) + 1) - (1)) \\ &= -\frac{2y \ln(x)}{x}\end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned}B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= -\frac{1}{y(y \ln(x) - 1)} ((1) - (-2y \ln(x) + 1)) \\ &= -\frac{2 \ln(x)}{y \ln(x) - 1}\end{aligned}$$

Since B depends on x , it can not be used to obtain an integrating factor. We will now try a third method to find an integrating factor. Let

$$R = \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN}$$

R is now checked to see if it is a function of only $t = xy$. Therefore

$$\begin{aligned} R &= \frac{\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y}}{xM - yN} \\ &= \frac{(1) - (-2y \ln(x) + 1)}{x(-y^2 \ln(x) + y) - y(x)} \\ &= -\frac{2}{xy} \end{aligned}$$

Replacing all powers of terms xy by t gives

$$R = -\frac{2}{t}$$

Since R depends on t only, then it can be used to find an integrating factor. Let the integrating factor be μ then

$$\begin{aligned} \mu &= e^{\int R dt} \\ &= e^{\int (-\frac{2}{t}) dt} \end{aligned}$$

The result of integrating gives

$$\begin{aligned} \mu &= e^{-2 \ln(t)} \\ &= \frac{1}{t^2} \end{aligned}$$

Now t is replaced back with xy giving

$$\mu = \frac{1}{x^2 y^2}$$

Multiplying M and N by this integrating factor gives new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N

$$\begin{aligned} \bar{M} &= \mu M \\ &= \frac{1}{x^2 y^2} (-y^2 \ln(x) + y) \\ &= \frac{-y \ln(x) + 1}{x^2 y} \end{aligned}$$

And

$$\begin{aligned} \bar{N} &= \mu N \\ &= \frac{1}{x^2 y^2} (x) \\ &= \frac{1}{x y^2} \end{aligned}$$

A modified ODE is now obtained from the original ODE, which is exact and can be solved. The modified ODE is

$$\begin{aligned} \overline{M} + \overline{N} \frac{dy}{dx} &= 0 \\ \left(\frac{-y \ln(x) + 1}{x^2 y} \right) + \left(\frac{1}{x y^2} \right) \frac{dy}{dx} &= 0 \end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \overline{M} \quad (1)$$

$$\frac{\partial \phi}{\partial y} = \overline{N} \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int \overline{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{-y \ln(x) + 1}{x^2 y} dx \\ \phi &= \frac{y \ln(x) + y - 1}{xy} + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\begin{aligned} \frac{\partial \phi}{\partial y} &= \frac{\ln(x) + 1}{xy} - \frac{y \ln(x) + y - 1}{x y^2} + f'(y) \\ &= \frac{1}{x y^2} + f'(y) \end{aligned} \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{1}{x y^2}$. Therefore equation (4) becomes

$$\frac{1}{x y^2} = \frac{1}{x y^2} + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{y \ln(x) + y - 1}{xy} + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{y \ln(x) + y - 1}{xy}$$

The solution becomes

$$y = \frac{1}{-c_1 x + \ln(x) + 1}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{-c_1 x + \ln(x) + 1} \tag{1}$$




Figure 85: Slope field plot

Verification of solutions

$$y = \frac{1}{-c_1x + \ln(x) + 1}$$

Verified OK.

1.44.4 Solving as riccati ode

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= \frac{y(y \ln(x) - 1)}{x} \end{aligned}$$

This is a Riccati ODE. Comparing the ODE to solve

$$y' = \frac{y^2 \ln(x)}{x} - \frac{y}{x}$$

With Riccati ODE standard form

$$y' = f_0(x) + f_1(x)y + f_2(x)y^2$$

Shows that $f_0(x) = 0$, $f_1(x) = -\frac{1}{x}$ and $f_2(x) = \frac{\ln(x)}{x}$. Let

$$\begin{aligned} y &= \frac{-u'}{f_2 u} \\ &= \frac{-u'}{\frac{\ln(x)u}{x}} \end{aligned} \tag{1}$$

Using the above substitution in the given ODE results (after some simplification) in a second order ODE to solve for $u(x)$ which is

$$f_2 u''(x) - (f_2' + f_1 f_2) u'(x) + f_2^2 f_0 u(x) = 0 \tag{2}$$

But

$$\begin{aligned} f_2' &= -\frac{\ln(x)}{x^2} + \frac{1}{x^2} \\ f_1 f_2 &= -\frac{\ln(x)}{x^2} \\ f_2^2 f_0 &= 0 \end{aligned}$$

Substituting the above terms back in equation (2) gives

$$\frac{\ln(x) u''(x)}{x} - \left(-\frac{2 \ln(x)}{x^2} + \frac{1}{x^2} \right) u'(x) = 0$$

Solving the above ODE (this ode solved using Maple, not this program), gives

$$u(x) = \frac{-\ln(x) c_2 + c_1 x - c_2}{x}$$

The above shows that

$$u'(x) = \frac{\ln(x) c_2}{x^2}$$

Using the above in (1) gives the solution

$$y = -\frac{c_2}{-\ln(x) c_2 + c_1 x - c_2}$$

Dividing both numerator and denominator by c_1 gives, after renaming the constant $\frac{c_2}{c_1} = c_3$ the following solution

$$y = \frac{1}{-c_3 x + \ln(x) + 1}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{-c_3 x + \ln(x) + 1} \tag{1}$$




Figure 86: Slope field plot

Verification of solutions

$$y = \frac{1}{-c_3x + \ln(x) + 1}$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
<- Bernoulli successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 13

```
dsolve(x*diff(y(x),x)-y(x)^2*ln(x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{1}{1 + c_1x + \ln(x)}$$

✓ Solution by Mathematica

Time used: 0.228 (sec). Leaf size: 20

```
DSolve[x*y'[x]-y[x]^2*Log[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{\log(x) + c_1x + 1}$$
$$y(x) \rightarrow 0$$

1.45 problem Problem 59

1.45.1 Solving as linear ode	435
1.45.2 Solving as first order ode lie symmetry lookup ode	437
1.45.3 Solving as exact ode	441
1.45.4 Maple step by step solution	445

Internal problem ID [12156]

Internal file name [OUTPUT/10808_Thursday_September_21_2023_05_46_14_AM_39879519/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 59.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "linear", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

`[_linear]`

$$(x^2 - 1) y' + 2yx = \cos(x)$$

1.45.1 Solving as linear ode

Entering Linear first order ODE solver. In canonical form a linear first order is

$$y' + p(x)y = q(x)$$

Where here

$$p(x) = \frac{2x}{x^2 - 1}$$
$$q(x) = \frac{\cos(x)}{x^2 - 1}$$

Hence the ode is

$$y' + \frac{2xy}{x^2 - 1} = \frac{\cos(x)}{x^2 - 1}$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{-2x}{x^2-1} dx} \\ &= e^{\ln(x-1)+\ln(x+1)}\end{aligned}$$

Which simplifies to

$$\mu = x^2 - 1$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu y) &= (\mu) \left(\frac{\cos(x)}{x^2-1} \right) \\ \frac{d}{dx}((x^2-1)y) &= (x^2-1) \left(\frac{\cos(x)}{x^2-1} \right) \\ d((x^2-1)y) &= \cos(x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}(x^2-1)y &= \int \cos(x) dx \\ (x^2-1)y &= \sin(x) + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = x^2 - 1$ results in

$$y = \frac{\sin(x)}{x^2-1} + \frac{c_1}{x^2-1}$$

which simplifies to

$$y = \frac{\sin(x) + c_1}{x^2-1}$$

Summary

The solution(s) found are the following

$$y = \frac{\sin(x) + c_1}{x^2-1} \tag{1}$$




Figure 87: Slope field plot

Verification of solutions

$$y = \frac{\sin(x) + c_1}{x^2 - 1}$$

Verified OK.

1.45.2 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = \frac{-2xy + \cos(x)}{x^2 - 1}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **linear**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 50: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= e^{-\ln(x-1)-\ln(x+1)}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{e^{-\ln(x-1) - \ln(x+1)}} dy \end{aligned}$$

Which results in

$$S = (x - 1)(x + 1)y$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{-2xy + \cos(x)}{x^2 - 1}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= 2xy \\ S_y &= x^2 - 1 \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \cos(x) \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \cos(R)$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by

integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \sin(R) + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$x^2 y - y = \sin(x) + c_1$$



Which simplifies to

$$x^2 y - y = \sin(x) + c_1$$

Which gives

$$y = \frac{\sin(x) + c_1}{x^2 - 1}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{-2xy + \cos(x)}{x^2 - 1}$ 	$R = x$ $S = x^2 y - y$	$\frac{dS}{dR} = \cos(R)$ 

Summary

The solution(s) found are the following

$$y = \frac{\sin(x) + c_1}{x^2 - 1} \quad (1)$$




Figure 88: Slope field plot

Verification of solutions

$$y = \frac{\sin(x) + c_1}{x^2 - 1}$$

Verified OK.

1.45.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \tag{A}$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \tag{B}$$

Comparing (A,B) shows that

$$\begin{aligned}\frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N\end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \tag{1A}$$

Therefore

$$\begin{aligned}(x^2 - 1) dy &= (-2xy + \cos(x)) dx \\ (2xy - \cos(x)) dx + (x^2 - 1) dy &= 0\end{aligned} \tag{2A}$$

Comparing (1A) and (2A) shows that

$$\begin{aligned}M(x, y) &= 2xy - \cos(x) \\ N(x, y) &= x^2 - 1\end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(2xy - \cos(x)) \\ &= 2x\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(x^2 - 1) \\ &= 2x\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int 2xy - \cos(x) dx \\ \phi &= x^2y - \sin(x) + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = x^2 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = x^2 - 1$. Therefore equation (4) becomes

$$x^2 - 1 = x^2 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = -1$$

Integrating the above w.r.t y gives

$$\begin{aligned} \int f'(y) dy &= \int (-1) dy \\ f(y) &= -y + c_1 \end{aligned}$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = x^2y - \sin(x) - y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = x^2 y - \sin(x) - y$$

The solution becomes

$$y = \frac{\sin(x) + c_1}{x^2 - 1}$$

Summary

The solution(s) found are the following

$$y = \frac{\sin(x) + c_1}{x^2 - 1} \tag{1}$$




Figure 89: Slope field plot

Verification of solutions

$$y = \frac{\sin(x) + c_1}{x^2 - 1}$$

Verified OK.

1.45.4 Maple step by step solution

Let's solve

$$(x^2 - 1)y' + 2yx = \cos(x)$$

- Highest derivative means the order of the ODE is 1

$$y'$$

- Isolate the derivative

$$y' = -\frac{2xy}{x^2-1} + \frac{\cos(x)}{x^2-1}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE

$$y' + \frac{2xy}{x^2-1} = \frac{\cos(x)}{x^2-1}$$

- The ODE is linear; multiply by an integrating factor $\mu(x)$

$$\mu(x) \left(y' + \frac{2xy}{x^2-1} \right) = \frac{\mu(x)\cos(x)}{x^2-1}$$

- Assume the lhs of the ODE is the total derivative $\frac{d}{dx}(\mu(x)y)$

$$\mu(x) \left(y' + \frac{2xy}{x^2-1} \right) = \mu'(x)y + \mu(x)y'$$

- Isolate $\mu'(x)$

$$\mu'(x) = \frac{2\mu(x)x}{x^2-1}$$

- Solve to find the integrating factor

$$\mu(x) = (x-1)(x+1)$$

- Integrate both sides with respect to x

$$\int \left(\frac{d}{dx}(\mu(x)y) \right) dx = \int \frac{\mu(x)\cos(x)}{x^2-1} dx + c_1$$

- Evaluate the integral on the lhs

$$\mu(x)y = \int \frac{\mu(x)\cos(x)}{x^2-1} dx + c_1$$

- Solve for y

$$y = \frac{\int \frac{\mu(x)\cos(x)}{x^2-1} dx + c_1}{\mu(x)}$$

- Substitute $\mu(x) = (x-1)(x+1)$

$$y = \frac{\int \frac{(x-1)(x+1)\cos(x)}{x^2-1} dx + c_1}{(x-1)(x+1)}$$

- Evaluate the integrals on the rhs

$$y = \frac{\sin(x) + c_1}{(x-1)(x+1)}$$

- Simplify

$$y = \frac{\sin(x) + c_1}{x^2 - 1}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
<- 1st order linear successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve((x^2-1)*diff(y(x),x)+2*x*y(x)-cos(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\sin(x) + c_1}{x^2 - 1}$$

✓ Solution by Mathematica

Time used: 0.063 (sec). Leaf size: 18

```
DSolve[(x^2-1)*y'[x]+2*x*y[x]-Cos[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sin(x) + c_1}{x^2 - 1}$$

1.46 problem Problem 60

1.46.1 Solving as first order ode lie symmetry calculated ode 447

Internal problem ID [12157]

Internal file name [OUTPUT/10809_Thursday_September_21_2023_05_46_14_AM_58898071/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 60.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_homogeneous, `class C`], _rational, [_Abel, `2nd type`, `class A`]]
```

$$(4y + 2x + 3)y' - 2y = x + 1$$

1.46.1 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = \frac{2y + x + 1}{4y + 2x + 3}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \quad (\text{A})$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \quad (\text{1E})$$

$$\eta = xb_2 + yb_3 + b_1 \quad (\text{2E})$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 + \frac{(2y+x+1)(b_3-a_2)}{4y+2x+3} - \frac{(2y+x+1)^2 a_3}{(4y+2x+3)^2} \\ - \left(\frac{1}{4y+2x+3} - \frac{2(2y+x+1)}{(4y+2x+3)^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(\frac{2}{4y+2x+3} - \frac{4(2y+x+1)}{(4y+2x+3)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{2x^2a_2 + x^2a_3 - 4x^2b_2 - 2x^2b_3 + 8xya_2 + 4xya_3 - 16xyb_2 - 8xyb_3 + 8y^2a_2 + 4y^2a_3 - 16y^2b_2 - 8y^2b_3 + \dots}{(4y+2x+3)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} -2x^2a_2 - x^2a_3 + 4x^2b_2 + 2x^2b_3 - 8xya_2 - 4xya_3 + 16xyb_2 + 8xyb_3 \\ - 8y^2a_2 - 4y^2a_3 + 16y^2b_2 + 8y^2b_3 - 6xa_2 - 2xa_3 + 10xb_2 + 5xb_3 \\ - 10ya_2 - 5ya_3 + 24yb_2 + 8yb_3 - a_1 - 3a_2 - a_3 - 2b_1 + 9b_2 + 3b_3 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} -2a_2v_1^2 - 8a_2v_1v_2 - 8a_2v_2^2 - a_3v_1^2 - 4a_3v_1v_2 - 4a_3v_2^2 + 4b_2v_1^2 + 16b_2v_1v_2 \\ + 16b_2v_2^2 + 2b_3v_1^2 + 8b_3v_1v_2 + 8b_3v_2^2 - 6a_2v_1 - 10a_2v_2 - 2a_3v_1 - 5a_3v_2 \\ + 10b_2v_1 + 24b_2v_2 + 5b_3v_1 + 8b_3v_2 - a_1 - 3a_2 - a_3 - 2b_1 + 9b_2 + 3b_3 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned} &(-2a_2 - a_3 + 4b_2 + 2b_3)v_1^2 + (-8a_2 - 4a_3 + 16b_2 + 8b_3)v_1v_2 \\ &+ (-6a_2 - 2a_3 + 10b_2 + 5b_3)v_1 + (-8a_2 - 4a_3 + 16b_2 + 8b_3)v_2^2 \\ &+ (-10a_2 - 5a_3 + 24b_2 + 8b_3)v_2 - a_1 - 3a_2 - a_3 - 2b_1 + 9b_2 + 3b_3 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} -10a_2 - 5a_3 + 24b_2 + 8b_3 &= 0 \\ -8a_2 - 4a_3 + 16b_2 + 8b_3 &= 0 \\ -6a_2 - 2a_3 + 10b_2 + 5b_3 &= 0 \\ -2a_2 - a_3 + 4b_2 + 2b_3 &= 0 \\ -a_1 - 3a_2 - a_3 - 2b_1 + 9b_2 + 3b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 5b_2 - 2b_1 \\ a_2 &= 2b_2 \\ a_3 &= 4b_2 \\ b_1 &= b_1 \\ b_2 &= b_2 \\ b_3 &= 2b_2 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= -2 \\ \eta &= 1 \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= 1 - \left(\frac{2y + x + 1}{4y + 2x + 3} \right) (-2) \\ &= \frac{4x + 8y + 5}{4y + 2x + 3} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{4x+8y+5}{4y+2x+3}} dy \end{aligned}$$

Which results in

$$S = \frac{y}{2} + \frac{\ln(4x + 8y + 5)}{16}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = \frac{2y + x + 1}{4y + 2x + 3}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{1}{16x + 32y + 20} \\ S_y &= \frac{4y + 2x + 3}{4x + 8y + 5} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{1}{4} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{1}{4}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{R}{4} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{y}{2} + \frac{\ln(4x + 8y + 5)}{16} = \frac{x}{4} + c_1$$



Which simplifies to

$$\frac{y}{2} + \frac{\ln(4x + 8y + 5)}{16} = \frac{x}{4} + c_1$$

Which gives

$$y = \frac{\text{LambertW}(e^{8x+5+16c_1})}{8} - \frac{x}{2} - \frac{5}{8}$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = \frac{2y+x+1}{4y+2x+3}$ 	$R = x$ $S = \frac{y}{2} + \frac{\ln(4x + 8y + 5)}{16}$	$\frac{dS}{dR} = \frac{1}{4}$ 

Summary

The solution(s) found are the following

$$y = \frac{\text{LambertW}(e^{8x+5+16c_1})}{8} - \frac{x}{2} - \frac{5}{8} \quad (1)$$




Figure 90: Slope field plot

Verification of solutions

$$y = \frac{\text{LambertW}(e^{8x+5+16c_1})}{8} - \frac{x}{2} - \frac{5}{8}$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous C
1st order, trying the canonical coordinates of the invariance group
<- 1st order, canonical coordinates successful
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 20

```
dsolve((4*y(x)+2*x+3)*diff(y(x),x)-2*y(x)-x-1=0,y(x), singsol=all)
```

$$y(x) = -\frac{x}{2} + \frac{\text{LambertW}(c_1 e^{5+8x})}{8} - \frac{5}{8}$$

✓ Solution by Mathematica

Time used: 6.325 (sec). Leaf size: 39

```
DSolve[(4*y[x]+2*x+3)*y'[x]-2*y[x]-x-1==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{8}(W(-e^{8x-1+c_1}) - 4x - 5)$$
$$y(x) \rightarrow \frac{1}{8}(-4x - 5)$$

1.47 problem Problem 61

1.47.1 Solving as differentialType ode	455
1.47.2 Solving as exact ode	459
1.47.3 Maple step by step solution	463

Internal problem ID [12158]

Internal file name [OUTPUT/10810_Thursday_September_21_2023_05_46_16_AM_9194465/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 61.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "differentialType"

Maple gives the following as the ode type

```
[_exact, _rational]
```

$$(-x + y^2) y' - y = -x^2$$

1.47.1 Solving as differentialType ode

Writing the ode as

$$y' = \frac{-x^2 + y}{-x + y^2} \tag{1}$$

Which becomes

$$(-y^2) dy = (-x) dy + (x^2 - y) dx \tag{2}$$

But the RHS is complete differential because

$$(-x) dy + (x^2 - y) dx = d\left(\frac{1}{3}x^3 - xy\right)$$

Hence (2) becomes

$$(-y^2) dy = d\left(\frac{1}{3}x^3 - xy\right)$$

Integrating both sides gives gives these solutions

$$y = \frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2} + \frac{2x}{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}} + c_1$$

$$y = -\frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{4} - \frac{x}{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}} +$$

$$y = -\frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{4} - \frac{x}{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}} -$$

Summary

The solution(s) found are the following

$$y = \frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2} + \frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2x} + c_1 \quad (1)$$

$$y = -\frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{\frac{4}{x}} - \frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2} + \frac{i\sqrt{3} \left(\frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2} - \frac{2x}{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}} \right)}{2} + c_1 \quad (2)$$

$$y = -\frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{\frac{4}{x}} - \frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2} + \frac{i\sqrt{3} \left(\frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2} - \frac{2x}{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}} \right)}{2} + c_1 \quad (3)$$




Figure 91: Slope field plot

Verification of solutions

$$y = \frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2} + \frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2x} + c_1$$

Verified OK.

$$y = -\frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{\frac{4}{x}} - \frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2} + \frac{i\sqrt{3} \left(\frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2} - \frac{2x}{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}} \right)}{2} + c_1$$

Verified OK.

$$y = -\frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{\frac{4}{x}} - \frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2} - \frac{i\sqrt{3} \left(\frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2} - \frac{2x}{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}} \right)}{2} + c_1$$

Verified OK.

1.47.2 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the

ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (y^2 - x) dy &= (-x^2 + y) dx \\ (x^2 - y) dx + (y^2 - x) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= x^2 - y \\ N(x, y) &= y^2 - x \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(x^2 - y) \\ &= -1\end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(y^2 - x) \\ &= -1\end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int x^2 - y dx \\ \phi &= \frac{1}{3}x^3 - xy + f(y)\end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = -x + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = y^2 - x$. Therefore equation (4) becomes

$$y^2 - x = -x + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = y^2$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int (y^2) dy$$

$$f(y) = \frac{y^3}{3} + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{1}{3}x^3 - xy + \frac{1}{3}y^3 + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{1}{3}x^3 - xy + \frac{1}{3}y^3$$

Summary

The solution(s) found are the following

$$\frac{x^3}{3} - yx + \frac{y^3}{3} = c_1 \tag{1}$$




Figure 92: Slope field plot

Verification of solutions

$$\frac{x^3}{3} - yx + \frac{y^3}{3} = c_1$$

Verified OK.

1.47.3 Maple step by step solution

Let's solve

$$(-x + y^2)y' - y = -x^2$$

- Highest derivative means the order of the ODE is 1
 y'
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function
 $F'(x, y) = 0$
 - Compute derivative of lhs
 $F'(x, y) + \left(\frac{\partial}{\partial y} F(x, y)\right) y' = 0$
 - Evaluate derivatives
 $-1 = -1$
 - Condition met, ODE is exact
- Exact ODE implies solution will be of this form
 $\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$
- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x
 $F(x, y) = \int (x^2 - y) dx + f_1(y)$
- Evaluate integral
 $F(x, y) = \frac{x^3}{3} - xy + f_1(y)$
- Take derivative of $F(x, y)$ with respect to y
 $N(x, y) = \frac{\partial}{\partial y} F(x, y)$
- Compute derivative
 $y^2 - x = -x + \frac{d}{dy} f_1(y)$
- Isolate for $\frac{d}{dy} f_1(y)$

$$\frac{d}{dy} f_1(y) = y^2$$

- Solve for $f_1(y)$

$$f_1(y) = \frac{y^3}{3}$$

- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = \frac{1}{3}x^3 - xy + \frac{1}{3}y^3$$

- Substitute $F(x, y)$ into the solution of the ODE

$$\frac{1}{3}x^3 - xy + \frac{1}{3}y^3 = c_1$$

- Solve for y

$$\left\{ y = \frac{\left(-4x^3 + 12c_1 + 4\sqrt{x^6 - 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{2} + \frac{2x}{\left(-4x^3 + 12c_1 + 4\sqrt{x^6 - 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}, y = -\frac{\left(-4x^3 + 12c_1 + 4\sqrt{x^6 - 6c_1x^3 - 4x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{4} \right.$$

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying Chini
differential order: 1; looking for linear symmetries
trying exact
<- exact successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 318

```
dsolve((y(x)^2-x)*diff(y(x),x)-y(x)+x^2=0,y(x), singsol=all)
```

$$y(x) = \frac{\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + (6c_1 - 4)x^3 + 9c_1^2}\right)^{\frac{2}{3}} + 4x}{2\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + (6c_1 - 4)x^3 + 9c_1^2}\right)^{\frac{1}{3}}}$$

$$y(x) = \frac{i\left(-\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + (6c_1 - 4)x^3 + 9c_1^2}\right)^{\frac{2}{3}} + 4x\right)\sqrt{3} - \left(-4x^3 - 12c_1 + 4\sqrt{x^6 + (6c_1 - 4)x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{4\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + (6c_1 - 4)x^3 + 9c_1^2}\right)^{\frac{1}{3}}}$$

$$y(x) = \frac{i\left(\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + (6c_1 - 4)x^3 + 9c_1^2}\right)^{\frac{2}{3}} - 4x\right)\sqrt{3} - \left(-4x^3 - 12c_1 + 4\sqrt{x^6 + (6c_1 - 4)x^3 + 9c_1^2}\right)^{\frac{1}{3}}}{4\left(-4x^3 - 12c_1 + 4\sqrt{x^6 + (6c_1 - 4)x^3 + 9c_1^2}\right)^{\frac{1}{3}}}$$

✓ Solution by Mathematica

Time used: 4.856 (sec). Leaf size: 326

```
DSolve[(y[x]^2-x)*y'[x]-y[x]+x^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{2x + \sqrt[3]{2}\left(x^3 + \sqrt{x^6 + (-4 + 6c_1)x^3 + 9c_1^2} + 3c_1\right)^{2/3}}{2^{2/3}\sqrt[3]{x^3 + \sqrt{x^6 + (-4 + 6c_1)x^3 + 9c_1^2} + 3c_1}}$$

$$y(x) \rightarrow \frac{2^{2/3}(1 - i\sqrt{3})\left(x^3 + \sqrt{x^6 + (-4 + 6c_1)x^3 + 9c_1^2} + 3c_1\right)^{2/3} + \sqrt[3]{2}(2 + 2i\sqrt{3})x}{4\sqrt[3]{x^3 + \sqrt{x^6 + (-4 + 6c_1)x^3 + 9c_1^2} + 3c_1}}$$

$$y(x) \rightarrow \frac{2^{2/3}(1 + i\sqrt{3})\left(x^3 + \sqrt{x^6 + (-4 + 6c_1)x^3 + 9c_1^2} + 3c_1\right)^{2/3} + \sqrt[3]{2}(2 - 2i\sqrt{3})x}{4\sqrt[3]{x^3 + \sqrt{x^6 + (-4 + 6c_1)x^3 + 9c_1^2} + 3c_1}}$$

1.48 problem Problem 62

1.48.1 Solving as homogeneousTypeD2 ode	466
1.48.2 Solving as first order ode lie symmetry calculated ode	468
1.48.3 Solving as exact ode	473

Internal problem ID [12159]

Internal file name [OUTPUT/10811_Thursday_September_21_2023_05_46_17_AM_82658170/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 62.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**homogeneousTypeD2**", "**exactWithIntegrationFactor**", "**first_order_ode_lie_symmetry_calculated**"

Maple gives the following as the ode type

```
[[_homogeneous, `class A`], _rational, _dAlembert]
```

$$(y^2 - x^2)y' + 2yx = 0$$

1.48.1 Solving as homogeneousTypeD2 ode

Using the change of variables $y = u(x)x$ on the above ode results in new ode in $u(x)$

$$(u(x)^2 x^2 - x^2)(u'(x)x + u(x)) + 2u(x)x^2 = 0$$

In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{u(u^2 + 1)}{(u^2 - 1)x}\end{aligned}$$

Where $f(x) = -\frac{1}{x}$ and $g(u) = \frac{u(u^2+1)}{u^2-1}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{u(u^2+1)}{u^2-1}} du &= -\frac{1}{x} dx \\ \int \frac{1}{\frac{u(u^2+1)}{u^2-1}} du &= \int -\frac{1}{x} dx \\ -\ln(u) + \ln(u^2 + 1) &= -\ln(x) + c_2\end{aligned}$$

Raising both side to exponential gives

$$e^{-\ln(u)+\ln(u^2+1)} = e^{-\ln(x)+c_2}$$

Which simplifies to

$$\frac{u^2 + 1}{u} = \frac{c_3}{x}$$

The solution is

$$\frac{u(x)^2 + 1}{u(x)} = \frac{c_3}{x}$$

Replacing $u(x)$ in the above solution by $\frac{y}{x}$ results in the solution for y in implicit form

$$\begin{aligned}\frac{x\left(\frac{y^2}{x^2} + 1\right)}{y} &= \frac{c_3}{x} \\ \frac{x^2 + y^2}{yx} &= \frac{c_3}{x}\end{aligned}$$

Which simplifies to

$$\frac{x^2 + y^2}{y} = c_3$$

Summary

The solution(s) found are the following

$$\frac{x^2 + y^2}{y} = c_3 \tag{1}$$




Figure 93: Slope field plot

Verification of solutions

$$\frac{x^2 + y^2}{y} = c_3$$

Verified OK.

1.48.2 Solving as first order ode lie symmetry calculated ode

Writing the ode as

$$y' = -\frac{2yx}{-x^2 + y^2}$$

$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2 \xi_y - \omega_x \xi - \omega_y \eta = 0 \tag{A}$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = xa_2 + ya_3 + a_1 \tag{1E}$$

$$\eta = xb_2 + yb_3 + b_1 \tag{2E}$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 - \frac{2yx(b_3 - a_2)}{-x^2 + y^2} - \frac{4y^2x^2a_3}{(-x^2 + y^2)^2} \\ - \left(-\frac{2y}{-x^2 + y^2} - \frac{4yx^2}{(-x^2 + y^2)^2} \right) (xa_2 + ya_3 + a_1) \\ - \left(-\frac{2x}{-x^2 + y^2} + \frac{4y^2x}{(-x^2 + y^2)^2} \right) (xb_2 + yb_3 + b_1) = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{-x^4b_2 + 2y^2x^2a_3 + 4x^2y^2b_2 - 4xy^3a_2 + 4xy^3b_3 - 2y^4a_3 - y^4b_2 + 2x^3b_1 - 2x^2ya_1 + 2xy^2b_1 - 2y^3a_1}{(x^2 - y^2)^2} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} -x^4b_2 - 2y^2x^2a_3 - 4x^2y^2b_2 + 4xy^3a_2 - 4xy^3b_3 + 2y^4a_3 \\ + y^4b_2 - 2x^3b_1 + 2x^2ya_1 - 2xy^2b_1 + 2y^3a_1 = 0 \end{aligned} \quad (6E)$$

Looking at the above PDE shows the following are all the terms with $\{x, y\}$ in them.

$$\{x, y\}$$

The following substitution is now made to be able to collect on all terms with $\{x, y\}$ in them

$$\{x = v_1, y = v_2\}$$

The above PDE (6E) now becomes

$$\begin{aligned} 4a_2v_1v_2^3 - 2a_3v_1^2v_2^2 + 2a_3v_2^4 - b_2v_1^4 - 4b_2v_1^2v_2^2 + b_2v_2^4 \\ - 4b_3v_1v_2^3 + 2a_1v_1^2v_2 + 2a_1v_2^3 - 2b_1v_1^3 - 2b_1v_1v_2^2 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2\}$$

Equation (7E) now becomes

$$\begin{aligned} -b_2v_1^4 - 2b_1v_1^3 + (-2a_3 - 4b_2)v_1^2v_2^2 + 2a_1v_1^2v_2 \\ + (4a_2 - 4b_3)v_1v_2^3 - 2b_1v_1v_2^2 + (2a_3 + b_2)v_2^4 + 2a_1v_2^3 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} 2a_1 &= 0 \\ -2b_1 &= 0 \\ -b_2 &= 0 \\ 4a_2 - 4b_3 &= 0 \\ -2a_3 - 4b_2 &= 0 \\ 2a_3 + b_2 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 0 \\ a_2 &= b_3 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= 0 \\ b_3 &= b_3 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= x \\ \eta &= y \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(x, y) \xi \\ &= y - \left(-\frac{2yx}{-x^2 + y^2} \right) (x) \\ &= \frac{-x^2y - y^3}{x^2 - y^2} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right) S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{-x^2y - y^3}{x^2 - y^2}} dy \end{aligned}$$

Which results in

$$S = \ln(x^2 + y^2) - \ln(y)$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{2yx}{-x^2 + y^2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{2x}{x^2 + y^2} \\ S_y &= \frac{2y}{x^2 + y^2} - \frac{1}{y} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = 0 \tag{2A}$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = 0$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = c_1 \tag{4}$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\ln(x^2 + y^2) - \ln(y) = c_1$$

Which simplifies to

$$\ln(x^2 + y^2) - \ln(y) = c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{2yx}{-x^2+y^2}$ 	$R = x$ $S = \ln(x^2 + y^2) - \ln(y)$	$\frac{dS}{dR} = 0$ 

Summary

The solution(s) found are the following

$$\ln(x^2 + y^2) - \ln(y) = c_1 \quad (1)$$




Figure 94: Slope field plot

Verification of solutions

$$\ln(x^2 + y^2) - \ln(y) = c_1$$

Verified OK.

1.48.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (A)$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx} \phi(x, y) = 0$$

Hence

$$\frac{\partial \phi}{\partial x} + \frac{\partial \phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial \phi}{\partial x} &= M \\ \frac{\partial \phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition $\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (\text{1A})$$

Therefore

$$\begin{aligned} (-x^2 + y^2) dy &= (-2xy) dx \\ (2xy) dx + (-x^2 + y^2) dy &= 0 \end{aligned} \quad (\text{2A})$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= 2xy \\ N(x, y) &= -x^2 + y^2 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y}(2xy) \\ &= 2x \end{aligned}$$

And

$$\begin{aligned}\frac{\partial N}{\partial x} &= \frac{\partial}{\partial x}(-x^2 + y^2) \\ &= -2x\end{aligned}$$

Since $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$, then the ODE is not exact. Since the ODE is not exact, we will try to find an integrating factor to make it exact. Let

$$\begin{aligned}A &= \frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) \\ &= \frac{1}{-x^2 + y^2} ((2x) - (-2x)) \\ &= -\frac{4x}{x^2 - y^2}\end{aligned}$$

Since A depends on y , it can not be used to obtain an integrating factor. We will now try a second method to find an integrating factor. Let

$$\begin{aligned}B &= \frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \\ &= \frac{1}{2xy} ((-2x) - (2x)) \\ &= -\frac{2}{y}\end{aligned}$$

Since B does not depend on x , it can be used to obtain an integrating factor. Let the integrating factor be μ . Then

$$\begin{aligned}\mu &= e^{\int B \, dy} \\ &= e^{\int -\frac{2}{y} \, dy}\end{aligned}$$

The result of integrating gives

$$\begin{aligned}\mu &= e^{-2 \ln(y)} \\ &= \frac{1}{y^2}\end{aligned}$$

M and N are now multiplied by this integrating factor, giving new M and new N which are called \bar{M} and \bar{N} so not to confuse them with the original M and N .

$$\begin{aligned}\bar{M} &= \mu M \\ &= \frac{1}{y^2} (2xy) \\ &= \frac{2x}{y}\end{aligned}$$

And

$$\begin{aligned}\bar{N} &= \mu N \\ &= \frac{1}{y^2}(-x^2 + y^2) \\ &= \frac{-x^2 + y^2}{y^2}\end{aligned}$$

So now a modified ODE is obtained from the original ODE which will be exact and can be solved using the standard method. The modified ODE is

$$\begin{aligned}\bar{M} + \bar{N} \frac{dy}{dx} &= 0 \\ \left(\frac{2x}{y}\right) + \left(\frac{-x^2 + y^2}{y^2}\right) \frac{dy}{dx} &= 0\end{aligned}$$

The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = \bar{M} \tag{1}$$

$$\frac{\partial \phi}{\partial y} = \bar{N} \tag{2}$$

Integrating (1) w.r.t. x gives

$$\begin{aligned}\int \frac{\partial \phi}{\partial x} dx &= \int \bar{M} dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int \frac{2x}{y} dx \\ \phi &= \frac{x^2}{y} + f(y)\end{aligned} \tag{3}$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = -\frac{x^2}{y^2} + f'(y) \tag{4}$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = \frac{-x^2 + y^2}{y^2}$. Therefore equation (4) becomes

$$\frac{-x^2 + y^2}{y^2} = -\frac{x^2}{y^2} + f'(y) \tag{5}$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 1$$

Integrating the above w.r.t y gives

$$\int f'(y) dy = \int (1) dy$$

$$f(y) = y + c_1$$

Where c_1 is constant of integration. Substituting result found above for $f(y)$ into equation (3) gives ϕ

$$\phi = \frac{x^2}{y} + y + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = \frac{x^2}{y} + y$$

Summary

The solution(s) found are the following

$$\frac{x^2}{y} + y = c_1 \tag{1}$$




Figure 95: Slope field plot

Verification of solutions

$$\frac{x^2}{y} + y = c_1$$

Verified OK.

Maple trace

```

`Methods for first order ODEs:
--- Trying classification methods ---
trying a quadrature
trying 1st order linear
trying Bernoulli
trying separable
trying inverse linear
trying homogeneous types:
trying homogeneous D
<- homogeneous successful`

```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 47

```
dsolve((y(x)^2-x^2)*diff(y(x),x)+2*x*y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{1 - \sqrt{-4c_1^2x^2 + 1}}{2c_1}$$
$$y(x) = \frac{1 + \sqrt{-4c_1^2x^2 + 1}}{2c_1}$$

✓ Solution by Mathematica

Time used: 1.683 (sec). Leaf size: 66

```
DSolve[(y[x]^2-x^2)*y'[x]+2*x*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left(e^{c_1} - \sqrt{-4x^2 + e^{2c_1}} \right)$$
$$y(x) \rightarrow \frac{1}{2} \left(\sqrt{-4x^2 + e^{2c_1}} + e^{c_1} \right)$$
$$y(x) \rightarrow 0$$

1.49 problem Problem 63

1.49.1 Solving as first order ode lie symmetry lookup ode	480
1.49.2 Solving as bernoulli ode	484
1.49.3 Solving as exact ode	488
1.49.4 Maple step by step solution	491

Internal problem ID [12160]

Internal file name [OUTPUT/10812_Thursday_September_21_2023_05_46_19_AM_35121215/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 63.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "exact", "bernoulli", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

```
[[_homogeneous, `class G`], _exact, _rational, _Bernoulli]
```

$$3xy^2y' + y^3 = 2x$$

1.49.1 Solving as first order ode lie symmetry lookup ode

Writing the ode as

$$y' = -\frac{y^3 - 2x}{3y^2x}$$
$$y' = \omega(x, y)$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_x + \omega(\eta_y - \xi_x) - \omega^2\xi_y - \omega_x\xi - \omega_y\eta = 0 \quad (\text{A})$$

The type of this ode is known. It is of type **Bernoulli**. Therefore we do not need to solve the PDE (A), and can just use the lookup table shown below to find ξ, η

Table 54: Lie symmetry infinitesimal lookup table for known first order ODE's

ODE class	Form	ξ	η
linear ode	$y' = f(x)y(x) + g(x)$	0	$e^{\int f dx}$
separable ode	$y' = f(x)g(y)$	$\frac{1}{f}$	0
quadrature ode	$y' = f(x)$	0	1
quadrature ode	$y' = g(y)$	1	0
homogeneous ODEs of Class A	$y' = f\left(\frac{y}{x}\right)$	x	y
homogeneous ODEs of Class C	$y' = (a + bx + cy)^{\frac{n}{m}}$	1	$-\frac{b}{c}$
homogeneous class D	$y' = \frac{y}{x} + g(x)F\left(\frac{y}{x}\right)$	x^2	xy
First order special form ID 1	$y' = g(x)e^{h(x)+by} + f(x)$	$\frac{e^{-\int bf(x)dx-h(x)}}{g(x)}$	$\frac{f(x)e^{-\int bf(x)dx-h(x)}}{g(x)}$
polynomial type ode	$y' = \frac{a_1x+b_1y+c_1}{a_2x+b_2y+c_2}$	$\frac{a_1b_2x-a_2b_1x-b_1c_2+b_2c_1}{a_1b_2-a_2b_1}$	$\frac{a_1b_2y-a_2b_1y-a_1c_2-a_2c_1}{a_1b_2-a_2b_1}$
Bernoulli ode	$y' = f(x)y + g(x)y^n$	0	$e^{-\int (n-1)f(x)dx}y^n$
Reduced Riccati	$y' = f_1(x)y + f_2(x)y^2$	0	$e^{-\int f_1 dx}$

The above table shows that

$$\begin{aligned}\xi(x, y) &= 0 \\ \eta(x, y) &= \frac{1}{xy^2}\end{aligned}\tag{A1}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(x, y) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dx}{\xi} = \frac{dy}{\eta} = dS\tag{1}$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial y}\right)S(x, y) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the

canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = x$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\frac{1}{xy^2}} dy \end{aligned}$$

Which results in

$$S = \frac{xy^3}{3}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_x + \omega(x, y)S_y}{R_x + \omega(x, y)R_y} \quad (2)$$

Where in the above R_x, R_y, S_x, S_y are all partial derivatives and $\omega(x, y)$ is the right hand side of the original ode given by

$$\omega(x, y) = -\frac{y^3 - 2x}{3y^2x}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_x &= 1 \\ R_y &= 0 \\ S_x &= \frac{y^3}{3} \\ S_y &= xy^2 \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = \frac{2x}{3} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for x, y in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = \frac{2R}{3}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = \frac{R^2}{3} + c_1 \quad (4)$$



To complete the solution, we just need to transform (4) back to x, y coordinates. This results in

$$\frac{y^3 x}{3} = \frac{x^2}{3} + c_1$$

Which simplifies to

$$\frac{y^3 x}{3} = \frac{x^2}{3} + c_1$$

The following diagram shows solution curves of the original ode and how they transform in the canonical coordinates space using the mapping shown.

Original ode in x, y coordinates	Canonical coordinates transformation	ODE in canonical coordinates (R, S)
$\frac{dy}{dx} = -\frac{y^3 - 2x}{3y^2 x}$ 	$R = x$ $S = \frac{x y^3}{3}$	$\frac{dS}{dR} = \frac{2R}{3}$ 

Summary

The solution(s) found are the following

$$\frac{y^3 x}{3} = \frac{x^2}{3} + c_1 \quad (1)$$




Figure 96: Slope field plot

Verification of solutions

$$\frac{y^3 x}{3} = \frac{x^2}{3} + c_1$$

Verified OK.

1.49.2 Solving as bernoulli ode

In canonical form, the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= -\frac{y^3 - 2x}{3y^2 x} \end{aligned}$$

This is a Bernoulli ODE.

$$y' = -\frac{1}{3x}y + \frac{2}{3} \frac{1}{y^2} \tag{1}$$

The standard Bernoulli ODE has the form

$$y' = f_0(x)y + f_1(x)y^n \tag{2}$$

The first step is to divide the above equation by y^n which gives

$$\frac{y'}{y^n} = f_0(x)y^{1-n} + f_1(x) \tag{3}$$

The next step is use the substitution $w = y^{1-n}$ in equation (3) which generates a new ODE in $w(x)$ which will be linear and can be easily solved using an integrating factor. Backsubstitution then gives the solution $y(x)$ which is what we want.

This method is now applied to the ODE at hand. Comparing the ODE (1) With (2) Shows that

$$\begin{aligned}f_0(x) &= -\frac{1}{3x} \\f_1(x) &= \frac{2}{3} \\n &= -2\end{aligned}$$

Dividing both sides of ODE (1) by $y^n = \frac{1}{y^2}$ gives

$$y'y^2 = -\frac{y^3}{3x} + \frac{2}{3} \quad (4)$$

Let

$$\begin{aligned}w &= y^{1-n} \\&= y^3\end{aligned} \quad (5)$$

Taking derivative of equation (5) w.r.t x gives

$$w' = 3y^2y' \quad (6)$$

Substituting equations (5) and (6) into equation (4) gives

$$\begin{aligned}\frac{w'(x)}{3} &= -\frac{w(x)}{3x} + \frac{2}{3} \\w' &= -\frac{w}{x} + 2\end{aligned} \quad (7)$$

The above now is a linear ODE in $w(x)$ which is now solved.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$w'(x) + p(x)w(x) = q(x)$$

Where here

$$\begin{aligned}p(x) &= \frac{1}{x} \\q(x) &= 2\end{aligned}$$

Hence the ode is

$$w'(x) + \frac{w(x)}{x} = 2$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{1}{x} dx} \\ &= x\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dx}(\mu w) &= (\mu) (2) \\ \frac{d}{dx}(xw) &= (x) (2) \\ d(xw) &= (2x) dx\end{aligned}$$

Integrating gives

$$\begin{aligned}xw &= \int 2x dx \\ xw &= x^2 + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = x$ results in

$$w(x) = x + \frac{c_1}{x}$$

Replacing w in the above by y^3 using equation (5) gives the final solution.

$$y^3 = x + \frac{c_1}{x}$$

Solving for y gives

$$\begin{aligned}y(x) &= \frac{((x^2 + c_1) x^2)^{\frac{1}{3}}}{x} \\ y(x) &= \frac{((x^2 + c_1) x^2)^{\frac{1}{3}} (i\sqrt{3} - 1)}{2x} \\ y(x) &= -\frac{((x^2 + c_1) x^2)^{\frac{1}{3}} (1 + i\sqrt{3})}{2x}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{((x^2 + c_1) x^2)^{\frac{1}{3}}}{x} \quad (1)$$

$$y = \frac{((x^2 + c_1) x^2)^{\frac{1}{3}} (i\sqrt{3} - 1)}{2x} \quad (2)$$

$$y = -\frac{((x^2 + c_1) x^2)^{\frac{1}{3}} (1 + i\sqrt{3})}{2x} \quad (3)$$




Figure 97: Slope field plot

Verification of solutions

$$y = \frac{((x^2 + c_1)x^2)^{\frac{1}{3}}}{x}$$

Verified OK.

$$y = \frac{((x^2 + c_1)x^2)^{\frac{1}{3}}(i\sqrt{3} - 1)}{2x}$$

Verified OK.

$$y = -\frac{((x^2 + c_1)x^2)^{\frac{1}{3}}(1 + i\sqrt{3})}{2x}$$

Verified OK.

1.49.3 Solving as exact ode

Entering Exact first order ODE solver. (Form one type)

To solve an ode of the form

$$M(x, y) + N(x, y) \frac{dy}{dx} = 0 \quad (\text{A})$$

We assume there exists a function $\phi(x, y) = c$ where c is constant, that satisfies the ode. Taking derivative of ϕ w.r.t. x gives

$$\frac{d}{dx}\phi(x, y) = 0$$

Hence

$$\frac{\partial\phi}{\partial x} + \frac{\partial\phi}{\partial y} \frac{dy}{dx} = 0 \quad (\text{B})$$

Comparing (A,B) shows that

$$\begin{aligned} \frac{\partial\phi}{\partial x} &= M \\ \frac{\partial\phi}{\partial y} &= N \end{aligned}$$

But since $\frac{\partial^2\phi}{\partial x\partial y} = \frac{\partial^2\phi}{\partial y\partial x}$ then for the above to be valid, we require that

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

If the above condition is satisfied, then the original ode is called exact. We still need to determine $\phi(x, y)$ but at least we know now that we can do that since the condition

$\frac{\partial^2 \phi}{\partial x \partial y} = \frac{\partial^2 \phi}{\partial y \partial x}$ is satisfied. If this condition is not satisfied then this method will not work and we have to now look for an integrating factor to force this condition, which might or might not exist. The first step is to write the ODE in standard form to check for exactness, which is

$$M(x, y) dx + N(x, y) dy = 0 \quad (1A)$$

Therefore

$$\begin{aligned} (3x y^2) dy &= (-y^3 + 2x) dx \\ (y^3 - 2x) dx + (3x y^2) dy &= 0 \end{aligned} \quad (2A)$$

Comparing (1A) and (2A) shows that

$$\begin{aligned} M(x, y) &= y^3 - 2x \\ N(x, y) &= 3x y^2 \end{aligned}$$

The next step is to determine if the ODE is exact or not. The ODE is exact when the following condition is satisfied

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

Using result found above gives

$$\begin{aligned} \frac{\partial M}{\partial y} &= \frac{\partial}{\partial y} (y^3 - 2x) \\ &= 3y^2 \end{aligned}$$

And

$$\begin{aligned} \frac{\partial N}{\partial x} &= \frac{\partial}{\partial x} (3x y^2) \\ &= 3y^2 \end{aligned}$$

Since $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$, then the ODE is exact. The following equations are now set up to solve for the function $\phi(x, y)$

$$\frac{\partial \phi}{\partial x} = M \quad (1)$$

$$\frac{\partial \phi}{\partial y} = N \quad (2)$$

Integrating (1) w.r.t. x gives

$$\begin{aligned} \int \frac{\partial \phi}{\partial x} dx &= \int M dx \\ \int \frac{\partial \phi}{\partial x} dx &= \int y^3 - 2x dx \\ \phi &= x y^3 - x^2 + f(y) \end{aligned} \quad (3)$$

Where $f(y)$ is used for the constant of integration since ϕ is a function of both x and y . Taking derivative of equation (3) w.r.t y gives

$$\frac{\partial \phi}{\partial y} = 3x y^2 + f'(y) \quad (4)$$

But equation (2) says that $\frac{\partial \phi}{\partial y} = 3x y^2$. Therefore equation (4) becomes

$$3x y^2 = 3x y^2 + f'(y) \quad (5)$$

Solving equation (5) for $f'(y)$ gives

$$f'(y) = 0$$

Therefore

$$f(y) = c_1$$

Where c_1 is constant of integration. Substituting this result for $f(y)$ into equation (3) gives ϕ

$$\phi = x y^3 - x^2 + c_1$$

But since ϕ itself is a constant function, then let $\phi = c_2$ where c_2 is new constant and combining c_1 and c_2 constants into new constant c_1 gives the solution as

$$c_1 = x y^3 - x^2$$

Summary

The solution(s) found are the following

$$y^3 x - x^2 = c_1 \quad (1)$$




Figure 98: Slope field plot

Verification of solutions

$$y^3x - x^2 = c_1$$

Verified OK.

1.49.4 Maple step by step solution

Let's solve

$$3xy^2y' + y^3 = 2x$$

- Highest derivative means the order of the ODE is 1
 y'
- Check if ODE is exact
 - ODE is exact if the lhs is the total derivative of a C^2 function
 $F'(x, y) = 0$
 - Compute derivative of lhs
 $F'(x, y) + \left(\frac{\partial}{\partial y}F(x, y)\right) y' = 0$

- Evaluate derivatives

$$3y^2 = 3y^2$$

- Condition met, ODE is exact

- Exact ODE implies solution will be of this form

$$\left[F(x, y) = c_1, M(x, y) = F'(x, y), N(x, y) = \frac{\partial}{\partial y} F(x, y) \right]$$

- Solve for $F(x, y)$ by integrating $M(x, y)$ with respect to x

$$F(x, y) = \int (y^3 - 2x) dx + f_1(y)$$

- Evaluate integral

$$F(x, y) = x y^3 - x^2 + f_1(y)$$

- Take derivative of $F(x, y)$ with respect to y

$$N(x, y) = \frac{\partial}{\partial y} F(x, y)$$

- Compute derivative

$$3x y^2 = 3x y^2 + \frac{d}{dy} f_1(y)$$

- Isolate for $\frac{d}{dy} f_1(y)$

$$\frac{d}{dy} f_1(y) = 0$$

- Solve for $f_1(y)$

$$f_1(y) = 0$$

- Substitute $f_1(y)$ into equation for $F(x, y)$

$$F(x, y) = x y^3 - x^2$$

- Substitute $F(x, y)$ into the solution of the ODE

$$x y^3 - x^2 = c_1$$

- Solve for y

$$\left\{ y = \frac{((x^2+c_1)x^2)^{\frac{1}{3}}}{x}, y = -\frac{((x^2+c_1)x^2)^{\frac{1}{3}}}{2x} - \frac{I\sqrt{3}((x^2+c_1)x^2)^{\frac{1}{3}}}{2x}, y = -\frac{((x^2+c_1)x^2)^{\frac{1}{3}}}{2x} + \frac{I\sqrt{3}((x^2+c_1)x^2)^{\frac{1}{3}}}{2x} \right\}$$

Maple trace

```
`Methods for first order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying 1st order linear  
trying Bernoulli  
<- Bernoulli successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 73

```
dsolve(3*x*y(x)^2*diff(y(x),x)+y(x)^3-2*x=0,y(x), singsol=all)
```

$$y(x) = \frac{((x^2 + c_1) x^2)^{\frac{1}{3}}}{x}$$
$$y(x) = -\frac{((x^2 + c_1) x^2)^{\frac{1}{3}} (1 + i\sqrt{3})}{2x}$$
$$y(x) = \frac{((x^2 + c_1) x^2)^{\frac{1}{3}} (i\sqrt{3} - 1)}{2x}$$

✓ Solution by Mathematica

Time used: 0.352 (sec). Leaf size: 72

```
DSolve[3*x*y[x]^2*y'[x]+y[x]^3-2*x==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\sqrt[3]{x^2 + c_1}}{\sqrt[3]{x}}$$
$$y(x) \rightarrow -\frac{\sqrt[3]{-1} \sqrt[3]{x^2 + c_1}}{\sqrt[3]{x}}$$
$$y(x) \rightarrow \frac{(-1)^{2/3} \sqrt[3]{x^2 + c_1}}{\sqrt[3]{x}}$$

1.50 problem Problem 64

1.50.1 Solving as clairaut ode 494

Internal problem ID [12161]

Internal file name [OUTPUT/10813_Thursday_September_21_2023_05_46_20_AM_58836891/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 64.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "**clairaut**"

Maple gives the following as the ode type

```
[[_1st_order , _with_linear_symmetries] , _Clairaut]
```

$$y'^2 + (x + a)y' - y = 0$$

1.50.1 Solving as clairaut ode

This is Clairaut ODE. It has the form

$$y = y'x + g(y')$$

Where g is function of $y'(x)$. Let $p = y'$ the ode becomes

$$p^2 + (x + a)p - y = 0$$

Solving for y from the above results in

$$y = pa + p^2 + px \tag{1A}$$

The above ode is a Clairaut ode which is now solved. We start by replacing y' by p which gives

$$\begin{aligned} y &= pa + p^2 + px \\ &= pa + p^2 + px \end{aligned}$$

Writing the ode as

$$y = px + g(p)$$

We now write $g \equiv g(p)$ to make notation simpler but we should always remember that g is function of p which in turn is function of x . Hence the above becomes

$$y = px + g \tag{1}$$

Then we see that

$$g = pa + p^2$$

Taking derivative of (1) w.r.t. x gives

$$\begin{aligned} p &= \frac{d}{dx}(xp + g) \\ p &= \left(p + x \frac{dp}{dx} \right) + \left(g' \frac{dp}{dx} \right) \\ p &= p + (x + g') \frac{dp}{dx} \\ 0 &= (x + g') \frac{dp}{dx} \end{aligned}$$

Where g' is derivative of $g(p)$ w.r.t. p . The general solution is given by

$$\begin{aligned} \frac{dp}{dx} &= 0 \\ p &= c_1 \end{aligned}$$

Substituting this in (1) gives the general solution as

$$y = ac_1 + c_1^2 + c_1x$$

The singular solution is found from solving for p from

$$x + g'(p) = 0$$

And substituting the result back in (1). Since we found above that $g = pa + p^2$, then the above equation becomes

$$\begin{aligned} x + g'(p) &= x + a + 2p \\ &= 0 \end{aligned}$$

Solving the above for p results in

$$p_1 = -\frac{a}{2} - \frac{x}{2}$$

Substituting the above back in (1) results in

$$y_1 = -\frac{(x+a)^2}{4}$$

Summary

The solution(s) found are the following

$$y = ac_1 + c_1^2 + c_1x \quad (1)$$

$$y = -\frac{(x+a)^2}{4} \quad (2)$$

Verification of solutions

$$y = ac_1 + c_1^2 + c_1x$$

Verified OK.

$$y = -\frac{(x+a)^2}{4}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:
*** Sublevel 2 ***
Methods for first order ODEs:
-> Solving 1st order ODE of high degree, 1st attempt
trying 1st order WeierstrassP solution for high degree ODE
trying 1st order WeierstrassPPrime solution for high degree ODE
trying 1st order JacobiSN solution for high degree ODE
trying 1st order ODE linearizable_by_differentiation
trying differential order: 1; missing variables
trying dAlembert
<- dAlembert successful`
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 20

```
dsolve(diff(y(x),x)^2+(x+a)*diff(y(x),x)-y(x)=0,y(x), singsol=all)
```

$$y(x) = -\frac{(a+x)^2}{4}$$
$$y(x) = c_1(c_1 + a + x)$$

✓ Solution by Mathematica

Time used: 0.011 (sec). Leaf size: 26

```
DSolve[y'[x]^2+(x+a)*y'[x]-y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1(a + x + c_1)$$
$$y(x) \rightarrow -\frac{1}{4}(a + x)^2$$

1.51 problem Problem 65

1.51.1 Solving as dAlembert ode 498

Internal problem ID [12162]

Internal file name [OUTPUT/10814_Thursday_September_21_2023_05_46_20_AM_97807292/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 65.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "dAlembert"

Maple gives the following as the ode type

```
[[_1st_order , _with_linear_symmetries] , _dAlembert]
```

$$y'^2 - 2y'x + y = 0$$

1.51.1 Solving as dAlembert ode

Let $p = y'$ the ode becomes

$$p^2 - 2px + y = 0$$

Solving for y from the above results in

$$y = -p^2 + 2px \tag{1A}$$

This has the form

$$y = xf(p) + g(p) \tag{*}$$

Where f, g are functions of $p = y'(x)$. The above ode is dAlembert ode which is now solved. Taking derivative of (*) w.r.t. x gives

$$\begin{aligned} p &= f + (xf' + g')\frac{dp}{dx} \\ p - f &= (xf' + g')\frac{dp}{dx} \end{aligned} \tag{2}$$

Comparing the form $y = xf + g$ to (1A) shows that

$$\begin{aligned}f &= 2p \\g &= -p^2\end{aligned}$$

Hence (2) becomes

$$-p = (2x - 2p)p'(x) \tag{2A}$$

The singular solution is found by setting $\frac{dp}{dx} = 0$ in the above which gives

$$-p = 0$$

Solving for p from the above gives

$$p = 0$$

Substituting these in (1A) gives

$$y = 0$$

The general solution is found when $\frac{dp}{dx} \neq 0$. From eq. (2A). This results in

$$p'(x) = -\frac{p(x)}{2x - 2p(x)} \tag{3}$$

This ODE is now solved for $p(x)$.

Inverting the above ode gives

$$\frac{d}{dp}x(p) = -\frac{2x(p) - 2p}{p} \tag{4}$$

This ODE is now solved for $x(p)$.

Entering Linear first order ODE solver. In canonical form a linear first order is

$$\frac{d}{dp}x(p) + p(p)x(p) = q(p)$$

Where here

$$\begin{aligned}p(p) &= \frac{2}{p} \\q(p) &= 2\end{aligned}$$

Hence the ode is

$$\frac{d}{dp}x(p) + \frac{2x(p)}{p} = 2$$

The integrating factor μ is

$$\begin{aligned}\mu &= e^{\int \frac{2}{p} dp} \\ &= p^2\end{aligned}$$

The ode becomes

$$\begin{aligned}\frac{d}{dp}(\mu x) &= (\mu)(2) \\ \frac{d}{dp}(p^2 x) &= (p^2)(2) \\ d(p^2 x) &= (2p^2) dp\end{aligned}$$

Integrating gives

$$\begin{aligned}p^2 x &= \int 2p^2 dp \\ p^2 x &= \frac{2p^3}{3} + c_1\end{aligned}$$

Dividing both sides by the integrating factor $\mu = p^2$ results in

$$x(p) = \frac{2p}{3} + \frac{c_1}{p^2}$$

Now we need to eliminate p between the above and (1A). One way to do this is by solving (1) for p . This results in

$$\begin{aligned}p &= x + \sqrt{x^2 - y} \\ p &= x - \sqrt{x^2 - y}\end{aligned}$$

Substituting the above in the solution for x found above gives

$$\begin{aligned}x &= \frac{(8x^2 - 2y)\sqrt{x^2 - y} + 8x^3 - 6yx + 3c_1}{3(x + \sqrt{x^2 - y})^2} \\ x &= \frac{(-8x^2 + 2y)\sqrt{x^2 - y} + 8x^3 - 6yx + 3c_1}{3(x - \sqrt{x^2 - y})^2}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = 0 \tag{1}$$

$$x = \frac{(8x^2 - 2y)\sqrt{x^2 - y} + 8x^3 - 6yx + 3c_1}{3(x + \sqrt{x^2 - y})^2} \tag{2}$$

$$x = \frac{(-8x^2 + 2y)\sqrt{x^2 - y} + 8x^3 - 6yx + 3c_1}{3(x - \sqrt{x^2 - y})^2} \tag{3}$$

Verification of solutions

$$y = 0$$

Verified OK.

$$x = \frac{(8x^2 - 2y)\sqrt{x^2 - y} + 8x^3 - 6yx + 3c_1}{3(x + \sqrt{x^2 - y})^2}$$

Verified OK.

$$x = \frac{(-8x^2 + 2y)\sqrt{x^2 - y} + 8x^3 - 6yx + 3c_1}{3(x - \sqrt{x^2 - y})^2}$$

Verified OK.

Maple trace

```
`Methods for first order ODEs:  
-> Solving 1st order ODE of high degree, 1st attempt  
trying 1st order WeierstrassP solution for high degree ODE  
trying 1st order WeierstrassPPrime solution for high degree ODE  
trying 1st order JacobiSN solution for high degree ODE  
trying 1st order ODE linearizable_by_differentiation  
trying differential order: 1; missing variables  
trying dAlembert  
<- dAlembert successful`
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 611

```
dsolve(diff(y(x),x)^2-2*x*diff(y(x),x)+y(x)=0,y(x), singsol=all)
```

$$y(x) = \frac{\left(x^2 + x\left(x^3 + 2\sqrt{3}\sqrt{-c_1(x^3 - 3c_1)} - 6c_1\right)^{\frac{1}{3}} + \left(x^3 + 2\sqrt{3}\sqrt{-c_1(x^3 - 3c_1)} - 6c_1\right)^{\frac{2}{3}}\right)\left(x^2 - 3x\left(x^3 + 2\sqrt{3}\sqrt{-c_1(x^3 - 3c_1)} - 6c_1\right)^{\frac{1}{3}} - 4\left(x^3 + 2\sqrt{3}\sqrt{-c_1(x^3 - 3c_1)} - 6c_1\right)^{\frac{2}{3}}\right)}{4\left(x^3 + 2\sqrt{3}\sqrt{-c_1(x^3 - 3c_1)} - 6c_1\right)^{\frac{2}{3}}}$$

$$y(x) = \frac{\left(i\sqrt{3}\left(x^3 + 2\sqrt{3}\sqrt{-c_1(x^3 - 3c_1)} - 6c_1\right)^{\frac{2}{3}} - i\sqrt{3}x^2 + \left(x^3 + 2\sqrt{3}\sqrt{-c_1(x^3 - 3c_1)} - 6c_1\right)^{\frac{2}{3}} - 2x\left(x^3 + 2\sqrt{3}\sqrt{-c_1(x^3 - 3c_1)} - 6c_1\right)^{\frac{1}{3}}\right)\left(x^2 - 3x\left(x^3 + 2\sqrt{3}\sqrt{-c_1(x^3 - 3c_1)} - 6c_1\right)^{\frac{1}{3}} - 4\left(x^3 + 2\sqrt{3}\sqrt{-c_1(x^3 - 3c_1)} - 6c_1\right)^{\frac{2}{3}}\right)}{4\left(x^3 + 2\sqrt{3}\sqrt{-c_1(x^3 - 3c_1)} - 6c_1\right)^{\frac{2}{3}}}$$

$$y(x) = \frac{\left(i\sqrt{3}x^2 - i\sqrt{3}\left(x^3 + 2\sqrt{3}\sqrt{-c_1(x^3 - 3c_1)} - 6c_1\right)^{\frac{2}{3}} + x^2 - 2x\left(x^3 + 2\sqrt{3}\sqrt{-c_1(x^3 - 3c_1)} - 6c_1\right)^{\frac{1}{3}}\right)\left(x^2 - 3x\left(x^3 + 2\sqrt{3}\sqrt{-c_1(x^3 - 3c_1)} - 6c_1\right)^{\frac{1}{3}} - 4\left(x^3 + 2\sqrt{3}\sqrt{-c_1(x^3 - 3c_1)} - 6c_1\right)^{\frac{2}{3}}\right)}{4\left(x^3 + 2\sqrt{3}\sqrt{-c_1(x^3 - 3c_1)} - 6c_1\right)^{\frac{2}{3}}}$$

✓ Solution by Mathematica

Time used: 60.164 (sec). Leaf size: 954

`DSolve[y'[x]^2-2*x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]`

$$y(x) \rightarrow \frac{1}{4} \left(x^2 + \frac{x(x^3 + 8e^{3c_1})}{\sqrt[3]{x^6 - 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3} - 8e^{6c_1}}} + \sqrt[3]{x^6 - 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3} - 8e^{6c_1}} \right)$$

$$y(x) \rightarrow \frac{1}{72} \left(18x^2 - \frac{9i(\sqrt{3} - i)x(x^3 + 8e^{3c_1})}{\sqrt[3]{x^6 - 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3} - 8e^{6c_1}}} + 9i(\sqrt{3} + i) \sqrt[3]{x^6 - 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3} - 8e^{6c_1}} \right)$$

$$y(x) \rightarrow \frac{1}{72} \left(18x^2 + \frac{9i(\sqrt{3} + i)x(x^3 + 8e^{3c_1})}{\sqrt[3]{x^6 - 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3} - 8e^{6c_1}}} - 9(1 + i\sqrt{3}) \sqrt[3]{x^6 - 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(-x^3 + e^{3c_1})^3} - 8e^{6c_1}} \right)$$

$$y(x) \rightarrow \frac{x^4 + (x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(x^3 + e^{3c_1})^3} - 8e^{6c_1})^{2/3} + x^2 \sqrt[3]{x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(x^3 + e^{3c_1})^3} - 8e^{6c_1}}}{4 \sqrt[3]{x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(x^3 + e^{3c_1})^3} - 8e^{6c_1}}}$$

$$y(x) \rightarrow \frac{1}{72} \left(18x^2 + \frac{9(1 + i\sqrt{3})x(-x^3 + 8e^{3c_1})}{\sqrt[3]{x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(x^3 + e^{3c_1})^3} - 8e^{6c_1}}} + 9i(\sqrt{3} + i) \sqrt[3]{x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(x^3 + e^{3c_1})^3} - 8e^{6c_1}} \right)$$

$$y(x) \rightarrow \frac{1}{72} \left(18x^2 + \frac{9i(\sqrt{3} + i)x(x^3 - 8e^{3c_1})}{\sqrt[3]{x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(x^3 + e^{3c_1})^3} - 8e^{6c_1}}} - 9(1 + i\sqrt{3}) \sqrt[3]{x^6 + 20e^{3c_1}x^3 + 8\sqrt{e^{3c_1}(x^3 + e^{3c_1})^3} - 8e^{6c_1}} \right)$$

1.52 problem Problem 66

1.52.1 Maple step by step solution 506

Internal problem ID [12163]

Internal file name [OUTPUT/10815_Thursday_September_21_2023_05_47_13_AM_95217676/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 1, First-Order Differential Equations. Problems page 88

Problem number: Problem 66.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "exact", "linear", "separable", "homogeneousTypeD2", "first_order_ode_lie_symmetry_lookup"

Maple gives the following as the ode type

[_separable]

$$y'^2 + 2yy' \cot(x) - y^2 = 0$$

Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \frac{\left(-1 + \sqrt{\tan(x)^2 + 1}\right) y}{\tan(x)} \quad (1)$$

$$y' = -\frac{\left(1 + \sqrt{\tan(x)^2 + 1}\right) y}{\tan(x)} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{\left(-1 + \sqrt{\tan(x)^2 + 1}\right) y}{\tan(x)} \end{aligned}$$

Where $f(x) = \frac{-1 + \sqrt{\tan(x)^2 + 1}}{\tan(x)}$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned} \frac{1}{y} dy &= \frac{-1 + \sqrt{\tan(x)^2 + 1}}{\tan(x)} dx \\ \int \frac{1}{y} dy &= \int \frac{-1 + \sqrt{\tan(x)^2 + 1}}{\tan(x)} dx \\ \ln(y) &= \cos(x) \ln(\csc(x) - \cot(x)) \sqrt{\sec(x)^2} - \ln(\csc(x) - \cot(x)) + \ln\left(\frac{2}{\cos(x) + 1}\right) + c_1 \\ y &= e^{\cos(x) \ln(\csc(x) - \cot(x)) \sqrt{\sec(x)^2} - \ln(\csc(x) - \cot(x)) + \ln\left(\frac{2}{\cos(x) + 1}\right) + c_1} \\ &= c_1 e^{\cos(x) \ln(\csc(x) - \cot(x)) \sqrt{\sec(x)^2} - \ln(\csc(x) - \cot(x)) + \ln\left(\frac{2}{\cos(x) + 1}\right)} \end{aligned}$$

Which simplifies to

$$y = \frac{2c_1 (\csc(x) - \cot(x))^{\cos(x) \sqrt{\sec(x)^2}}}{(\csc(x) - \cot(x)) (\cos(x) + 1)}$$

Summary

The solution(s) found are the following

$$y = \frac{2c_1 (\csc(x) - \cot(x))^{\cos(x) \sqrt{\sec(x)^2}}}{(\csc(x) - \cot(x)) (\cos(x) + 1)} \quad (1)$$

Verification of solutions

$$y = \frac{2c_1 (\csc(x) - \cot(x))^{\cos(x) \sqrt{\sec(x)^2}}}{(\csc(x) - \cot(x)) (\cos(x) + 1)}$$

Verified OK.

Solving equation (2)

In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= -\frac{\left(1 + \sqrt{\tan(x)^2 + 1}\right) y}{\tan(x)} \end{aligned}$$

Where $f(x) = -\frac{1+\sqrt{\tan(x)^2+1}}{\tan(x)}$ and $g(y) = y$. Integrating both sides gives

$$\frac{1}{y} dy = -\frac{1 + \sqrt{\tan(x)^2 + 1}}{\tan(x)} dx$$

$$\int \frac{1}{y} dy = \int -\frac{1 + \sqrt{\tan(x)^2 + 1}}{\tan(x)} dx$$

$$\ln(y) = -\cos(x) \ln(\csc(x) - \cot(x)) \sqrt{\sec(x)^2} - \ln(\csc(x) - \cot(x)) + \ln\left(\frac{2}{\cos(x) + 1}\right) + c_2$$

$$y = e^{-\cos(x) \ln(\csc(x) - \cot(x)) \sqrt{\sec(x)^2} - \ln(\csc(x) - \cot(x)) + \ln\left(\frac{2}{\cos(x) + 1}\right) + c_2}$$

$$= c_2 e^{-\cos(x) \ln(\csc(x) - \cot(x)) \sqrt{\sec(x)^2} - \ln(\csc(x) - \cot(x)) + \ln\left(\frac{2}{\cos(x) + 1}\right)}$$

Which simplifies to

$$y = \frac{2c_2(\csc(x) - \cot(x))^{-\cos(x)\sqrt{\sec(x)^2}}}{(\csc(x) - \cot(x))(\cos(x) + 1)}$$

Summary

The solution(s) found are the following

$$y = \frac{2c_2(\csc(x) - \cot(x))^{-\cos(x)\sqrt{\sec(x)^2}}}{(\csc(x) - \cot(x))(\cos(x) + 1)} \quad (1)$$

Verification of solutions

$$y = \frac{2c_2(\csc(x) - \cot(x))^{-\cos(x)\sqrt{\sec(x)^2}}}{(\csc(x) - \cot(x))(\cos(x) + 1)}$$

Verified OK.

1.52.1 Maple step by step solution

Let's solve

$$y'^2 + 2yy' \cot(x) - y^2 = 0$$

- Highest derivative means the order of the ODE is 1

y'

- Separate variables

$$\frac{y'}{y} = \frac{-1 + \sqrt{\tan(x)^2 + 1}}{\tan(x)}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{y} dx = \int \frac{-1 + \sqrt{\tan(x)^2 + 1}}{\tan(x)} dx + c_1$$

- Evaluate integral

$$\ln(y) = -\operatorname{arctanh}\left(\frac{1}{\sqrt{\tan(x)^2 + 1}}\right) - \ln(\tan(x)) + \frac{\ln(\tan(x)^2 + 1)}{2} + c_1$$

- Solve for y

$$y = \frac{e^{c_1} \cos(x) \left(-1 + \sqrt{\frac{1}{\cos(x)^2}}\right)}{\sqrt{\sin(x)^2} \sin(x)}$$

Maple trace

```

`Methods for first order ODEs:
-> Solving 1st order ODE of high degree, 1st attempt
trying 1st order WeierstrassP solution for high degree ODE
trying 1st order WeierstrassPPrime solution for high degree ODE
trying 1st order JacobiSN solution for high degree ODE
trying 1st order ODE linearizable_by_differentiation
trying differential order: 1; missing variables
trying simple symmetries for implicit equations
<- symmetries for implicit equations successful`

```

✓ Solution by Maple

Time used: 0.125 (sec). Leaf size: 39

```
dsolve(diff(y(x),x)^2+2*y(x)*diff(y(x),x)*cot(x)-y(x)^2=0,y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = \frac{\operatorname{csgn}(\sin(x)) c_1}{\cos(x) + \operatorname{csgn}(\sec(x))}$$

$$y(x) = \csc(x)^2 (\cos(x) + \operatorname{csgn}(\sec(x))) \operatorname{csgn}(\sin(x)) c_1$$

✓ Solution by Mathematica

Time used: 0.251 (sec). Leaf size: 36

```
DSolve[y'[x]^2+2*y[x]*y'[x]*Cot[x]-y[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \csc^2\left(\frac{x}{2}\right)$$

$$y(x) \rightarrow c_1 \sec^2\left(\frac{x}{2}\right)$$

$$y(x) \rightarrow 0$$

2 Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER.

Problems page 172

2.1	problem Problem 1	511
2.2	problem Problem 2	524
2.3	problem Problem 3	535
2.4	problem Problem 4	540
2.5	problem Problem 5	553
2.6	problem Problem 6	583
2.7	problem Problem 7	597
2.8	problem Problem 8	601
2.9	problem Problem 9	614
2.10	problem Problem 10	618
2.11	problem Problem 11	624
2.12	problem Problem 12	633
2.13	problem Problem 13	636
2.14	problem Problem 14	640
2.15	problem Problem 15	644
2.16	problem Problem 16	653
2.17	problem Problem 17	657
2.18	problem Problem 18	664
2.19	problem Problem 19	671
2.20	problem Problem 20	684
2.21	problem Problem 30	698
2.22	problem Problem 31	702
2.23	problem Problem 32	715
2.24	problem Problem 33	726
2.25	problem Problem 34	741
2.26	problem Problem 35	749
2.27	problem Problem 36	760
2.28	problem Problem 40(a)	770
2.29	problem Problem 40(b)	773
2.30	problem Problem 41	777
2.31	problem Problem 42	781
2.32	problem Problem 43	787
2.33	problem Problem 47	807
2.34	problem Problem 49	810

2.35	problem Problem 50	821
2.36	problem Problem 51	832
2.37	problem Problem 52	848
2.38	problem Problem 53	852
2.39	problem Problem 54	856
2.40	problem Problem 55	860
2.41	problem Problem 56	862
2.42	problem Problem 57	865
2.43	problem Problem 58	876
2.44	problem Problem 59	887

2.1 problem Problem 1

2.1.1	Existence and uniqueness analysis	511
2.1.2	Solving as second order linear constant coeff ode	512
2.1.3	Solving using Kovacic algorithm	516
2.1.4	Maple step by step solution	521

Internal problem ID [12164]

Internal file name [OUTPUT/10816_Thursday_September_21_2023_05_47_17_AM_10735052/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 1.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$y'' - 6y' + 10y = 100$$

With initial conditions

$$[y(0) = 10, y'(0) = 5]$$

2.1.1 Existence and uniqueness analysis

This is a linear ODE. In canonical form it is written as

$$y'' + p(x)y' + q(x)y = F$$

Where here

$$p(x) = -6$$

$$q(x) = 10$$

$$F = 100$$

Hence the ode is

$$y'' - 6y' + 10y = 100$$

The domain of $p(x) = -6$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is inside this domain. The domain of $q(x) = 10$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. The domain of $F = 100$ is

$$\{-\infty < x < \infty\}$$

And the point $x_0 = 0$ is also inside this domain. Hence solution exists and is unique.

2.1.2 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -6, C = 10, f(x) = 100$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 6y' + 10y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -6, C = 10$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 6\lambda e^{\lambda x} + 10e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 6\lambda + 10 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -6, C = 10$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{6}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-6^2 - (4)(1)(10)} \\ &= 3 \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= 3 + i \\ \lambda_2 &= 3 - i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= 3 + i \\ \lambda_2 &= 3 - i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 3$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^{3x} (c_1 \cos(x) + c_2 \sin(x))$$

Therefore the homogeneous solution y_h is

$$y_h = e^{3x} (c_1 \cos(x) + c_2 \sin(x))$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

1

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x) e^{3x}, e^{3x} \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$10A_1 = 100$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 10]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 10$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (e^{3x}(c_1 \cos(x) + c_2 \sin(x))) + (10) \end{aligned}$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{3x}(c_1 \cos(x) + c_2 \sin(x)) + 10 \tag{1}$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 10$ and $x = 0$ in the above gives

$$10 = c_1 + 10 \tag{1A}$$

Taking derivative of the solution gives

$$y' = 3e^{3x}(c_1 \cos(x) + c_2 \sin(x)) + e^{3x}(-\sin(x)c_1 + c_2 \cos(x))$$

substituting $y' = 5$ and $x = 0$ in the above gives

$$5 = 3c_1 + c_2 \tag{2A}$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

$$c_2 = 5$$


Substituting these values back in above solution results in

$$y = 10 + 5e^{3x} \sin(x)$$


Summary

The solution(s) found are the following

$$y = 10 + 5e^{3x} \sin(x) \tag{1}$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 10 + 5e^{3x} \sin(x)$$

Verified OK.

2.1.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 6y' + 10y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -6 \\ C &= 10 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 58: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-6}{1} dx} \end{aligned}$$

$$\begin{aligned}
&= z_1 e^{3x} \\
&= z_1 (e^{3x})
\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x) e^{3x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}
y_2 &= y_1 \int \frac{e^{\int -\frac{-6}{1} dx}}{(y_1)^2} dx \\
&= y_1 \int \frac{e^{6x}}{(y_1)^2} dx \\
&= y_1 (\tan(x))
\end{aligned}$$

Therefore the solution is

$$\begin{aligned}
y &= c_1 y_1 + c_2 y_2 \\
&= c_1 (\cos(x) e^{3x}) + c_2 (\cos(x) e^{3x} (\tan(x)))
\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 6y' + 10y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) e^{3x} + e^{3x} \sin(x) c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x) e^{3x}, e^{3x} \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$10A_1 = 100$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 10]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = 10$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) e^{3x} + e^{3x} \sin(x) c_2) + (10) \end{aligned}$$

Which simplifies to

$$y = e^{3x}(c_1 \cos(x) + c_2 \sin(x)) + 10$$

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$y = e^{3x}(c_1 \cos(x) + c_2 \sin(x)) + 10 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 10$ and $x = 0$ in the above gives

$$10 = c_1 + 10 \quad (1A)$$

Taking derivative of the solution gives

$$y' = 3e^{3x}(c_1 \cos(x) + c_2 \sin(x)) + e^{3x}(-\sin(x)c_1 + c_2 \cos(x))$$

substituting $y' = 5$ and $x = 0$ in the above gives

$$5 = 3c_1 + c_2 \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. Solving for the constants gives

$$c_1 = 0$$

$$c_2 = 5$$


Substituting these values back in above solution results in

$$y = 10 + 5e^{3x} \sin(x)$$


Summary

The solution(s) found are the following

$$y = 10 + 5e^{3x} \sin(x) \quad (1)$$



(a) Solution plot



(b) Slope field plot

Verification of solutions

$$y = 10 + 5 e^{3x} \sin(x)$$

Verified OK.

2.1.4 Maple step by step solution

Let's solve

$$\left[y'' - 6y' + 10y = 100, y(0) = 10, y'|_{\{x=0\}} = 5 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 6r + 10 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{6 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (3 - I, 3 + I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x) e^{3x}$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^{3x} \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(x) e^{3x} + e^{3x} \sin(x) c_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = 100 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) e^{3x} & e^{3x} \sin(x) \\ -e^{3x} \sin(x) + 3 \cos(x) e^{3x} & 3 e^{3x} \sin(x) + \cos(x) e^{3x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{6x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -100 e^{3x} (\cos(x) (\int e^{-3x} \sin(x) dx) - \sin(x) (\int e^{-3x} \cos(x) dx))$$

- Compute integrals

$$y_p(x) = 10$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(x) e^{3x} + e^{3x} \sin(x) c_2 + 10$$

- Check validity of solution $y = c_1 \cos(x) e^{3x} + e^{3x} \sin(x) c_2 + 10$

- Use initial condition $y(0) = 10$

$$10 = c_1 + 10$$

- Compute derivative of the solution

$$y' = -c_1 \sin(x) e^{3x} + 3c_1 \cos(x) e^{3x} + 3 e^{3x} \sin(x) c_2 + e^{3x} \cos(x) c_2$$

- Use the initial condition $y' \Big|_{\{x=0\}} = 5$

$$5 = 3c_1 + c_2$$

- Solve for c_1 and c_2

$$\{c_1 = 0, c_2 = 5\}$$

- Substitute constant values into general solution and simplify

$$y = 10 + 5 e^{3x} \sin(x)$$

- Solution to the IVP

$$y = 10 + 5 e^{3x} \sin(x)$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 14

```
dsolve([diff(y(x),x$2)-6*diff(y(x),x)+10*y(x)=100,y(0) = 10, D(y)(0) = 5],y(x), singsol=all)
```

$$y(x) = 5e^{3x} \sin(x) + 10$$

✓ Solution by Mathematica

Time used: 0.029 (sec). Leaf size: 17

```
DSolve[{y''[x]-6*y'[x]+10*y[x]==100,{y[0]==10,y'[0]==5}},y[x],x,IncludeSingularSolutions ->
```

$$y(x) \rightarrow 5(e^{3x} \sin(x) + 2)$$

2.2 problem Problem 2

2.2.1	Solving as second order linear constant coeff ode	524
2.2.2	Solving using Kovacic algorithm	528
2.2.3	Maple step by step solution	532

Internal problem ID [12165]

Internal file name [OUTPUT/10817_Thursday_September_21_2023_05_47_20_AM_99087045/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 2.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$x'' + x = \sin(t) - \cos(2t)$$

2.2.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ax''(t) + Bx'(t) + Cx(t) = f(t)$$

Where $A = 1, B = 0, C = 1, f(t) = \sin(t) - \cos(2t)$. Let the solution be

$$x = x_h + x_p$$

Where x_h is the solution to the homogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = 0$, and x_p is a particular solution to the non-homogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = f(t)$. x_h is the solution to

$$x'' + x = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ax''(t) + Bx'(t) + Cx(t) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $x = e^{\lambda t}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda t} + e^{\lambda t} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda t}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$x = e^{\alpha t} (c_1 \cos(\beta t) + c_2 \sin(\beta t))$$

Which becomes

$$x = e^0 (c_1 \cos(t) + c_2 \sin(t))$$

Or

$$x = c_1 \cos(t) + c_2 \sin(t)$$

Therefore the homogeneous solution x_h is

$$x_h = c_1 \cos(t) + c_2 \sin(t)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(t) - \cos(2t)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(t), \sin(t)\}, \{\cos(2t), \sin(2t)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(t), \sin(t)\}$$

Since $\cos(t)$ is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{\cos(t)t, \sin(t)t\}, \{\cos(2t), \sin(2t)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$x_p = A_1 \cos(t)t + A_2 \sin(t)t + A_3 \cos(2t) + A_4 \sin(2t)$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution x_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(t) + 2A_2 \cos(t) - 3A_3 \cos(2t) - 3A_4 \sin(2t) = \sin(t) - \cos(2t)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0, A_3 = \frac{1}{3}, A_4 = 0 \right]$$

Substituting the above back in the above trial solution x_p , gives the particular solution

$$x_p = -\frac{\cos(t)t}{2} + \frac{\cos(2t)}{3}$$

Therefore the general solution is

$$\begin{aligned}x &= x_h + x_p \\ &= (c_1 \cos(t) + c_2 \sin(t)) + \left(-\frac{\cos(t)t}{2} + \frac{\cos(2t)}{3}\right)\end{aligned}$$

Summary

The solution(s) found are the following

$$x = c_1 \cos(t) + c_2 \sin(t) - \frac{\cos(t)t}{2} + \frac{\cos(2t)}{3} \quad (1)$$




Figure 101: Slope field plot

Verification of solutions

$$x = c_1 \cos(t) + c_2 \sin(t) - \frac{\cos(t)t}{2} + \frac{\cos(2t)}{3}$$

Verified OK.

2.2.2 Solving using Kovacic algorithm

Writing the ode as

$$x'' + x = 0 \quad (1)$$

$$Ax'' + Bx' + Cx = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = xe^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = -z(t) \quad (7)$$

Equation (7) is now solved. After finding $z(t)$ then x is found using the inverse transformation

$$x = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 60: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of t , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(t) = \cos(t)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in x is found from

$$x_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}x_1 &= z_1 \\ &= \cos(t)\end{aligned}$$

Which simplifies to

$$x_1 = \cos(t)$$

The second solution x_2 to the original ode is found using reduction of order

$$x_2 = x_1 \int \frac{e^{\int -\frac{B}{A} dt}}{x_1^2} dt$$

Since $B = 0$ then the above becomes

$$\begin{aligned}x_2 &= x_1 \int \frac{1}{x_1^2} dt \\ &= \cos(t) \int \frac{1}{\cos(t)^2} dt \\ &= \cos(t) (\tan(t))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}x &= c_1 x_1 + c_2 x_2 \\ &= c_1 (\cos(t)) + c_2 (\cos(t) (\tan(t)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$x = x_h + x_p$$

Where x_h is the solution to the homogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = 0$, and x_p is a particular solution to the nonhomogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = f(t)$.
 x_h is the solution to

$$x'' + x = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$x_h = c_1 \cos(t) + c_2 \sin(t)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(t) - \cos(2t)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(t), \sin(t)\}, \{\cos(2t), \sin(2t)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(t), \sin(t)\}$$

Since $\cos(t)$ is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{\cos(t)t, \sin(t)t\}, \{\cos(2t), \sin(2t)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$x_p = A_1 \cos(t)t + A_2 \sin(t)t + A_3 \cos(2t) + A_4 \sin(2t)$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution x_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-2A_1 \sin(t) + 2A_2 \cos(t) - 3A_3 \cos(2t) - 3A_4 \sin(2t) = \sin(t) - \cos(2t)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = 0, A_3 = \frac{1}{3}, A_4 = 0 \right]$$

Substituting the above back in the above trial solution x_p , gives the particular solution

$$x_p = -\frac{\cos(t)t}{2} + \frac{\cos(2t)}{3}$$

Therefore the general solution is

$$\begin{aligned} x &= x_h + x_p \\ &= (c_1 \cos(t) + c_2 \sin(t)) + \left(-\frac{\cos(t)t}{2} + \frac{\cos(2t)}{3} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$x = c_1 \cos(t) + c_2 \sin(t) - \frac{\cos(t)t}{2} + \frac{\cos(2t)}{3} \quad (1)$$




Figure 102: Slope field plot

Verification of solutions

$$x = c_1 \cos(t) + c_2 \sin(t) - \frac{\cos(t)t}{2} + \frac{\cos(2t)}{3}$$

Verified OK.

2.2.3 Maple step by step solution

Let's solve

$$x'' + x = \sin(t) - \cos(2t)$$

- Highest derivative means the order of the ODE is 2
 x''
- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$x_1(t) = \cos(t)$$

- 2nd solution of the homogeneous ODE

$$x_2(t) = \sin(t)$$

- General solution of the ODE

$$x = c_1 x_1(t) + c_2 x_2(t) + x_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$x = c_1 \cos(t) + c_2 \sin(t) + x_p(t)$$

- Find a particular solution $x_p(t)$ of the ODE

- Use variation of parameters to find x_p here $f(t)$ is the forcing function

$$\left[x_p(t) = -x_1(t) \left(\int \frac{x_2(t)f(t)}{W(x_1(t), x_2(t))} dt \right) + x_2(t) \left(\int \frac{x_1(t)f(t)}{W(x_1(t), x_2(t))} dt \right), f(t) = \sin(t) - \cos(2t) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(x_1(t), x_2(t)) = \begin{bmatrix} \cos(t) & \sin(t) \\ -\sin(t) & \cos(t) \end{bmatrix}$$

- Compute Wronskian

$$W(x_1(t), x_2(t)) = 1$$

- Substitute functions into equation for $x_p(t)$

$$x_p(t) = -\cos(t) \left(\int \sin(t) (\sin(t) - 2\cos(t)^2 + 1) dt \right) + \sin(t) \left(\int \cos(t) (\sin(t) - \cos(2t)) dt \right)$$

- Compute integrals

$$x_p(t) = \frac{2\cos(t)^2}{3} - \frac{\cos(t)t}{2} + \frac{\sin(t)}{4} - \frac{1}{3}$$

- Substitute particular solution into general solution to ODE

$$x = c_1 \cos(t) + c_2 \sin(t) + \frac{2\cos(t)^2}{3} - \frac{\cos(t)t}{2} + \frac{\sin(t)}{4} - \frac{1}{3}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(x(t),t$2)+x(t)=sin(t)-cos(2*t),x(t), singsol=all)
```

$$x(t) = \frac{\cos(2t)}{3} + \frac{(-t + 2c_1)\cos(t)}{2} + \frac{(1 + 4c_2)\sin(t)}{4}$$

✓ Solution by Mathematica

Time used: 0.076 (sec). Leaf size: 30

```
DSolve[x''[t]+x[t]==Sin[t]-Cos[2*t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \frac{1}{3} \cos(2t) + \left(-\frac{t}{2} + c_1\right) \cos(t) + c_2 \sin(t)$$

2.3 problem Problem 3

2.3.1 Maple step by step solution 536

Internal problem ID [12166]

Internal file name [OUTPUT/10818_Thursday_September_21_2023_05_47_23_AM_40160976/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 3.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _missing_x]]
```

$$y' + y''' - 3y'' = 0$$

The characteristic equation is

$$\lambda^3 - 3\lambda^2 + \lambda = 0$$

The roots of the above equation are

$$\begin{aligned}\lambda_1 &= 0 \\ \lambda_2 &= \frac{3}{2} + \frac{\sqrt{5}}{2} \\ \lambda_3 &= \frac{3}{2} - \frac{\sqrt{5}}{2}\end{aligned}$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 + e^{\left(\frac{3}{2} - \frac{\sqrt{5}}{2}\right)x} c_2 + e^{\left(\frac{3}{2} + \frac{\sqrt{5}}{2}\right)x} c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned}y_1 &= 1 \\ y_2 &= e^{\left(\frac{3}{2} - \frac{\sqrt{5}}{2}\right)x} \\ y_3 &= e^{\left(\frac{3}{2} + \frac{\sqrt{5}}{2}\right)x}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 + e^{\left(\frac{3}{2} - \frac{\sqrt{5}}{2}\right)x} c_2 + e^{\left(\frac{3}{2} + \frac{\sqrt{5}}{2}\right)x} c_3 \quad (1)$$

Verification of solutions

$$y = c_1 + e^{\left(\frac{3}{2} - \frac{\sqrt{5}}{2}\right)x} c_2 + e^{\left(\frac{3}{2} + \frac{\sqrt{5}}{2}\right)x} c_3$$

Verified OK.

2.3.1 Maple step by step solution

Let's solve

$$y' + y''' - 3y'' = 0$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = -y_2(x) + 3y_3(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = -y_2(x) + 3y_3(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 3 \end{bmatrix} \cdot \vec{y}(x)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & -1 & 3 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x)$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\begin{array}{c} \left[\begin{array}{c} 1 \\ 0 \\ 0 \end{array} \right] \\ 0, \left[\begin{array}{c} 1 \\ 0 \\ 0 \end{array} \right] \end{array} \right], \left[\begin{array}{c} \frac{3}{2} - \frac{\sqrt{5}}{2}, \left[\begin{array}{c} \frac{1}{\left(\frac{3}{2} - \frac{\sqrt{5}}{2}\right)^2} \\ \frac{1}{\frac{3}{2} - \frac{\sqrt{5}}{2}} \\ 1 \end{array} \right] \end{array} \right], \left[\begin{array}{c} \frac{3}{2} + \frac{\sqrt{5}}{2}, \left[\begin{array}{c} \frac{1}{\left(\frac{3}{2} + \frac{\sqrt{5}}{2}\right)^2} \\ \frac{1}{\frac{3}{2} + \frac{\sqrt{5}}{2}} \\ 1 \end{array} \right] \end{array} \right] \right]$$

- Consider eigenpair

$$\left[\begin{array}{c} \left[\begin{array}{c} 1 \\ 0 \\ 0 \end{array} \right] \\ 0, \left[\begin{array}{c} 1 \\ 0 \\ 0 \end{array} \right] \end{array} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

- Consider eigenpair

$$\left[\begin{array}{c} \frac{3}{2} - \frac{\sqrt{5}}{2}, \left[\begin{array}{c} \frac{1}{\left(\frac{3}{2} - \frac{\sqrt{5}}{2}\right)^2} \\ \frac{1}{\frac{3}{2} - \frac{\sqrt{5}}{2}} \\ 1 \end{array} \right] \end{array} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^{\left(\frac{3}{2} - \frac{\sqrt{5}}{2}\right)x} \cdot \begin{bmatrix} \frac{1}{\left(\frac{3}{2} - \frac{\sqrt{5}}{2}\right)^2} \\ \frac{1}{\frac{3}{2} - \frac{\sqrt{5}}{2}} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[\frac{3}{2} + \frac{\sqrt{5}}{2}, \begin{bmatrix} \frac{1}{\left(\frac{3}{2} + \frac{\sqrt{5}}{2}\right)^2} \\ \frac{1}{\frac{3}{2} + \frac{\sqrt{5}}{2}} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_3 = e^{\left(\frac{3}{2} + \frac{\sqrt{5}}{2}\right)x} \cdot \begin{bmatrix} \frac{1}{\left(\frac{3}{2} + \frac{\sqrt{5}}{2}\right)^2} \\ \frac{1}{\frac{3}{2} + \frac{\sqrt{5}}{2}} \\ 1 \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{y} = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3$$

- Substitute solutions into the general solution

$$\vec{y} = e^{\left(\frac{3}{2} - \frac{\sqrt{5}}{2}\right)x} c_2 \cdot \begin{bmatrix} \frac{1}{\left(\frac{3}{2} - \frac{\sqrt{5}}{2}\right)^2} \\ \frac{1}{\frac{3}{2} - \frac{\sqrt{5}}{2}} \\ 1 \end{bmatrix} + e^{\left(\frac{3}{2} + \frac{\sqrt{5}}{2}\right)x} c_3 \cdot \begin{bmatrix} \frac{1}{\left(\frac{3}{2} + \frac{\sqrt{5}}{2}\right)^2} \\ \frac{1}{\frac{3}{2} + \frac{\sqrt{5}}{2}} \\ 1 \end{bmatrix} + \begin{bmatrix} c_1 \\ 0 \\ 0 \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = \frac{c_3(7-3\sqrt{5})e^{\frac{(3+\sqrt{5})x}{2}}}{2} + \frac{c_2(3\sqrt{5}+7)e^{-\frac{(\sqrt{5}-3)x}{2}}}{2} + c_1$$

Maple trace

```
`Methods for third order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve(diff(y(x),x)+diff(y(x),x$3)-3*diff(y(x),x$2)=0,y(x), singsol=all)
```

$$y(x) = c_1 + c_2 e^{\frac{(3+\sqrt{5})x}{2}} + c_3 e^{-\frac{(\sqrt{5}-3)x}{2}}$$

✓ Solution by Mathematica

Time used: 0.384 (sec). Leaf size: 57

```
DSolve[y'[x]+y'''[x]-3*y''[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} e^{-\frac{1}{2}(\sqrt{5}-3)x} \left((3 + \sqrt{5}) c_1 - (\sqrt{5} - 3) c_2 e^{\sqrt{5}x} \right) + c_3$$

2.4 problem Problem 4

2.4.1	Solving as second order linear constant coeff ode	540
2.4.2	Solving using Kovacic algorithm	545
2.4.3	Maple step by step solution	550

Internal problem ID [12167]

Internal file name [OUTPUT/10819_Thursday_September_21_2023_05_47_23_AM_3328183/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 4.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic", "second_order_linear_constant_coeff"**

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \frac{1}{\sin(x)^3}$$

2.4.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \csc(x)^3$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \csc(x)^3}{1} dx$$

Which simplifies to

$$u_1 = - \int \csc(x)^2 dx$$

Hence

$$u_1 = \cot(x)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos(x) \csc(x)^3}{1} dx$$

Which simplifies to

$$u_2 = \int \csc(x)^2 \cot(x) dx$$

Hence

$$u_2 = - \frac{\cot(x)^2}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \cos(x) \cot(x) - \frac{\cot(x)^2 \sin(x)}{2}$$

Which simplifies to

$$y_p(x) = \frac{\cos(x) \cot(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(\frac{\cos(x) \cot(x)}{2} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{\cos(x) \cot(x)}{2} \quad (1)$$




Figure 103: Slope field plot

Verification of solutions

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{\cos(x) \cot(x)}{2}$$

Verified OK.

2.4.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 63: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \csc(x)^3}{1} dx$$

Which simplifies to

$$u_1 = - \int \csc(x)^2 dx$$

Hence

$$u_1 = \cot(x)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos(x) \csc(x)^3}{1} dx$$

Which simplifies to

$$u_2 = \int \csc(x)^2 \cot(x) dx$$

Hence

$$u_2 = - \frac{\cot(x)^2}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \cos(x) \cot(x) - \frac{\cot(x)^2 \sin(x)}{2}$$

Which simplifies to

$$y_p(x) = \frac{\cos(x) \cot(x)}{2}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(\frac{\cos(x) \cot(x)}{2} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{\cos(x) \cot(x)}{2} \quad (1)$$




Figure 104: Slope field plot

Verification of solutions

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{\cos(x) \cot(x)}{2}$$

Verified OK.

2.4.3 Maple step by step solution

Let's solve

$$y'' + y = \csc(x)^3$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = \csc(x)^3 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \csc(x)^2 dx \right) + \sin(x) \left(\int \csc(x)^2 \cot(x) dx \right)$$

- Compute integrals

$$y_p(x) = \frac{\cos(x)\cot(x)}{2}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{\cos(x)\cot(x)}{2}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 20

```
dsolve(diff(y(x),x$2)+y(x)=1/sin(x)^3,y(x), singsol=all)
```

$$y(x) = (c_1 + \cot(x)) \cos(x) + \sin(x) c_2 - \frac{\csc(x)}{2}$$

✓ Solution by Mathematica

Time used: 0.09 (sec). Leaf size: 25

```
DSolve[y''[x]+y[x]==1/Sin[x]^3,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{\csc(x)}{2} + c_2 \sin(x) + \cos(x)(\cot(x) + c_1)$$

2.5 problem Problem 5

2.5.1	Solving as second order euler ode	554
2.5.2	Solving as linear second order ode solved by an integrating factor ode	557
2.5.3	Solving as second order change of variable on x method 2 ode .	558
2.5.4	Solving as second order change of variable on x method 1 ode .	563
2.5.5	Solving as second order change of variable on y method 1 ode .	568
2.5.6	Solving as second order change of variable on y method 2 ode .	572
2.5.7	Solving using Kovacic algorithm	576

Internal problem ID [12168]

Internal file name [OUTPUT/10820_Thursday_September_21_2023_05_47_25_AM_597492/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 5.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_euler_ode", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2", "second_order_change_of_variable_on_y_method_1", "second_order_change_of_variable_on_y_method_2", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' - 4y'/x + 6y = 2$$

2.5.1 Solving as second order euler ode ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = -4x$, $C = 6$, $f(x) = 2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' - 4y'x + 6y = 0$$

This is Euler second order ODE. Let the solution be $y = x^r$, then $y' = rx^{r-1}$ and $y'' = r(r-1)x^{r-2}$. Substituting these back into the given ODE gives

$$x^2(r(r-1))x^{r-2} - 4rx^{r-1} + 6x^r = 0$$

Simplifying gives

$$r(r-1)x^r - 4rx^r + 6x^r = 0$$

Since $x^r \neq 0$ then dividing throughout by x^r gives

$$r(r-1) - 4r + 6 = 0$$

Or

$$r^2 - 5r + 6 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = 2$$

$$r_2 = 3$$

Since the roots are real and distinct, then the general solution is

$$y = c_1y_1 + c_2y_2$$

Where $y_1 = x^{r_1}$ and $y_2 = x^{r_2}$. Hence

$$y = c_2x^3 + c_1x^2$$

Next, we find the particular solution to the ODE

$$x^2y'' - 4y'x + 6y = 2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x^2$$

$$y_2 = x^3$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x^2 & x^3 \\ \frac{d}{dx}(x^2) & \frac{d}{dx}(x^3) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x^2 & x^3 \\ 2x & 3x^2 \end{vmatrix}$$

Therefore

$$W = (x^2)(3x^2) - (x^3)(2x)$$

Which simplifies to

$$W = x^4$$

Which simplifies to

$$W = x^4$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{2x^3}{x^6} dx$$

Which simplifies to

$$u_1 = - \int \frac{2}{x^3} dx$$

Hence

$$u_1 = \frac{1}{x^2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{2x^2}{x^6} dx$$

Which simplifies to

$$u_2 = \int \frac{2}{x^4} dx$$

Hence

$$u_2 = -\frac{2}{3x^3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{1}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \frac{1}{3} + c_2 x^3 + c_1 x^2 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{1}{3} + c_2x^3 + c_1x^2 \quad (1)$$

Verification of solutions

$$y = \frac{1}{3} + c_2x^3 + c_1x^2$$

Verified OK.

2.5.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = -\frac{4}{x}$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -\frac{4}{x} dx} \\ &= \frac{1}{x^2} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned} (M(x)y)'' &= \frac{2}{x^4} \\ \left(\frac{y}{x^2}\right)'' &= \frac{2}{x^4} \end{aligned}$$

Integrating once gives

$$\left(\frac{y}{x^2}\right)' = -\frac{2}{3x^3} + c_1$$

Integrating again gives

$$\left(\frac{y}{x^2}\right) = c_1x + \frac{1}{3x^2} + c_2$$

Hence the solution is

$$y = \frac{c_1x + \frac{1}{3x^2} + c_2}{\frac{1}{x^2}}$$

Or

$$y = c_1x^3 + c_2x^2 + \frac{1}{3}$$

Summary

The solution(s) found are the following

$$y = c_1x^3 + c_2x^2 + \frac{1}{3} \quad (1)$$

Verification of solutions

$$y = c_1x^3 + c_2x^2 + \frac{1}{3}$$

Verified OK.

2.5.3 Solving as second order change of variable on x method 2 ode

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$.
 y_h is the solution to

$$x^2y'' - 4y'x + 6y = 0$$

In normal form the ode

$$x^2y'' - 4y'x + 6y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{4}{x}$$
$$q(x) = \frac{6}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int -\frac{4}{x}dx)} dx \\ &= \int e^{4\ln(x)} dx \\ &= \int x^4 dx \\ &= \frac{x^5}{5} \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{6}{x^2} \\ &= \frac{6}{x^{10}} \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned} \frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + \frac{6y(\tau)}{x^{10}} &= 0 \end{aligned}$$

But in terms of τ

$$\frac{6}{x^{10}} = \frac{6}{25\tau^2}$$

Hence the above ode becomes

$$\frac{d^2}{d\tau^2}y(\tau) + \frac{6y(\tau)}{25\tau^2} = 0$$

The above ode is now solved for $y(\tau)$. The ode can be written as

$$25\left(\frac{d^2}{d\tau^2}y(\tau)\right)\tau^2 + 6y(\tau) = 0$$

Which shows it is a Euler ODE. This is Euler second order ODE. Let the solution be $y(\tau) = \tau^r$, then $y' = r\tau^{r-1}$ and $y'' = r(r-1)\tau^{r-2}$. Substituting these back into the given ODE gives

$$25\tau^2(r(r-1))\tau^{r-2} + 0r\tau^{r-1} + 6\tau^r = 0$$

Simplifying gives

$$25r(r-1)\tau^r + 0\tau^r + 6\tau^r = 0$$

Since $\tau^r \neq 0$ then dividing throughout by τ^r gives

$$25r(r-1) + 0 + 6 = 0$$

Or

$$25r^2 - 25r + 6 = 0 \tag{1}$$

Equation (1) is the characteristic equation. Its roots determine the form of the general solution. Using the quadratic equation the roots are

$$r_1 = \frac{2}{5}$$
$$r_2 = \frac{3}{5}$$

Since the roots are real and distinct, then the general solution is

$$y(\tau) = c_1y_1 + c_2y_2$$

Where $y_1 = \tau^{r_1}$ and $y_2 = \tau^{r_2}$. Hence

$$y(\tau) = c_1\tau^{\frac{2}{5}} + c_2\tau^{\frac{3}{5}}$$

The above solution is now transformed back to y using (6) which results in

$$y = \frac{c_15^{\frac{3}{5}}(x^5)^{\frac{2}{5}}}{5} + \frac{c_25^{\frac{2}{5}}(x^5)^{\frac{3}{5}}}{5}$$

Therefore the homogeneous solution y_h is

$$y_h = \frac{c_1 5^{\frac{3}{5}} (x^5)^{\frac{2}{5}}}{5} + \frac{c_2 5^{\frac{2}{5}} (x^5)^{\frac{3}{5}}}{5}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (x^5)^{\frac{2}{5}}$$

$$y_2 = (x^5)^{\frac{3}{5}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x^5)^{\frac{2}{5}} & (x^5)^{\frac{3}{5}} \\ \frac{d}{dx} \left((x^5)^{\frac{2}{5}} \right) & \frac{d}{dx} \left((x^5)^{\frac{3}{5}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x^5)^{\frac{2}{5}} & (x^5)^{\frac{3}{5}} \\ \frac{2x^4}{(x^5)^{\frac{3}{5}}} & \frac{3x^4}{(x^5)^{\frac{2}{5}}} \end{vmatrix}$$

Therefore

$$W = \left((x^5)^{\frac{2}{5}} \right) \left(\frac{3x^4}{(x^5)^{\frac{2}{5}}} \right) - \left((x^5)^{\frac{3}{5}} \right) \left(\frac{2x^4}{(x^5)^{\frac{3}{5}}} \right)$$

Which simplifies to

$$W = x^4$$

Which simplifies to

$$W = x^4$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{2(x^5)^{\frac{3}{5}}}{x^6} dx$$

Which simplifies to

$$u_1 = - \int \frac{2(x^5)^{\frac{3}{5}}}{x^6} dx$$

Hence

$$u_1 = \frac{(x^5)^{\frac{3}{5}}}{x^5}$$

And Eq. (3) becomes

$$u_2 = \int \frac{2(x^5)^{\frac{2}{5}}}{x^6} dx$$

Which simplifies to

$$u_2 = \int \frac{2(x^5)^{\frac{2}{5}}}{x^6} dx$$

Hence

$$u_2 = - \frac{2(x^5)^{\frac{2}{5}}}{3x^5}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{1}{3}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= \left(\frac{c_1 5^{\frac{3}{5}} (x^5)^{\frac{2}{5}}}{5} + \frac{c_2 5^{\frac{2}{5}} (x^5)^{\frac{3}{5}}}{5} \right) + \left(\frac{1}{3} \right)$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 5^{\frac{3}{5}} (x^5)^{\frac{2}{5}}}{5} + \frac{c_2 5^{\frac{2}{5}} (x^5)^{\frac{3}{5}}}{5} + \frac{1}{3} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 5^{\frac{3}{5}} (x^5)^{\frac{2}{5}}}{5} + \frac{c_2 5^{\frac{2}{5}} (x^5)^{\frac{3}{5}}}{5} + \frac{1}{3}$$

Verified OK.

2.5.4 Solving as second order change of variable on x method 1 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = -4x$, $C = 6$, $f(x) = 2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2 y'' - 4y'x + 6y = 0$$

In normal form the ode

$$x^2 y'' - 4y'x + 6y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{4}{x}$$

$$q(x) = \frac{6}{x^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\begin{aligned} \tau' &= \frac{1}{c}\sqrt{q} \\ &= \frac{\sqrt{6}\sqrt{\frac{1}{x^2}}}{c} \\ \tau'' &= -\frac{\sqrt{6}}{c\sqrt{\frac{1}{x^2}}x^3} \end{aligned} \quad (6)$$

Substituting the above into (4) results in

$$\begin{aligned} p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\ &= \frac{-\frac{\sqrt{6}}{c\sqrt{\frac{1}{x^2}}x^3} - \frac{4}{x}\frac{\sqrt{6}\sqrt{\frac{1}{x^2}}}{c}}{\left(\frac{\sqrt{6}\sqrt{\frac{1}{x^2}}}{c}\right)^2} \\ &= -\frac{5c\sqrt{6}}{6} \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned} y(\tau)'' + p_1y(\tau)' + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) - \frac{5c\sqrt{6}}{6}\left(\frac{d}{d\tau}y(\tau)\right) + c^2y(\tau) &= 0 \end{aligned} \quad (7)$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = e^{\frac{5\sqrt{6}c\tau}{12}} \left(c_1 \cosh\left(\frac{\sqrt{6}c\tau}{12}\right) + ic_2 \sinh\left(\frac{\sqrt{6}c\tau}{12}\right) \right)$$

Now from (6)

$$\begin{aligned}\tau &= \int \frac{1}{c} \sqrt{q} dx \\ &= \frac{\int \sqrt{6} \sqrt{\frac{1}{x^2}} dx}{c} \\ &= \frac{\sqrt{6} \sqrt{\frac{1}{x^2}} x \ln(x)}{c}\end{aligned}$$

Substituting the above into the solution obtained gives

$$y = x^{\frac{5}{2}} \left(c_1 \cosh \left(\frac{\ln(x)}{2} \right) + i c_2 \sinh \left(\frac{\ln(x)}{2} \right) \right)$$

Now the particular solution to this ODE is found

$$x^2 y'' - 4y'x + 6y = 2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (x^5)^{\frac{2}{5}}$$

$$y_2 = (x^5)^{\frac{3}{5}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x^5)^{\frac{2}{5}} & (x^5)^{\frac{3}{5}} \\ \frac{d}{dx} \left((x^5)^{\frac{2}{5}} \right) & \frac{d}{dx} \left((x^5)^{\frac{3}{5}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x^5)^{\frac{2}{5}} & (x^5)^{\frac{3}{5}} \\ \frac{2x^4}{(x^5)^{\frac{3}{5}}} & \frac{3x^4}{(x^5)^{\frac{2}{5}}} \end{vmatrix}$$

Therefore

$$W = \left((x^5)^{\frac{2}{5}} \right) \left(\frac{3x^4}{(x^5)^{\frac{2}{5}}} \right) - \left((x^5)^{\frac{3}{5}} \right) \left(\frac{2x^4}{(x^5)^{\frac{3}{5}}} \right)$$

Which simplifies to

$$W = x^4$$

Which simplifies to

$$W = x^4$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{2(x^5)^{\frac{3}{5}}}{x^6} dx$$

Which simplifies to

$$u_1 = - \int \frac{2(x^5)^{\frac{3}{5}}}{x^6} dx$$

Hence

$$u_1 = \frac{(x^5)^{\frac{3}{5}}}{x^5}$$

And Eq. (3) becomes

$$u_2 = \int \frac{2(x^5)^{\frac{2}{5}}}{x^6} dx$$

Which simplifies to

$$u_2 = \int \frac{2(x^5)^{\frac{2}{5}}}{x^6} dx$$

Hence

$$u_2 = -\frac{2(x^5)^{\frac{2}{5}}}{3x^5}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{1}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(x^{\frac{5}{2}} \left(c_1 \cosh \left(\frac{\ln(x)}{2} \right) + ic_2 \sinh \left(\frac{\ln(x)}{2} \right) \right) \right) + \left(\frac{1}{3} \right) \\ &= \frac{1}{3} + x^{\frac{5}{2}} \left(c_1 \cosh \left(\frac{\ln(x)}{2} \right) + ic_2 \sinh \left(\frac{\ln(x)}{2} \right) \right) \end{aligned}$$

Which simplifies to

$$y = ix^{\frac{5}{2}} \sinh \left(\frac{\ln(x)}{2} \right) c_2 + x^{\frac{5}{2}} \cosh \left(\frac{\ln(x)}{2} \right) c_1 + \frac{1}{3}$$

Summary

The solution(s) found are the following

$$y = ix^{\frac{5}{2}} \sinh \left(\frac{\ln(x)}{2} \right) c_2 + x^{\frac{5}{2}} \cosh \left(\frac{\ln(x)}{2} \right) c_1 + \frac{1}{3} \quad (1)$$

Verification of solutions

$$y = ix^{\frac{5}{2}} \sinh \left(\frac{\ln(x)}{2} \right) c_2 + x^{\frac{5}{2}} \cosh \left(\frac{\ln(x)}{2} \right) c_1 + \frac{1}{3}$$

Verified OK.

2.5.5 Solving as second order change of variable on y method 1 ode

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^2y'' - 4y'x + 6y = 0$$

In normal form the given ode is written as

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{4}{x}$$
$$q(x) = \frac{6}{x^2}$$

Calculating the Liouville ode invariant Q given by

$$\begin{aligned} Q &= q - \frac{p'}{2} - \frac{p^2}{4} \\ &= \frac{6}{x^2} - \frac{\left(-\frac{4}{x}\right)'}{2} - \frac{\left(-\frac{4}{x}\right)^2}{4} \\ &= \frac{6}{x^2} - \frac{\left(\frac{4}{x^2}\right)}{2} - \frac{\left(\frac{16}{x^2}\right)}{4} \\ &= \frac{6}{x^2} - \left(\frac{2}{x^2}\right) - \frac{4}{x^2} \\ &= 0 \end{aligned}$$

Since the Liouville ode invariant does not depend on the independent variable x then the transformation

$$y = v(x)z(x) \quad (3)$$

is used to change the original ode to a constant coefficients ode in v . In (3) the term $z(x)$ is given by

$$\begin{aligned} z(x) &= e^{-\left(\int \frac{p(x)}{2} dx\right)} \\ &= e^{-\int \frac{-4}{2x} dx} \\ &= x^2 \end{aligned} \quad (5)$$

Hence (3) becomes

$$y = v(x) x^2 \quad (4)$$

Applying this change of variable to the original ode results in

$$x^4 v''(x) = 2$$

Which is now solved for $v(x)$ Simplifying the ode gives

$$v''(x) = \frac{2}{x^4}$$

Integrating once gives

$$v'(x) = -\frac{2}{3x^3} + c_1$$

Integrating again gives

$$v(x) = \frac{1}{3x^2} + c_1 x + c_2$$

Now that $v(x)$ is known, then

$$\begin{aligned} y &= v(x) z(x) \\ &= \left(c_1 x + \frac{1}{3x^2} + c_2 \right) (z(x)) \end{aligned} \quad (7)$$

But from (5)

$$z(x) = x^2$$

Hence (7) becomes

$$y = \left(c_1 x + \frac{1}{3x^2} + c_2 \right) x^2$$

Therefore the homogeneous solution y_h is

$$y_h = \left(c_1 x + \frac{1}{3x^2} + c_2 \right) x^2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (x^5)^{\frac{2}{5}}$$

$$y_2 = (x^5)^{\frac{3}{5}}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x^5)^{\frac{2}{5}} & (x^5)^{\frac{3}{5}} \\ \frac{d}{dx} \left((x^5)^{\frac{2}{5}} \right) & \frac{d}{dx} \left((x^5)^{\frac{3}{5}} \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x^5)^{\frac{2}{5}} & (x^5)^{\frac{3}{5}} \\ \frac{2x^4}{(x^5)^{\frac{3}{5}}} & \frac{3x^4}{(x^5)^{\frac{2}{5}}} \end{vmatrix}$$

Therefore

$$W = \left((x^5)^{\frac{2}{5}} \right) \left(\frac{3x^4}{(x^5)^{\frac{2}{5}}} \right) - \left((x^5)^{\frac{3}{5}} \right) \left(\frac{2x^4}{(x^5)^{\frac{3}{5}}} \right)$$

Which simplifies to

$$W = x^4$$

Which simplifies to

$$W = x^4$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{2(x^5)^{\frac{3}{5}}}{x^6} dx$$

Which simplifies to

$$u_1 = - \int \frac{2(x^5)^{\frac{3}{5}}}{x^6} dx$$

Hence

$$u_1 = \frac{(x^5)^{\frac{3}{5}}}{x^5}$$

And Eq. (3) becomes

$$u_2 = \int \frac{2(x^5)^{\frac{2}{5}}}{x^6} dx$$

Which simplifies to

$$u_2 = \int \frac{2(x^5)^{\frac{2}{5}}}{x^6} dx$$

Hence

$$u_2 = -\frac{2(x^5)^{\frac{2}{5}}}{3x^5}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{1}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(\left(c_1 x + \frac{1}{3x^2} + c_2 \right) x^2 \right) + \left(\frac{1}{3} \right) \end{aligned}$$

Which simplifies to

$$y = c_1 x^3 + c_2 x^2 + \frac{2}{3}$$

Summary

The solution(s) found are the following

$$y = c_1x^3 + c_2x^2 + \frac{2}{3} \quad (1)$$

Verification of solutions

$$y = c_1x^3 + c_2x^2 + \frac{2}{3}$$

Warning, solution could not be verified

2.5.6 Solving as second order change of variable on y method 2 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = x^2$, $B = -4x$, $C = 6$, $f(x) = 2$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$x^2y'' - 4y'x + 6y = 0$$

In normal form the ode

$$x^2y'' - 4y'x + 6y = 0 \quad (1)$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = -\frac{4}{x}$$

$$q(x) = \frac{6}{x^2}$$

Applying change of variables on the dependent variable $y = v(x)x^n$ to (2) gives the following ode where the dependent variables is $v(x)$ and not y .

$$v''(x) + \left(\frac{2n}{x} + p\right)v'(x) + \left(\frac{n(n-1)}{x^2} + \frac{np}{x} + q\right)v(x) = 0 \quad (3)$$

Let the coefficient of $v(x)$ above be zero. Hence

$$\frac{n(n-1)}{x^2} + \frac{np}{x} + q = 0 \quad (4)$$

Substituting the earlier values found for $p(x)$ and $q(x)$ into (4) gives

$$\frac{n(n-1)}{x^2} - \frac{4n}{x^2} + \frac{6}{x^2} = 0 \quad (5)$$

Solving (5) for n gives

$$n = 3 \quad (6)$$

Substituting this value in (3) gives

$$\begin{aligned} v''(x) + \frac{2v'(x)}{x} &= 0 \\ v''(x) + \frac{2v'(x)}{x} &= 0 \end{aligned} \quad (7)$$

Using the substitution

$$u(x) = v'(x)$$

Then (7) becomes

$$u'(x) + \frac{2u(x)}{x} = 0 \quad (8)$$

The above is now solved for $u(x)$. In canonical form the ODE is

$$\begin{aligned} u' &= F(x, u) \\ &= f(x)g(u) \\ &= -\frac{2u}{x} \end{aligned}$$

Where $f(x) = -\frac{2}{x}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= -\frac{2}{x} dx \\ \int \frac{1}{u} du &= \int -\frac{2}{x} dx \\ \ln(u) &= -2 \ln(x) + c_1 \\ u &= e^{-2 \ln(x) + c_1} \\ &= \frac{c_1}{x^2} \end{aligned}$$

Now that $u(x)$ is known, then

$$\begin{aligned}v'(x) &= u(x) \\v(x) &= \int u(x) dx + c_2 \\&= -\frac{c_1}{x} + c_2\end{aligned}$$

Hence

$$\begin{aligned}y &= v(x) x^n \\&= \left(-\frac{c_1}{x} + c_2\right) x^3 \\&= (c_2 x - c_1) x^2\end{aligned}$$

Now the particular solution to this ODE is found

$$x^2 y'' - 4y'x + 6y = 2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= x^2 \\y_2 &= x^3\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x^2 & x^3 \\ \frac{d}{dx}(x^2) & \frac{d}{dx}(x^3) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x^2 & x^3 \\ 2x & 3x^2 \end{vmatrix}$$

Therefore

$$W = (x^2)(3x^2) - (x^3)(2x)$$

Which simplifies to

$$W = x^4$$

Which simplifies to

$$W = x^4$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{2x^3}{x^6} dx$$

Which simplifies to

$$u_1 = - \int \frac{2}{x^3} dx$$

Hence

$$u_1 = \frac{1}{x^2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{2x^2}{x^6} dx$$

Which simplifies to

$$u_2 = \int \frac{2}{x^4} dx$$

Hence

$$u_2 = -\frac{2}{3x^3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{1}{3}$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\&= \left(\left(-\frac{c_1}{x} + c_2 \right) x^3 \right) + \left(\frac{1}{3} \right) \\&= \frac{1}{3} + \left(-\frac{c_1}{x} + c_2 \right) x^3\end{aligned}$$

Which simplifies to

$$y = \frac{1}{3} + c_2x^3 - c_1x^2$$

Summary

The solution(s) found are the following

$$y = \frac{1}{3} + c_2x^3 - c_1x^2 \quad (1)$$

Verification of solutions

$$y = \frac{1}{3} + c_2x^3 - c_1x^2$$

Verified OK.

2.5.7 Solving using Kovacic algorithm

Writing the ode as

$$x^2y'' - 4y'x + 6y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= x^2 \\B &= -4x \\C &= 6\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= 0 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = 0\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 65: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - -\infty \\
 &= \infty
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned}
 y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\
 &= z_1 e^{-\int \frac{1}{2} \frac{-4x}{x^2} dx} \\
 &= z_1 e^{2 \ln(x)} \\
 &= z_1 (x^2)
 \end{aligned}$$

Which simplifies to

$$y_1 = x^2$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{4x}{x^2} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{4\ln(x)}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1(x^2) + c_2(x^2(x)) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$x^2 y'' - 4y'x + 6y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_2 x^3 + c_1 x^2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x^2$$

$$y_2 = x^3$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x^2 & x^3 \\ \frac{d}{dx}(x^2) & \frac{d}{dx}(x^3) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x^2 & x^3 \\ 2x & 3x^2 \end{vmatrix}$$

Therefore

$$W = (x^2)(3x^2) - (x^3)(2x)$$

Which simplifies to

$$W = x^4$$

Which simplifies to

$$W = x^4$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{2x^3}{x^6} dx$$

Which simplifies to

$$u_1 = - \int \frac{2}{x^3} dx$$

Hence

$$u_1 = \frac{1}{x^2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{2x^2}{x^6} dx$$

Which simplifies to

$$u_2 = \int \frac{2}{x^4} dx$$

Hence

$$u_2 = -\frac{2}{3x^3}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{1}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2x^3 + c_1x^2) + \left(\frac{1}{3}\right) \end{aligned}$$

Which simplifies to

$$y = x^2(c_2x + c_1) + \frac{1}{3}$$

Summary

The solution(s) found are the following

$$y = x^2(c_2x + c_1) + \frac{1}{3} \quad (1)$$

Verification of solutions

$$y = x^2(c_2x + c_1) + \frac{1}{3}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    checking if the LODE is of Euler type  
    <- LODE of Euler type successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve(x^2*diff(y(x),x$2)-4*x*diff(y(x),x)+6*y(x)=2,y(x), singsol=all)
```

$$y(x) = c_2x^2 + c_1x^3 + \frac{1}{3}$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 21

```
DSolve[x^2*y''[x]-4*x*y'[x]+6*y[x]==2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2x^3 + c_1x^2 + \frac{1}{3}$$

2.6 problem Problem 6

2.6.1	Solving as second order linear constant coeff ode	583
2.6.2	Solving using Kovacic algorithm	588
2.6.3	Maple step by step solution	594

Internal problem ID [12169]

Internal file name [OUTPUT/10821_Thursday_September_21_2023_05_47_27_AM_71251468/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 6.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \cosh(x)$$

2.6.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \cosh(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \cosh(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \sin(x) \cosh(x) dx$$

Hence

$$u_1 = \frac{e^x \cos(x)}{4} - \frac{e^x \sin(x)}{4} + \frac{e^{-x} \cos(x)}{4} + \frac{e^{-x} \sin(x)}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos(x) \cosh(x)}{1} dx$$

Which simplifies to

$$u_2 = \int \cos(x) \cosh(x) dx$$

Hence

$$u_2 = \frac{e^x \cos(x)}{4} + \frac{e^x \sin(x)}{4} - \frac{e^{-x} \cos(x)}{4} + \frac{e^{-x} \sin(x)}{4}$$

Which simplifies to

$$u_1 = \frac{(\cos(x) + \sin(x)) e^{-x}}{4} + \frac{e^x(\cos(x) - \sin(x))}{4}$$
$$u_2 = \frac{(-\cos(x) + \sin(x)) e^{-x}}{4} + \frac{e^x(\cos(x) + \sin(x))}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(\frac{(\cos(x) + \sin(x)) e^{-x}}{4} + \frac{e^x(\cos(x) - \sin(x))}{4} \right) \cos(x) \\ + \left(\frac{(-\cos(x) + \sin(x)) e^{-x}}{4} + \frac{e^x(\cos(x) + \sin(x))}{4} \right) \sin(x)$$

Which simplifies to

$$y_p(x) = \frac{e^x}{4} + \frac{e^{-x}}{4}$$

Therefore the general solution is

$$y = y_h + y_p \\ = (c_1 \cos(x) + c_2 \sin(x)) + \left(\frac{e^x}{4} + \frac{e^{-x}}{4} \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{e^x}{4} + \frac{e^{-x}}{4} \quad (1)$$




Figure 105: Slope field plot

Verification of solutions

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{e^x}{4} + \frac{e^{-x}}{4}$$

Verified OK.

2.6.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$A = 1$$

$$B = 0 \tag{3}$$

$$C = 1$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$
$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = -1$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 66: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= \cos(x) \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) \cosh(x)}{1} dx$$

Which simplifies to

$$u_1 = - \int \sin(x) \cosh(x) dx$$

Hence

$$u_1 = \frac{e^x \cos(x)}{4} - \frac{e^x \sin(x)}{4} + \frac{e^{-x} \cos(x)}{4} + \frac{e^{-x} \sin(x)}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos(x) \cosh(x)}{1} dx$$

Which simplifies to

$$u_2 = \int \cos(x) \cosh(x) dx$$

Hence

$$u_2 = \frac{e^x \cos(x)}{4} + \frac{e^x \sin(x)}{4} - \frac{e^{-x} \cos(x)}{4} + \frac{e^{-x} \sin(x)}{4}$$

Which simplifies to

$$u_1 = \frac{(\cos(x) + \sin(x)) e^{-x}}{4} + \frac{e^x(\cos(x) - \sin(x))}{4}$$
$$u_2 = \frac{(-\cos(x) + \sin(x)) e^{-x}}{4} + \frac{e^x(\cos(x) + \sin(x))}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(\frac{(\cos(x) + \sin(x)) e^{-x}}{4} + \frac{e^x(\cos(x) - \sin(x))}{4} \right) \cos(x) \\ + \left(\frac{(-\cos(x) + \sin(x)) e^{-x}}{4} + \frac{e^x(\cos(x) + \sin(x))}{4} \right) \sin(x)$$

Which simplifies to

$$y_p(x) = \frac{e^x}{4} + \frac{e^{-x}}{4}$$

Therefore the general solution is

$$y = y_h + y_p \\ = (c_1 \cos(x) + c_2 \sin(x)) + \left(\frac{e^x}{4} + \frac{e^{-x}}{4} \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{e^x}{4} + \frac{e^{-x}}{4} \quad (1)$$




Figure 106: Slope field plot

Verification of solutions

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{e^x}{4} + \frac{e^{-x}}{4}$$

Verified OK.

2.6.3 Maple step by step solution

Let's solve

$$y'' + y = \cosh(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \cosh(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int \sin(x) \cosh(x) dx \right) + \sin(x) \left(\int \cos(x) \cosh(x) dx \right)$$

- Compute integrals

$$y_p(x) = \frac{e^x}{4} + \frac{e^{-x}}{4}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + \frac{e^x}{4} + \frac{e^{-x}}{4}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x), x$2)+y(x)=cosh(x), y(x), singsol=all)
```

$$y(x) = \sin(x) c_2 + c_1 \cos(x) + \frac{e^x}{4} + \frac{e^{-x}}{4}$$

✓ Solution by Mathematica

Time used: 0.063 (sec). Leaf size: 22

```
DSolve[y''[x]+y[x]==Cosh[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{\cosh(x)}{2} + c_1 \cos(x) + c_2 \sin(x)$$

2.7 problem Problem 7

2.7.1	Solving as second order ode missing x ode	597
2.7.2	Maple step by step solution	599

Internal problem ID [12170]

Internal file name [OUTPUT/10822_Thursday_September_21_2023_05_47_30_AM_81996745/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 7.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_x**"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], _Liouville, [_2nd_order, _reducible,
    _mu_x_y1], [_2nd_order, _reducible, _mu_xy]]
```

$$y'' + \frac{2y'^2}{1-y} = 0$$

2.7.1 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable.

Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$(y - 1) p(y) \left(\frac{d}{dy} p(y) \right) - 2p(y)^2 = 0$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= f(y)g(p) \\ &= \frac{2p}{y - 1} \end{aligned}$$

Where $f(y) = \frac{2}{y-1}$ and $g(p) = p$. Integrating both sides gives

$$\begin{aligned} \frac{1}{p} dp &= \frac{2}{y - 1} dy \\ \int \frac{1}{p} dp &= \int \frac{2}{y - 1} dy \\ \ln(p) &= 2 \ln(y - 1) + c_1 \\ p &= e^{2 \ln(y-1) + c_1} \\ &= c_1 (y - 1)^2 \end{aligned}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = c_1 (y - 1)^2$$

Integrating both sides gives

$$\begin{aligned} \int \frac{1}{c_1 (y - 1)^2} dy &= x + c_2 \\ -\frac{1}{c_1 (y - 1)} &= x + c_2 \end{aligned}$$

Solving for y gives these solutions

$$y_1 = \frac{c_1 c_2 + c_1 x - 1}{c_1 (x + c_2)}$$

Summary

The solution(s) found are the following

$$y = \frac{c_1 c_2 + c_1 x - 1}{c_1 (x + c_2)} \quad (1)$$

Verification of solutions

$$y = \frac{c_1 c_2 + c_1 x - 1}{c_1 (x + c_2)}$$

Verified OK.

2.7.2 Maple step by step solution

Let's solve

$$(y - 1) y'' - 2y'^2 = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Define new dependent variable u

$$u(x) = y'$$

- Compute y''

$$u'(x) = y''$$

- Use chain rule on the lhs

$$y' \left(\frac{d}{dy} u(y) \right) = y''$$

- Substitute in the definition of u

$$u(y) \left(\frac{d}{dy} u(y) \right) = y''$$

- Make substitutions $y' = u(y)$, $y'' = u(y) \left(\frac{d}{dy} u(y) \right)$ to reduce order of ODE

$$(y - 1) u(y) \left(\frac{d}{dy} u(y) \right) - 2u(y)^2 = 0$$

- Separate variables

$$\frac{\frac{d}{dy} u(y)}{u(y)} = \frac{2}{y-1}$$

- Integrate both sides with respect to y

$$\int \frac{\frac{d}{dy} u(y)}{u(y)} dy = \int \frac{2}{y-1} dy + c_1$$

- Evaluate integral

$$\ln(u(y)) = 2 \ln(y - 1) + c_1$$

- Solve for $u(y)$

$$u(y) = e^{c_1} (y - 1)^2$$

- Solve 1st ODE for $u(y)$

$$u(y) = e^{c_1} (y - 1)^2$$

- Revert to original variables with substitution $u(y) = y'$, $y = y$

$$y' = e^{c_1} (y - 1)^2$$

- Separate variables

$$\frac{y'}{(y-1)^2} = e^{c_1}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{(y-1)^2} dx = \int e^{c_1} dx + c_2$$

- Evaluate integral

$$-\frac{1}{y-1} = e^{c_1}x + c_2$$

- Solve for y

$$y = \frac{e^{c_1}x + c_2 - 1}{e^{c_1}x + c_2}$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
<- 2nd_order Liouville successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 18

```
dsolve(diff(y(x), x$2)+2/(1-y(x))*diff(y(x), x)^2=0, y(x), singsol=all)
```

$$y(x) = \frac{c_1x + c_2 - 1}{c_1x + c_2}$$

✓ Solution by Mathematica

Time used: 0.298 (sec). Leaf size: 37

```
DSolve[y''[x]+2/(1-y[x])*y'[x]^2==0, y[x], x, IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{c_1x - 1 + c_2c_1}{c_1(x + c_2)}$$

$$y(x) \rightarrow 1$$

$$y(x) \rightarrow \text{Indeterminate}$$

2.8 problem Problem 8

2.8.1	Solving as second order linear constant coeff ode	601
2.8.2	Solving as linear second order ode solved by an integrating factor ode	604
2.8.3	Solving using Kovacic algorithm	606
2.8.4	Maple step by step solution	611

Internal problem ID [12171]

Internal file name [OUTPUT/10823_Thursday_September_21_2023_05_47_30_AM_99165603/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 8.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$x'' - 4x' + 4x = e^t + e^{2t} + 1$$

2.8.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ax''(t) + Bx'(t) + Cx(t) = f(t)$$

Where $A = 1, B = -4, C = 4, f(t) = e^t + e^{2t} + 1$. Let the solution be

$$x = x_h + x_p$$

Where x_h is the solution to the homogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = 0$, and x_p is a particular solution to the non-homogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = f(t)$. x_h is the solution to

$$x'' - 4x' + 4x = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ax''(t) + Bx'(t) + Cx(t) = 0$$

Where in the above $A = 1, B = -4, C = 4$. Let the solution be $x = e^{\lambda t}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda t} - 4\lambda e^{\lambda t} + 4e^{\lambda t} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda t}$ gives

$$\lambda^2 - 4\lambda + 4 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -4, C = 4$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{4}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(-4)^2 - (4)(1)(4)} \\ &= 2 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = -2$. Therefore the solution is

$$x = c_1 e^{2t} + c_2 e^{2t} t \quad (1)$$

Therefore the homogeneous solution x_h is

$$x_h = c_1 e^{2t} + c_2 e^{2t} t$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^t + e^{2t} + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}, \{e^t\}, \{e^{2t}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{2t} t, e^{2t}\}$$

Since e^{2t} is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{1\}, \{e^t\}, \{e^{2t}t\}]$$

Since $e^{2t}t$ is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{1\}, \{e^t\}, \{e^{2t}t^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$x_p = A_1 + A_2e^t + A_3e^{2t}t^2$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution x_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_2e^t + 2A_3e^{2t} + 4A_1 = e^t + e^{2t} + 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{4}, A_2 = 1, A_3 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution x_p , gives the particular solution

$$x_p = \frac{1}{4} + e^t + \frac{e^{2t}t^2}{2}$$

Therefore the general solution is

$$\begin{aligned} x &= x_h + x_p \\ &= (c_1e^{2t} + c_2e^{2t}t) + \left(\frac{1}{4} + e^t + \frac{e^{2t}t^2}{2} \right) \end{aligned}$$

Which simplifies to

$$x = e^{2t}(c_2t + c_1) + \frac{1}{4} + e^t + \frac{e^{2t}t^2}{2}$$

Summary

The solution(s) found are the following

$$x = e^{2t}(c_2t + c_1) + \frac{1}{4} + e^t + \frac{e^{2t}t^2}{2} \quad (1)$$




Figure 107: Slope field plot

Verification of solutions

$$x = e^{2t}(c_2t + c_1) + \frac{1}{4} + e^t + \frac{e^{2t}t^2}{2}$$

Verified OK.

2.8.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$x'' + p(t)x' + \frac{(p(t))^2 + p'(t)}{2}x = f(t)$$

Where $p(t) = -4$. Therefore, there is an integrating factor given by

$$\begin{aligned}M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int -4 dx} \\ &= e^{-2t}\end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned}(M(x)x)'' &= e^{-2t}(e^t + e^{2t} + 1) \\ (e^{-2t}x)'' &= e^{-2t}(e^t + e^{2t} + 1)\end{aligned}$$

Integrating once gives

$$(e^{-2t}x)' = t - \frac{e^{-2t}}{2} - e^{-t} + c_1$$

Integrating again gives

$$(e^{-2t}x) = \frac{t^2}{2} + c_1t + \frac{e^{-2t}}{4} + e^{-t} + c_2$$

Hence the solution is

$$x = \frac{\frac{t^2}{2} + c_1t + \frac{e^{-2t}}{4} + e^{-t} + c_2}{e^{-2t}}$$

Or

$$x = c_1t e^{2t} + \frac{e^{2t}t^2}{2} + c_2e^{2t} + e^t + \frac{1}{4}$$

Summary

The solution(s) found are the following

$$x = c_1t e^{2t} + \frac{e^{2t}t^2}{2} + c_2e^{2t} + e^t + \frac{1}{4} \quad (1)$$




Figure 108: Slope field plot

Verification of solutions

$$x = c_1 t e^{2t} + \frac{e^{2t} t^2}{2} + c_2 e^{2t} + e^t + \frac{1}{4}$$

Verified OK.

2.8.3 Solving using Kovacic algorithm

Writing the ode as

$$x'' - 4x' + 4x = 0 \tag{1}$$

$$Ax'' + Bx' + Cx = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -4 \\ C &= 4 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = x e^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(t)$ then x is found using the inverse transformation

$$x = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 69: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of t , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(t) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in x is found from

$$\begin{aligned} x_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-4}{1} dt} \\ &= z_1 e^{2t} \\ &= z_1 (e^{2t}) \end{aligned}$$

Which simplifies to

$$x_1 = e^{2t}$$

The second solution x_2 to the original ode is found using reduction of order

$$x_2 = x_1 \int \frac{e^{\int -\frac{B}{A} dt}}{x_1^2} dt$$

Substituting gives

$$\begin{aligned} x_2 &= x_1 \int \frac{e^{\int -\frac{-4}{1} dt}}{(x_1)^2} dt \\ &= x_1 \int \frac{e^{4t}}{(x_1)^2} dt \\ &= x_1(t) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}x &= c_1x_1 + c_2x_2 \\ &= c_1(e^{2t}) + c_2(e^{2t}(t))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$x = x_h + x_p$$

Where x_h is the solution to the homogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = 0$, and x_p is a particular solution to the nonhomogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = f(t)$. x_h is the solution to

$$x'' - 4x' + 4x = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$x_h = c_1e^{2t} + c_2e^{2t}t$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^t + e^{2t} + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}, \{e^t\}, \{e^{2t}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{2t}t, e^{2t}\}$$

Since e^{2t} is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{1\}, \{e^t\}, \{e^{2t}t\}]$$

Since $e^{2t}t$ is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{1\}, \{e^t\}, \{e^{2t}t^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$x_p = A_1 + A_2e^t + A_3e^{2t}t^2$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution x_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_2e^t + 2A_3e^{2t} + 4A_1 = e^t + e^{2t} + 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{4}, A_2 = 1, A_3 = \frac{1}{2} \right]$$

Substituting the above back in the above trial solution x_p , gives the particular solution

$$x_p = \frac{1}{4} + e^t + \frac{e^{2t}t^2}{2}$$

Therefore the general solution is

$$\begin{aligned} x &= x_h + x_p \\ &= (c_1e^{2t} + c_2e^{2t}t) + \left(\frac{1}{4} + e^t + \frac{e^{2t}t^2}{2} \right) \end{aligned}$$

Which simplifies to

$$x = e^{2t}(c_2t + c_1) + \frac{1}{4} + e^t + \frac{e^{2t}t^2}{2}$$

Summary

The solution(s) found are the following

$$x = e^{2t}(c_2t + c_1) + \frac{1}{4} + e^t + \frac{e^{2t}t^2}{2} \quad (1)$$




Figure 109: Slope field plot

Verification of solutions

$$x = e^{2t}(c_2t + c_1) + \frac{1}{4} + e^t + \frac{e^{2t}t^2}{2}$$

Verified OK.

2.8.4 Maple step by step solution

Let's solve

$$x'' - 4x' + 4x = e^t + e^{2t} + 1$$

- Highest derivative means the order of the ODE is 2

$$x''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 4r + 4 = 0$$

- Factor the characteristic polynomial

$$(r - 2)^2 = 0$$

- Root of the characteristic polynomial

$$r = 2$$

- 1st solution of the homogeneous ODE

$$x_1(t) = e^{2t}$$

- Repeated root, multiply $x_1(t)$ by t to ensure linear independence

$$x_2(t) = e^{2t}t$$

- General solution of the ODE

$$x = c_1x_1(t) + c_2x_2(t) + x_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$x = c_1e^{2t} + c_2e^{2t}t + x_p(t)$$

- Find a particular solution $x_p(t)$ of the ODE

- Use variation of parameters to find x_p here $f(t)$ is the forcing function

$$\left[x_p(t) = -x_1(t) \left(\int \frac{x_2(t)f(t)}{W(x_1(t),x_2(t))} dt \right) + x_2(t) \left(\int \frac{x_1(t)f(t)}{W(x_1(t),x_2(t))} dt \right), f(t) = e^t + e^{2t} + 1 \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(x_1(t), x_2(t)) = \begin{bmatrix} e^{2t} & e^{2t}t \\ 2e^{2t} & 2e^{2t}t + e^{2t} \end{bmatrix}$$

- Compute Wronskian

$$W(x_1(t), x_2(t)) = e^{4t}$$

- Substitute functions into equation for $x_p(t)$

$$x_p(t) = e^{2t} \left(- \left(\int (e^t + e^{2t} + 1) t e^{-2t} dt \right) + \left(\int e^{-2t} (e^t + e^{2t} + 1) dt \right) t \right)$$

- Compute integrals

$$x_p(t) = \frac{1}{4} + e^t + \frac{e^{2t}t^2}{2}$$

- Substitute particular solution into general solution to ODE

$$x = c_2e^{2t}t + c_1e^{2t} + \frac{1}{4} + e^t + \frac{e^{2t}t^2}{2}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(x(t),t$2)-4*diff(x(t),t)+4*x(t)=exp(t)+exp(2*t)+1,x(t), singsol=all)
```

$$x(t) = \frac{(4c_1t + 2t^2 + 4c_2)e^{2t}}{4} + e^t + \frac{1}{4}$$

✓ Solution by Mathematica

Time used: 0.245 (sec). Leaf size: 32

```
DSolve[x''[t]-4*x'[t]+4*x[t]==Exp[t]+Exp[2*t]+1,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow e^{2t} \left(\frac{t^2}{2} + c_2t + c_1 \right) + e^t + \frac{1}{4}$$

2.9 problem Problem 9

2.9.1 Solving as second order ode missing y ode	614
2.9.2 Maple step by step solution	616

Internal problem ID [12172]

Internal file name [OUTPUT/10824_Thursday_September_21_2023_05_47_32_AM_35019989/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 9.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_y**"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_y], [_2nd_order, _reducible, _mu_y_y1]]
```

$$(x^2 + 1) y'' + y'^2 = -1$$

2.9.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$(x^2 + 1) p'(x) + 1 + p(x)^2 = 0$$

Which is now solve for $p(x)$ as first order ode. In canonical form the ODE is

$$\begin{aligned} p' &= F(x, p) \\ &= f(x)g(p) \\ &= \frac{-p^2 - 1}{x^2 + 1} \end{aligned}$$

Where $f(x) = \frac{1}{x^2+1}$ and $g(p) = -p^2 - 1$. Integrating both sides gives

$$\begin{aligned}\frac{1}{-p^2-1} dp &= \frac{1}{x^2+1} dx \\ \int \frac{1}{-p^2-1} dp &= \int \frac{1}{x^2+1} dx \\ -\arctan(p) &= \arctan(x) + c_1\end{aligned}$$

The solution is

$$-\arctan(p(x)) - \arctan(x) - c_1 = 0$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$-\arctan(y') - \arctan(x) - c_1 = 0$$

Integrating both sides gives

$$\begin{aligned}y &= \int -\tan(\arctan(x) + c_1) dx \\ &= \frac{ie^{4ic_1}x}{(e^{2ic_1}-1)^2} - \frac{ix}{(e^{2ic_1}-1)^2} - \frac{4e^{2ic_1} \ln((-e^{2ic_1}+1)x + ie^{2ic_1}+i)}{(e^{2ic_1}-1)^2} + c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{ie^{4ic_1}x}{(e^{2ic_1}-1)^2} - \frac{ix}{(e^{2ic_1}-1)^2} - \frac{4e^{2ic_1} \ln((-e^{2ic_1}+1)x + ie^{2ic_1}+i)}{(e^{2ic_1}-1)^2} + c_2 \quad (1)$$

Verification of solutions

$$y = \frac{ie^{4ic_1}x}{(e^{2ic_1}-1)^2} - \frac{ix}{(e^{2ic_1}-1)^2} - \frac{4e^{2ic_1} \ln((-e^{2ic_1}+1)x + ie^{2ic_1}+i)}{(e^{2ic_1}-1)^2} + c_2$$

Verified OK.

2.9.2 Maple step by step solution

Let's solve

$$(x^2 + 1) y'' + y'^2 = -1$$

- Highest derivative means the order of the ODE is 2

y''

- Make substitution $u = y'$ to reduce order of ODE

$$(x^2 + 1) u'(x) + u(x)^2 = -1$$

- Separate variables

$$\frac{u'(x)}{-u(x)^2 - 1} = \frac{1}{x^2 + 1}$$

- Integrate both sides with respect to x

$$\int \frac{u'(x)}{-u(x)^2 - 1} dx = \int \frac{1}{x^2 + 1} dx + c_1$$

- Evaluate integral

$$-\arctan(u(x)) = \arctan(x) + c_1$$

- Solve for $u(x)$

$$u(x) = -\tan(\arctan(x) + c_1)$$

- Solve 1st ODE for $u(x)$

$$u(x) = -\tan(\arctan(x) + c_1)$$

- Make substitution $u = y'$

$$y' = -\tan(\arctan(x) + c_1)$$

- Integrate both sides to solve for y

$$\int y' dx = \int -\tan(\arctan(x) + c_1) dx + c_2$$

- Compute integrals

$$y = \frac{1 e^{4 I c_1} x}{(e^{2 I c_1} - 1)^2} - \frac{I x}{(e^{2 I c_1} - 1)^2} - \frac{4 e^{2 I c_1} \ln((-e^{2 I c_1} + 1)x + I e^{2 I c_1} + 1)}{(e^{2 I c_1} - 1)^2} + c_2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying 2nd order Liouville  
trying 2nd order WeierstrassP  
trying 2nd order JacobiSN  
differential order: 2; trying a linearization to 3rd order  
trying 2nd order ODE linearizable_by_differentiation  
trying 2nd order, 2 integrating factors of the form mu(x,y)  
trying differential order: 2; missing variables  
, `-> Computing symmetries using: way = 3  
, `-> Computing symmetries using: way = exp_sym  
<- differential order: 2; canonical coordinates successful  
<- differential order 2; missing variables successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 33

```
dsolve((1+x^2)*diff(y(x),x$2)+diff(y(x),x)^2+1=0,y(x), singsol=all)
```

$$y(x) = \frac{\ln(c_1x - 1)c_1^2 + c_2c_1^2 + c_1x + \ln(c_1x - 1)}{c_1^2}$$

✓ Solution by Mathematica

Time used: 12.07 (sec). Leaf size: 33

```
DSolve[(1+x^2)*y''[x]+y'[x]^2+1==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -x \cot(c_1) + \csc^2(c_1) \log(-x \sin(c_1) - \cos(c_1)) + c_2$$

2.10 problem Problem 10

2.10.1 Solving as second order ode missing x ode	618
2.10.2 Maple step by step solution	620

Internal problem ID [12173]

Internal file name [OUTPUT/10825_Thursday_September_21_2023_05_47_32_AM_45372491/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 10.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_x**"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]
```

$$x^3 x'' = -1$$

2.10.1 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable x an independent variable. Using

$$x' = p(x)$$

Then

$$\begin{aligned} x'' &= \frac{dp}{dt} \\ &= \frac{dx}{dt} \frac{dp}{dx} \\ &= p \frac{dp}{dx} \end{aligned}$$

Hence the ode becomes

$$x^3 p(x) \left(\frac{d}{dx} p(x) \right) = -1$$

Which is now solved as first order ode for $p(x)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(x, p) \\ &= f(x)g(p) \\ &= -\frac{1}{x^3 p} \end{aligned}$$

Where $f(x) = -\frac{1}{x^3}$ and $g(p) = \frac{1}{p}$. Integrating both sides gives

$$\begin{aligned} \frac{1}{p} dp &= -\frac{1}{x^3} dx \\ \int \frac{1}{p} dp &= \int -\frac{1}{x^3} dx \\ \frac{p^2}{2} &= \frac{1}{2x^2} + c_1 \end{aligned}$$

The solution is

$$\frac{p(x)^2}{2} - \frac{1}{2x^2} - c_1 = 0$$

For solution (1) found earlier, since $p = x'$ then we now have a new first order ode to solve which is

$$\frac{x'^2}{2} - \frac{1}{2x^2} - c_1 = 0$$

Solving the given ode for x' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$x' = \frac{\sqrt{2c_1 x^2 + 1}}{x} \tag{1}$$

$$x' = -\frac{\sqrt{2c_1 x^2 + 1}}{x} \tag{2}$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned} \int \frac{x}{\sqrt{2c_1 x^2 + 1}} dx &= \int dt \\ \frac{\sqrt{2c_1 x^2 + 1}}{2c_1} &= t + c_2 \end{aligned}$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{x}{\sqrt{2c_1x^2 + 1}} dx = \int dt$$
$$-\frac{\sqrt{2c_1x^2 + 1}}{2c_1} = t + c_3$$

Summary

The solution(s) found are the following

$$\frac{\sqrt{2c_1x^2 + 1}}{2c_1} = t + c_2 \quad (1)$$

$$-\frac{\sqrt{2c_1x^2 + 1}}{2c_1} = t + c_3 \quad (2)$$

Verification of solutions

$$\frac{\sqrt{2c_1x^2 + 1}}{2c_1} = t + c_2$$

Verified OK.

$$-\frac{\sqrt{2c_1x^2 + 1}}{2c_1} = t + c_3$$

Verified OK.

2.10.2 Maple step by step solution

Let's solve

$$x^3x'' = -1$$

- Highest derivative means the order of the ODE is 2

$$x''$$

- Define new dependent variable u

$$u(t) = x'$$

- Compute x''

$$u'(t) = x''$$

- Use chain rule on the lhs

$$x' \left(\frac{d}{dx} u(x) \right) = x''$$

- Substitute in the definition of u

$$u(x) \left(\frac{d}{dx} u(x) \right) = x''$$

- Make substitutions $x' = u(x)$, $x'' = u(x) \left(\frac{d}{dx} u(x) \right)$ to reduce order of ODE

$$x^3 u(x) \left(\frac{d}{dx} u(x) \right) = -1$$

- Separate variables

$$u(x) \left(\frac{d}{dx} u(x) \right) = -\frac{1}{x^3}$$

- Integrate both sides with respect to x

$$\int u(x) \left(\frac{d}{dx} u(x) \right) dx = \int -\frac{1}{x^3} dx + c_1$$

- Evaluate integral

$$\frac{u(x)^2}{2} = \frac{1}{2x^2} + c_1$$

- Solve for $u(x)$

$$\left\{ u(x) = \frac{\sqrt{2c_1 x^2 + 1}}{x}, u(x) = -\frac{\sqrt{2c_1 x^2 + 1}}{x} \right\}$$

- Solve 1st ODE for $u(x)$

$$u(x) = \frac{\sqrt{2c_1 x^2 + 1}}{x}$$

- Revert to original variables with substitution $u(x) = x'$, $x = x$

$$x' = \frac{\sqrt{2c_1 x^2 + 1}}{x}$$

- Separate variables

$$\frac{xx'}{\sqrt{2c_1 x^2 + 1}} = 1$$

- Integrate both sides with respect to t

$$\int \frac{xx'}{\sqrt{2c_1 x^2 + 1}} dt = \int 1 dt + c_2$$

- Evaluate integral

$$\frac{\sqrt{2c_1 x^2 + 1}}{2c_1} = t + c_2$$

- Solve for x

$$\left\{ x = -\frac{\sqrt{2} \sqrt{c_1 (4c_1^2 c_2^2 + 8c_1^2 c_2 t + 4c_1^2 t^2 - 1)}}{2c_1}, x = \frac{\sqrt{2} \sqrt{c_1 (4c_1^2 c_2^2 + 8c_1^2 c_2 t + 4c_1^2 t^2 - 1)}}{2c_1} \right\}$$

- Solve 2nd ODE for $u(x)$

$$u(x) = -\frac{\sqrt{2c_1 x^2 + 1}}{x}$$

- Revert to original variables with substitution $u(x) = x', x = x$

$$x' = -\frac{\sqrt{2c_1x^2+1}}{x}$$

- Separate variables

$$\frac{xx'}{\sqrt{2c_1x^2+1}} = -1$$

- Integrate both sides with respect to t

$$\int \frac{xx'}{\sqrt{2c_1x^2+1}} dt = \int (-1) dt + c_2$$

- Evaluate integral

$$\frac{\sqrt{2c_1x^2+1}}{2c_1} = -t + c_2$$

- Solve for x

$$\left\{ x = -\frac{\sqrt{2} \sqrt{c_1(4c_1^2c_2^2 - 8c_1^2c_2t + 4c_1^2t^2 - 1)}}{2c_1}, x = \frac{\sqrt{2} \sqrt{c_1(4c_1^2c_2^2 - 8c_1^2c_2t + 4c_1^2t^2 - 1)}}{2c_1} \right\}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)+1/_a^3 = 0, _b(_a), HINT = [[_a,
symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[_a, -_b]

```

✓ Solution by Maple

Time used: 0.094 (sec). Leaf size: 52

```
dsolve(x(t)^3*diff(x(t),t$2)+1=0,x(t), singsol=all)
```

$$x(t) = \frac{\sqrt{(1 + c_1 (c_2 + t)) (-1 + c_1 (c_2 + t))} c_1}{c_1}$$
$$x(t) = -\frac{\sqrt{(1 + c_1 (c_2 + t)) (-1 + c_1 (c_2 + t))} c_1}{c_1}$$

✓ Solution by Mathematica

Time used: 4.287 (sec). Leaf size: 93

```
DSolve[x[t]^3*x''[t]+1==0,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow -\frac{\sqrt{c_1^2 t^2 + 2c_2 c_1^2 t - 1 + c_2^2 c_1^2}}{\sqrt{c_1}}$$
$$x(t) \rightarrow \frac{\sqrt{c_1^2 t^2 + 2c_2 c_1^2 t - 1 + c_2^2 c_1^2}}{\sqrt{c_1}}$$
$$x(t) \rightarrow \text{Indeterminate}$$

2.11 problem Problem 11

2.11.1 Maple step by step solution 626

Internal problem ID [12174]

Internal file name [OUTPUT/10826_Thursday_September_21_2023_05_47_33_AM_71755707/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 11.

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _linear , _nonhomogeneous]]
```

$$y'''' - 16y = x^2 - e^x$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y'''' - 16y = 0$$

The characteristic equation is

$$\lambda^4 - 16 = 0$$

The roots of the above equation are

$$\lambda_1 = 2$$

$$\lambda_2 = -2$$

$$\lambda_3 = 2i$$

$$\lambda_4 = -2i$$

Therefore the homogeneous solution is

$$y_h(x) = c_1e^{-2x} + c_2e^{2x} + e^{2ix}c_3 + e^{-2ix}c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = e^{-2x}$$

$$y_2 = e^{2x}$$

$$y_3 = e^{2ix}$$

$$y_4 = e^{-2ix}$$

Now the particular solution to the given ODE is found

$$y'''' - 16y = x^2 - e^x$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x^2 - e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^x\}, \{1, x, x^2\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^{-2x}, e^{2x}, e^{-2ix}, e^{2ix}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1e^x + A_2 + A_3x + A_4x^2$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-15A_1e^x - 16A_2 - 16A_3x - 16A_4x^2 = x^2 - e^x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{15}, A_2 = 0, A_3 = 0, A_4 = -\frac{1}{16} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^x}{15} - \frac{x^2}{16}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 e^{-2x} + c_2 e^{2x} + e^{2ix} c_3 + e^{-2ix} c_4) + \left(\frac{e^x}{15} - \frac{x^2}{16} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-2x} + c_2 e^{2x} + e^{2ix} c_3 + e^{-2ix} c_4 + \frac{e^x}{15} - \frac{x^2}{16} \quad (1)$$

Verification of solutions

$$y = c_1 e^{-2x} + c_2 e^{2x} + e^{2ix} c_3 + e^{-2ix} c_4 + \frac{e^x}{15} - \frac{x^2}{16}$$

Verified OK.

2.11.1 Maple step by step solution

Let's solve

$$y'''' - 16y = x^2 - e^x$$

- Highest derivative means the order of the ODE is 4

$$y''''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Define new variable $y_4(x)$

$$y_4(x) = y'''$$

- Isolate for $y_4'(x)$ using original ODE

$$y_4'(x) = x^2 - e^x + 16y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_4(x) = y_3'(x), y_4'(x) = x^2 - e^x + 16y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \\ y_4(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 16 & 0 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x) + \begin{bmatrix} 0 \\ 0 \\ 0 \\ x^2 - e^x \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(x) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ x^2 - e^x \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 16 & 0 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A

- Eigenpairs of A

$$\left[\left[-2, \begin{bmatrix} -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} \right], \left[2, \begin{bmatrix} \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right], \left[-2I, \begin{bmatrix} -\frac{1}{8} \\ -\frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right], \left[2I, \begin{bmatrix} \frac{1}{8} \\ -\frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-2, \begin{bmatrix} -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^{-2x} \cdot \begin{bmatrix} -\frac{1}{8} \\ \frac{1}{4} \\ -\frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[2, \begin{bmatrix} \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_2 = e^{2x} \cdot \begin{bmatrix} \frac{1}{8} \\ \frac{1}{4} \\ \frac{1}{2} \\ 1 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-2I, \begin{bmatrix} -\frac{I}{8} \\ -\frac{1}{4} \\ \frac{I}{2} \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{-2Ix} \cdot \begin{bmatrix} -\frac{I}{8} \\ -\frac{1}{4} \\ \frac{I}{2} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$(\cos(2x) - I \sin(2x)) \cdot \begin{bmatrix} -\frac{I}{8} \\ -\frac{1}{4} \\ \frac{I}{2} \\ 1 \end{bmatrix}$$

- Simplify expression

$$\begin{bmatrix} -\frac{I}{8}(\cos(2x) - I \sin(2x)) \\ -\frac{\cos(2x)}{4} + \frac{I \sin(2x)}{4} \\ \frac{I}{2}(\cos(2x) - I \sin(2x)) \\ \cos(2x) - I \sin(2x) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_3(x) = \begin{bmatrix} -\frac{\sin(2x)}{8} \\ -\frac{\cos(2x)}{4} \\ \frac{\sin(2x)}{2} \\ \cos(2x) \end{bmatrix}, \vec{y}_4(x) = \begin{bmatrix} -\frac{\cos(2x)}{8} \\ \frac{\sin(2x)}{4} \\ \frac{\cos(2x)}{2} \\ -\sin(2x) \end{bmatrix}$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{y}_p(x)$

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x) + \vec{y}_p(x)$$

□ Fundamental matrix

- Let $\phi(x)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(x) = \begin{bmatrix} -\frac{e^{-2x}}{8} & \frac{e^{2x}}{8} & -\frac{\sin(2x)}{8} & -\frac{\cos(2x)}{8} \\ \frac{e^{-2x}}{4} & \frac{e^{2x}}{4} & -\frac{\cos(2x)}{4} & \frac{\sin(2x)}{4} \\ -\frac{e^{-2x}}{2} & \frac{e^{2x}}{2} & \frac{\sin(2x)}{2} & \frac{\cos(2x)}{2} \\ e^{-2x} & e^{2x} & \cos(2x) & -\sin(2x) \end{bmatrix}$$

- The fundamental matrix, $\Phi(x)$ is a normalized version of $\phi(x)$ satisfying $\Phi(0) = I$ where I is the identity matrix

$$\Phi(x) = \phi(x) \cdot \frac{1}{\phi(0)}$$

- Substitute the value of $\phi(x)$ and $\phi(0)$

$$\Phi(x) = \begin{bmatrix} -\frac{e^{-2x}}{8} & \frac{e^{2x}}{8} & -\frac{\sin(2x)}{8} & -\frac{\cos(2x)}{8} \\ \frac{e^{-2x}}{4} & \frac{e^{2x}}{4} & -\frac{\cos(2x)}{4} & \frac{\sin(2x)}{4} \\ -\frac{e^{-2x}}{2} & \frac{e^{2x}}{2} & \frac{\sin(2x)}{2} & \frac{\cos(2x)}{2} \\ e^{-2x} & e^{2x} & \cos(2x) & -\sin(2x) \end{bmatrix} \cdot \frac{1}{\begin{bmatrix} -\frac{1}{8} & \frac{1}{8} & 0 & -\frac{1}{8} \\ \frac{1}{4} & \frac{1}{4} & -\frac{1}{4} & 0 \\ -\frac{1}{2} & \frac{1}{2} & 0 & \frac{1}{2} \\ 1 & 1 & 1 & 0 \end{bmatrix}}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(x) = \begin{bmatrix} \frac{e^{-2x}}{4} + \frac{e^{2x}}{4} + \frac{\cos(2x)}{2} & -\frac{e^{-2x}}{8} + \frac{e^{2x}}{8} + \frac{\sin(2x)}{4} & \frac{e^{-2x}}{16} + \frac{e^{2x}}{16} - \frac{\cos(2x)}{8} & -\frac{e^{-2x}}{32} + \frac{e^{2x}}{32} \\ -\frac{e^{-2x}}{2} + \frac{e^{2x}}{2} - \sin(2x) & \frac{e^{-2x}}{4} + \frac{e^{2x}}{4} + \frac{\cos(2x)}{2} & -\frac{e^{-2x}}{8} + \frac{e^{2x}}{8} + \frac{\sin(2x)}{4} & \frac{e^{-2x}}{16} + \frac{e^{2x}}{16} \\ e^{-2x} + e^{2x} - 2\cos(2x) & -\frac{e^{-2x}}{2} + \frac{e^{2x}}{2} - \sin(2x) & \frac{e^{-2x}}{4} + \frac{e^{2x}}{4} + \frac{\cos(2x)}{2} & -\frac{e^{-2x}}{8} + \frac{e^{2x}}{8} \\ -2e^{-2x} + 2e^{2x} + 4\sin(2x) & e^{-2x} + e^{2x} - 2\cos(2x) & -\frac{e^{-2x}}{2} + \frac{e^{2x}}{2} - \sin(2x) & \frac{e^{-2x}}{4} + \frac{e^{2x}}{4} \end{bmatrix}$$

□ Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(x)$ and solve for $\vec{v}(x)$

$$\vec{y}_p(x) = \Phi(x) \cdot \vec{v}(x)$$

- Take the derivative of the particular solution

$$\vec{y}'_p(x) = \Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative

$$A \cdot \Phi(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- Cancel like terms

$$\Phi(x) \cdot \vec{v}'(x) = \vec{f}(x)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(x) = \frac{1}{\Phi(x)} \cdot \vec{f}(x)$$

- Integrate to solve for $\vec{v}(x)$

$$\vec{v}(x) = \int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(x)$ into the equation for the particular solution

$$\vec{y}_p(x) = \Phi(x) \cdot \left(\int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{y}_p(x) = \begin{bmatrix} -\frac{\left(\left(x^2 + \frac{13 \cos(2x)}{20} + \frac{\sin(2x)}{5}\right)e^{2x} - \frac{16e^{3x}}{15} + \frac{3e^{4x}}{8} + \frac{1}{24}\right)e^{-2x}}{16} \\ -\frac{e^{-2x}\left(\left(x + \frac{\cos(2x)}{5} - \frac{13 \sin(2x)}{20}\right)e^{2x} - \frac{8e^{3x}}{15} + \frac{3e^{4x}}{8} - \frac{1}{24}\right)}{8} \\ \frac{e^{-2x}\left((-10 + 13 \cos(2x) + 4 \sin(2x))e^{2x} + \frac{16e^{3x}}{3} - \frac{15e^{4x}}{2} - \frac{5}{6}\right)}{80} \\ \frac{(24e^{2x} \cos(2x) - 78e^{2x} \sin(2x) - 45e^{4x} + 16e^{3x} + 5)e^{-2x}}{240} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2 + c_3 \vec{y}_3(x) + c_4 \vec{y}_4(x) + \begin{bmatrix} -\frac{\left(\left(x^2 + \frac{13 \cos(2x)}{20} + \frac{\sin(2x)}{5}\right)e^{2x} - \frac{16e^{3x}}{15} + \frac{3e^{4x}}{8} + \frac{1}{24}\right)e^{-2x}}{16} \\ -\frac{e^{-2x}\left(\left(x + \frac{\cos(2x)}{5} - \frac{13 \sin(2x)}{20}\right)e^{2x} - \frac{8e^{3x}}{15} + \frac{3e^{4x}}{8} - \frac{1}{24}\right)}{8} \\ \frac{e^{-2x}\left((-10 + 13 \cos(2x) + 4 \sin(2x))e^{2x} + \frac{16e^{3x}}{3} - \frac{15e^{4x}}{2} - \frac{5}{6}\right)}{80} \\ \frac{(24e^{2x} \cos(2x) - 78e^{2x} \sin(2x) - 45e^{4x} + 16e^{3x} + 5)e^{-2x}}{240} \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = -\frac{\left(\left(2c_4 + \frac{13}{20}\right) \cos(2x) + \left(2c_3 + \frac{1}{5}\right) \sin(2x) + x^2\right)e^{2x} + \left(-2c_2 + \frac{3}{8}\right)e^{4x} + 2c_1 - \frac{16e^{3x}}{15} + \frac{1}{24}}{16}e^{-2x}$$

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 4; linear nonhomogeneous with symmetry [0,1]  
trying high order linear exact nonhomogeneous  
trying differential order: 4; missing the dependent variable  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 53

```
dsolve(diff(y(x),x$4)-16*y(x)=x^2-exp(x),y(x), singsol=all)
```

$$y(x) = \frac{\left(\left((-16c_1 + \frac{1}{4}) \cos(2x) + x^2 - 16c_4 \sin(2x) \right) e^{2x} - 16c_3 e^{4x} - 16c_2 - \frac{16e^{3x}}{15} \right) e^{-2x}}{16}$$

✓ Solution by Mathematica

Time used: 0.299 (sec). Leaf size: 50

```
DSolve[y''''[x]-16*y[x]==x^2-Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x^2}{16} + \frac{e^x}{15} + c_1 e^{2x} + c_3 e^{-2x} + c_2 \cos(2x) + c_4 \sin(2x)$$

2.12 problem Problem 12

Internal problem ID [12175]

Internal file name [OUTPUT/10827_Thursday_September_21_2023_05_47_33_AM_47536843/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 12.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_3rd_order, _missing_x], [_3rd_order, _missing_y], [
  _3rd_order, _with_linear_symmetries], [_3rd_order, _reducible
  , _mu_y2]]
```

Unable to solve or complete the solution.

Unable to parse ODE.

Maple trace

```
`Methods for third order ODEs:
  *** Sublevel 2 ***
  Methods for third order ODEs:
  Successful isolation of  $d^3y/dx^3$ : 2 solutions were found. Trying to solve each resulting
    *** Sublevel 3 ***
    Methods for third order ODEs:
    --- Trying classification methods ---
    trying 3rd order ODE linearizable_by_differentiation
    -> Calling odsolve with the ODE`,  $\text{diff}(\text{diff}(\text{diff}(\text{diff}(y(x), x), x), x), x), x) + \text{diff}(\text{diff}(y(x), x), x) = -1, y(x), \text{singsol} = \text{none}$ 
      Methods for high order ODEs:
      --- Trying classification methods ---
      trying a quadrature
      checking if the LODE has constant coefficients
      <- constant coefficients successful
      <- 3rd order ODE linearizable_by_differentiation successful
    -----
  * Tackling next ODE.
  *** Sublevel 3 ***
  Methods for third order ODEs:
  --- Trying classification methods ---
  trying 3rd order ODE linearizable_by_differentiation
  <- 3rd order ODE linearizable_by_differentiation successful
-> Calling odsolve with the ODE`,  $\text{diff}(\text{diff}(y(x), x), x) = -1, y(x), \text{singsol} = \text{none}$  *** S
  Methods for second order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  <- quadrature successful
-> Calling odsolve with the ODE`,  $\text{diff}(\text{diff}(y(x), x), x) = 1, y(x), \text{singsol} = \text{none}$  *** Su
  Methods for second order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  <- quadrature successful`
```

✓ Solution by Maple

Time used: 0.469 (sec). Leaf size: 51

```
dsolve(diff(y(x),x$3)^2+diff(y(x),x$2)^2=1,y(x), singsol=all)
```

$$y(x) = -\frac{1}{2}x^2 + c_1x + c_2$$

$$y(x) = c_2 + c_1x + \frac{1}{2}x^2$$

$$y(x) = c_1 + c_2x + \sqrt{-c_3^2 + 1} \sin(x) + c_3 \cos(x)$$

✓ Solution by Mathematica

Time used: 0.348 (sec). Leaf size: 54

```
DSolve[y'''[x]^2+y''[x]^2==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_3x - \cos(x - c_1) + c_2$$

$$y(x) \rightarrow c_3x - \cos(x + c_1) + c_2$$

$$y(x) \rightarrow \text{Interval}[\{-1, 1\}] + c_3x + c_2$$

2.13 problem Problem 13

Internal problem ID [12176]

Internal file name [OUTPUT/10828_Thursday_September_21_2023_05_47_33_AM_87019545/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 13.

ODE order: 6.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_x]]
```

$$x^{(6)} - x'''' = 1$$

This is higher order nonhomogeneous ODE. Let the solution be

$$x = x_h + x_p$$

Where x_h is the solution to the homogeneous ODE And x_p is a particular solution to the nonhomogeneous ODE. x_h is the solution to

$$x^{(6)} - x'''' = 0$$

The characteristic equation is

$$\lambda^6 - \lambda^4 = 0$$

The roots of the above equation are

$$\lambda_1 = 0$$

$$\lambda_2 = 0$$

$$\lambda_3 = 0$$

$$\lambda_4 = 0$$

$$\lambda_5 = 1$$

$$\lambda_6 = -1$$

Therefore the homogeneous solution is

$$x_h(t) = c_1 e^{-t} + c_2 + c_3 t + t^2 c_4 + t^3 c_5 + e^t c_6$$

The fundamental set of solutions for the homogeneous solution are the following

$$x_1 = e^{-t}$$

$$x_2 = 1$$

$$x_3 = t$$

$$x_4 = t^2$$

$$x_5 = t^3$$

$$x_6 = e^t$$

Now the particular solution to the given ODE is found

$$x^{(6)} - x'''' = 1$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, t, t^2, t^3, e^t, e^{-t}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{t\}]$$

Since t is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{t^2\}]$$

Since t^2 is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{t^3\}]$$

Since t^3 is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$\{t^4\}$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$x_p = A_1 t^4$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution x_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-24A_1 = 1$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{24} \right]$$

Substituting the above back in the above trial solution x_p , gives the particular solution

$$x_p = -\frac{t^4}{24}$$

Therefore the general solution is

$$\begin{aligned} x &= x_h + x_p \\ &= (c_1 e^{-t} + c_2 + c_3 t + t^2 c_4 + t^3 c_5 + e^t c_6) + \left(-\frac{t^4}{24} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$x = c_1 e^{-t} + c_2 + c_3 t + t^2 c_4 + t^3 c_5 + e^t c_6 - \frac{t^4}{24} \quad (1)$$

Verification of solutions

$$x = c_1 e^{-t} + c_2 + c_3 t + t^2 c_4 + t^3 c_5 + e^t c_6 - \frac{t^4}{24}$$

Verified OK.

Maple trace

```
`Methods for high order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 6; linear nonhomogeneous with symmetry [0,1]
-> Calling odsolve with the ODE`, diff(diff(_b(_a), _a), _a) = _b(_a)+1, _b(_a)` *** Suble
  Methods for second order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying high order exact linear fully integrable
  trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
  trying a double symmetry of the form [xi=0, eta=F(x)]
  -> Try solving first the homogeneous part of the ODE
      checking if the LODE has constant coefficients
      <- constant coefficients successful
      <- solving first the homogeneous part of the ODE successful
  <- differential order: 6; linear nonhomogeneous with symmetry [0,1] successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 36

```
dsolve(diff(x(t),t$6)-diff(x(t),t$4)=1,x(t), singsol=all)
```

$$x(t) = -\frac{t^4}{24} + e^{-t}c_1 + c_2e^t + \frac{c_3t^3}{6} + \frac{c_4t^2}{2} + c_5t + c_6$$

✓ Solution by Mathematica

Time used: 0.103 (sec). Leaf size: 45

```
DSolve[x''''''[t]-x''''[t]==1,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow -\frac{t^4}{24} + c_6t^3 + c_5t^2 + c_4t + c_1e^t + c_2e^{-t} + c_3$$

2.14 problem Problem 14

Internal problem ID [12177]

Internal file name [OUTPUT/10829_Thursday_September_21_2023_05_47_34_AM_26296989/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 14.

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _with_linear_symmetries]]
```

$$x'''' - 2x'' + x = t^2 - 3$$

This is higher order nonhomogeneous ODE. Let the solution be

$$x = x_h + x_p$$

Where x_h is the solution to the homogeneous ODE And x_p is a particular solution to the nonhomogeneous ODE. x_h is the solution to

$$x'''' - 2x'' + x = 0$$

The characteristic equation is

$$\lambda^4 - 2\lambda^2 + 1 = 0$$

The roots of the above equation are

$$\lambda_1 = 1$$

$$\lambda_2 = 1$$

$$\lambda_3 = -1$$

$$\lambda_4 = -1$$

Therefore the homogeneous solution is

$$x_h(t) = c_1 e^{-t} + c_2 t e^{-t} + c_3 e^t + t e^t c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$x_1 = e^{-t}$$

$$x_2 = t e^{-t}$$

$$x_3 = e^t$$

$$x_4 = t e^t$$

Now the particular solution to the given ODE is found

$$x'''' - 2x'' + x = t^2 - 3$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$t^2 + 1$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[1, t, t^2]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{t e^t, t e^{-t}, e^t, e^{-t}\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$x_p = A_3 t^2 + A_2 t + A_1$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution x_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_3 t^2 + A_2 t + A_1 - 4A_3 = t^2 - 3$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 1, A_2 = 0, A_3 = 1]$$

Substituting the above back in the above trial solution x_p , gives the particular solution

$$x_p = t^2 + 1$$

Therefore the general solution is

$$\begin{aligned} x &= x_h + x_p \\ &= (c_1 e^{-t} + c_2 t e^{-t} + c_3 e^t + t e^t c_4) + (t^2 + 1) \end{aligned}$$

Which simplifies to

$$x = e^{-t}(c_2 t + c_1) + e^t(c_4 t + c_3) + t^2 + 1$$

Summary

The solution(s) found are the following

$$x = e^{-t}(c_2 t + c_1) + e^t(c_4 t + c_3) + t^2 + 1 \quad (1)$$

Verification of solutions

$$x = e^{-t}(c_2 t + c_1) + e^t(c_4 t + c_3) + t^2 + 1$$

Verified OK.

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 4; linear nonhomogeneous with symmetry [0,1]  
trying high order linear exact nonhomogeneous  
trying differential order: 4; missing the dependent variable  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 27

```
dsolve(diff(x(t),t$4)-2*diff(x(t),t$2)+x(t)=t^2-3,x(t), singsol=all)
```

$$x(t) = (c_4 t + c_2) e^{-t} + (c_3 t + c_1) e^t + t^2 + 1$$

✓ Solution by Mathematica

Time used: 0.006 (sec). Leaf size: 38

```
DSolve[x''''[t]-2*x''[t]+x[t]==t^2-3,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow t^2 + c_2 e^{-t} + c_1 e^{-t} + e^t (c_4 t + c_3) + 1$$

2.15 problem Problem 15

2.15.1 Maple step by step solution 651

Internal problem ID [12178]

Internal file name [OUTPUT/10830_Thursday_September_21_2023_05_47_34_AM_85319751/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 15.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_airy", "second_order_bessel_ode", "second order series method. Ordinary point", "second order series method. Taylor series method"

Maple gives the following as the ode type

```
[[_Emden, _Fowler]]
```

$$y'' + 4yx = 0$$

With the expansion point for the power series method at $x = 0$.

Solving ode using Taylor series method. This gives review on how the Taylor series method works for solving second order ode.

Let

$$y'' = f(x, y, y')$$

Assuming expansion is at $x_0 = 0$ (we can always shift the actual expansion point to 0 by change of variables) and assuming $f(x, y, y')$ is analytic at x_0 which must be the case for an ordinary point. Let initial conditions be $y(x_0) = y_0$ and $y'(x_0) = y'_0$. Using

Taylor series gives

$$\begin{aligned}
 y(x) &= y(x_0) + (x - x_0)y'(x_0) + \frac{(x - x_0)^2}{2}y''(x_0) + \frac{(x - x_0)^3}{3!}y'''(x_0) + \dots \\
 &= y_0 + xy'_0 + \frac{x^2}{2}f|_{x_0, y_0, y'_0} + \frac{x^3}{3!}f'|_{x_0, y_0, y'_0} + \dots \\
 &= y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} \frac{d^n f}{dx^n} \Big|_{x_0, y_0, y'_0}
 \end{aligned}$$

But

$$\frac{df}{dx} = \frac{\partial f}{\partial x} \frac{dx}{dx} + \frac{\partial f}{\partial y} \frac{dy}{dx} + \frac{\partial f}{\partial y'} \frac{dy'}{dx} \quad (1)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \quad (148)$$

$$= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \quad (149)$$

$$\begin{aligned}
 \frac{d^2 f}{dx^2} &= \frac{d}{dx} \left(\frac{df}{dx} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{df}{dx} \right) + \frac{\partial}{\partial y} \left(\frac{df}{dx} \right) y' + \frac{\partial}{\partial y'} \left(\frac{df}{dx} \right) f \quad (2)
 \end{aligned}$$

$$\begin{aligned}
 \frac{d^3 f}{dx^3} &= \frac{d}{dx} \left(\frac{d^2 f}{dx^2} \right) \\
 &= \frac{\partial}{\partial x} \left(\frac{d^2 f}{dx^2} \right) + \left(\frac{\partial}{\partial y} \frac{d^2 f}{dx^2} \right) y' + \frac{\partial}{\partial y'} \left(\frac{d^2 f}{dx^2} \right) f \quad (3)
 \end{aligned}$$

⋮

And so on. Hence if we name $F_0 = f(x, y, y')$ then the above can be written as

$$F_0 = f(x, y, y') \quad (4)$$

$$\begin{aligned} F_1 &= \frac{df}{dx} \\ &= \frac{dF_0}{dx} \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} y'' \\ &= \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} y' + \frac{\partial f}{\partial y'} f \\ &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y} y' + \frac{\partial F_0}{\partial y'} F_0 \end{aligned} \quad (5)$$

$$\begin{aligned} F_2 &= \frac{d}{dx} \left(\frac{d}{dx} f \right) \\ &= \frac{d}{dx} (F_1) \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_1 + \left(\frac{\partial F_1}{\partial y} \right) y' + \left(\frac{\partial F_1}{\partial y'} \right) F_0 \end{aligned}$$

⋮

$$\begin{aligned} F_n &= \frac{d}{dx} (F_{n-1}) \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) y'' \\ &= \frac{\partial}{\partial x} F_{n-1} + \left(\frac{\partial F_{n-1}}{\partial y} \right) y' + \left(\frac{\partial F_{n-1}}{\partial y'} \right) F_0 \end{aligned} \quad (6)$$

Therefore (6) can be used from now on along with

$$y(x) = y_0 + xy'_0 + \sum_{n=0}^{\infty} \frac{x^{n+2}}{(n+2)!} F_n|_{x_0, y_0, y'_0} \quad (7)$$

To find $y(x)$ series solution around $x = 0$. Hence

$$\begin{aligned}
 F_0 &= -4yx \\
 F_1 &= \frac{dF_0}{dx} \\
 &= \frac{\partial F_0}{\partial x} + \frac{\partial F_0}{\partial y}y' + \frac{\partial F_0}{\partial y'}F_0 \\
 &= -4y'x - 4y \\
 F_2 &= \frac{dF_1}{dx} \\
 &= \frac{\partial F_1}{\partial x} + \frac{\partial F_1}{\partial y}y' + \frac{\partial F_1}{\partial y'}F_1 \\
 &= 16x^2y - 8y' \\
 F_3 &= \frac{dF_2}{dx} \\
 &= \frac{\partial F_2}{\partial x} + \frac{\partial F_2}{\partial y}y' + \frac{\partial F_2}{\partial y'}F_2 \\
 &= 16x(y'x + 4y) \\
 F_4 &= \frac{dF_3}{dx} \\
 &= \frac{\partial F_3}{\partial x} + \frac{\partial F_3}{\partial y}y' + \frac{\partial F_3}{\partial y'}F_3 \\
 &= -64yx^3 + 96y'x + 64y
 \end{aligned}$$

And so on. Evaluating all the above at initial conditions $x = 0$ and $y(0) = y(0)$ and $y'(0) = y'(0)$ gives

$$\begin{aligned}
 F_0 &= 0 \\
 F_1 &= -4y(0) \\
 F_2 &= -8y'(0) \\
 F_3 &= 0 \\
 F_4 &= 64y(0)
 \end{aligned}$$

Substituting all the above in (7) and simplifying gives the solution as

$$y = \left(1 - \frac{2}{3}x^3 + \frac{4}{45}x^6\right)y(0) + \left(x - \frac{1}{3}x^4\right)y'(0) + O(x^6)$$

Since the expansion point $x = 0$ is an ordinary, we can also solve this using standard

power series Let the solution be represented as power series of the form

$$y = \sum_{n=0}^{\infty} a_n x^n$$

Then

$$y' = \sum_{n=1}^{\infty} n a_n x^{n-1}$$

$$y'' = \sum_{n=2}^{\infty} n(n-1) a_n x^{n-2}$$

Substituting the above back into the ode gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = -4 \left(\sum_{n=0}^{\infty} a_n x^n \right) x \quad (1)$$

Which simplifies to

$$\left(\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} \right) + \left(\sum_{n=0}^{\infty} 4x^{1+n} a_n \right) = 0 \quad (2)$$

The next step is to make all powers of x be n in each summation term. Going over each summation term above with power of x in it which is not already x^n and adjusting the power and the corresponding index gives

$$\sum_{n=2}^{\infty} n(n-1) a_n x^{n-2} = \sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n$$

$$\sum_{n=0}^{\infty} 4x^{1+n} a_n = \sum_{n=1}^{\infty} 4a_{n-1} x^n$$

Substituting all the above in Eq (2) gives the following equation where now all powers of x are the same and equal to n .

$$\left(\sum_{n=0}^{\infty} (n+2) a_{n+2} (1+n) x^n \right) + \left(\sum_{n=1}^{\infty} 4a_{n-1} x^n \right) = 0 \quad (3)$$

For $1 \leq n$, the recurrence equation is

$$(n+2) a_{n+2} (1+n) + 4a_{n-1} = 0 \quad (4)$$

Solving for a_{n+2} , gives

$$a_{n+2} = -\frac{4a_{n-1}}{(n+2)(1+n)} \quad (5)$$

For $n = 1$ the recurrence equation gives

$$6a_3 + 4a_0 = 0$$

Which after substituting the earlier terms found becomes

$$a_3 = -\frac{2a_0}{3}$$

For $n = 2$ the recurrence equation gives

$$12a_4 + 4a_1 = 0$$

Which after substituting the earlier terms found becomes

$$a_4 = -\frac{a_1}{3}$$

For $n = 3$ the recurrence equation gives

$$20a_5 + 4a_2 = 0$$

Which after substituting the earlier terms found becomes

$$a_5 = 0$$

For $n = 4$ the recurrence equation gives

$$30a_6 + 4a_3 = 0$$

Which after substituting the earlier terms found becomes

$$a_6 = \frac{4a_0}{45}$$

For $n = 5$ the recurrence equation gives

$$42a_7 + 4a_4 = 0$$

Which after substituting the earlier terms found becomes

$$a_7 = \frac{2a_1}{63}$$

And so on. Therefore the solution is

$$\begin{aligned} y &= \sum_{n=0}^{\infty} a_n x^n \\ &= a_3 x^3 + a_2 x^2 + a_1 x + a_0 + \dots \end{aligned}$$

Substituting the values for a_n found above, the solution becomes

$$y = a_0 + a_1 x - \frac{2}{3} a_0 x^3 - \frac{1}{3} a_1 x^4 + \dots$$

Collecting terms, the solution becomes

$$y = \left(1 - \frac{2x^3}{3}\right) a_0 + \left(x - \frac{1}{3}x^4\right) a_1 + O(x^6) \quad (3)$$

At $x = 0$ the solution above becomes

$$y = \left(1 - \frac{2x^3}{3}\right) c_1 + \left(x - \frac{1}{3}x^4\right) c_2 + O(x^6)$$

Summary

The solution(s) found are the following

$$y = \left(1 - \frac{2}{3}x^3 + \frac{4}{45}x^6\right) y(0) + \left(x - \frac{1}{3}x^4\right) y'(0) + O(x^6) \quad (1)$$

$$y = \left(1 - \frac{2x^3}{3}\right) c_1 + \left(x - \frac{1}{3}x^4\right) c_2 + O(x^6) \quad (2)$$

Verification of solutions

$$y = \left(1 - \frac{2}{3}x^3 + \frac{4}{45}x^6\right) y(0) + \left(x - \frac{1}{3}x^4\right) y'(0) + O(x^6)$$

Verified OK.

$$y = \left(1 - \frac{2x^3}{3}\right) c_1 + \left(x - \frac{1}{3}x^4\right) c_2 + O(x^6)$$

Verified OK.

2.15.1 Maple step by step solution

Let's solve

$$y'' = -4yx$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + 4yx = 0$$

- Assume series solution for y

$$y = \sum_{k=0}^{\infty} a_k x^k$$

- Rewrite ODE with series expansions

- Convert $x \cdot y$ to series expansion

$$x \cdot y = \sum_{k=0}^{\infty} a_k x^{k+1}$$

- Shift index using $k \rightarrow k - 1$

$$x \cdot y = \sum_{k=1}^{\infty} a_{k-1} x^k$$

- Convert y'' to series expansion

$$y'' = \sum_{k=2}^{\infty} a_k k(k-1) x^{k-2}$$

- Shift index using $k \rightarrow k + 2$

$$y'' = \sum_{k=0}^{\infty} a_{k+2} (k+2)(k+1) x^k$$

Rewrite ODE with series expansions

$$2a_2 + \left(\sum_{k=1}^{\infty} (a_{k+2} (k+2)(k+1) + 4a_{k-1}) x^k \right) = 0$$

- Each term must be 0

$$2a_2 = 0$$

- Each term in the series must be 0, giving the recursion relation

$$(k^2 + 3k + 2) a_{k+2} + 4a_{k-1} = 0$$

- Shift index using $k \rightarrow k + 1$

$$((k+1)^2 + 3k + 5) a_{k+3} + 4a_k = 0$$

- Recursion relation that defines the series solution to the ODE

$$\left[y = \sum_{k=0}^{\infty} a_k x^k, a_{k+3} = -\frac{4a_k}{k^2+5k+6}, 2a_2 = 0 \right]$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 24

```

Order:=6;
dsolve(diff(y(x),x$2)+4*x*y(x)=0,y(x),type='series',x=0);

```

$$y(x) = \left(1 - \frac{2x^3}{3}\right) y(0) + \left(x - \frac{1}{3}x^4\right) D(y)(0) + O(x^6)$$

✓ Solution by Mathematica

Time used: 0.001 (sec). Leaf size: 28

```

AsymptoticDSolveValue[y'[x]+4*x*y[x]==0,y[x],{x,0,5}]

```

$$y(x) \rightarrow c_2 \left(x - \frac{x^4}{3}\right) + c_1 \left(1 - \frac{2x^3}{3}\right)$$

2.16 problem Problem 16

2.16.1 Solving as second order bessel ode ode	653
2.16.2 Maple step by step solution	654

Internal problem ID [12179]

Internal file name [OUTPUT/10831_Thursday_September_21_2023_05_47_34_AM_92910118/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 16.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_bessel_ode**"

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$x^2y'' + y'x + \left(9x^2 - \frac{1}{25}\right)y = 0$$

2.16.1 Solving as second order bessel ode ode

Writing the ode as

$$x^2y'' + y'x + \left(9x^2 - \frac{1}{25}\right)y = 0 \tag{1}$$

Bessel ode has the form

$$x^2y'' + y'x + (-n^2 + x^2)y = 0 \tag{2}$$

The generalized form of Bessel ode is given by Bowman (1958) as the following

$$x^2y'' + (1 - 2\alpha)xy' + (\beta^2\gamma^2x^{2\gamma} - n^2\gamma^2 + \alpha^2)y = 0 \tag{3}$$

With the standard solution

$$y = x^\alpha(c_1 \text{BesselJ}(n, \beta x^\gamma) + c_2 \text{BesselY}(n, \beta x^\gamma)) \tag{4}$$

Comparing (3) to (1) and solving for α, β, n, γ gives

$$\begin{aligned}\alpha &= 0 \\ \beta &= 3 \\ n &= -\frac{1}{5} \\ \gamma &= 1\end{aligned}$$

Substituting all the above into (4) gives the solution as

$$y = c_1 \text{BesselJ}\left(-\frac{1}{5}, 3x\right) + c_2 \text{BesselY}\left(-\frac{1}{5}, 3x\right)$$

Summary

The solution(s) found are the following

$$y = c_1 \text{BesselJ}\left(-\frac{1}{5}, 3x\right) + c_2 \text{BesselY}\left(-\frac{1}{5}, 3x\right) \quad (1)$$

Verification of solutions

$$y = c_1 \text{BesselJ}\left(-\frac{1}{5}, 3x\right) + c_2 \text{BesselY}\left(-\frac{1}{5}, 3x\right)$$

Verified OK.

2.16.2 Maple step by step solution

Let's solve

$$y''x^2 + y'x + \left(9x^2 - \frac{1}{25}\right)y = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = -\frac{(225x^2-1)y}{25x^2} - \frac{y'}{x}$$

- Group terms with y on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$y'' + \frac{y'}{x} + \frac{(225x^2-1)y}{25x^2} = 0$$

- Simplify ODE

$$y''x^2 + y'x + 9x^2y - \frac{y}{25} = 0$$

- Make a change of variables

$$t = 3x$$

- Compute y'

$$y' = 3 \frac{d}{dt} y(t)$$

- Compute second derivative

$$y'' = 9 \frac{d^2}{dt^2} y(t)$$

- Apply change of variables to the ODE

$$\left(\frac{d^2}{dt^2} y(t) \right) t^2 + \left(\frac{d}{dt} y(t) \right) t + t^2 y(t) - \frac{y(t)}{25} = 0$$

- ODE is now of the Bessel form

- Solution to Bessel ODE

$$y(t) = c_1 \text{Bessel}J\left(\frac{1}{5}, t\right) + c_2 \text{Bessel}Y\left(\frac{1}{5}, t\right)$$

- Make the change from t back to x

$$y = c_1 \text{Bessel}J\left(\frac{1}{5}, 3x\right) + c_2 \text{Bessel}Y\left(\frac{1}{5}, 3x\right)$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
checking if the LODE has constant coefficients
checking if the LODE is of Euler type
trying a symmetry of the form [xi=0, eta=F(x)]
checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacic's algorithm
<- No Liouvillian solutions exist
-> Trying a solution in terms of special functions:
    -> Bessel
    <- Bessel successful
<- special function solution successful`

```


✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 19

```
dsolve(x^2*diff(y(x),x$2)+x*diff(y(x),x)+(9*x^2-1/25)*y(x)=0,y(x), singsol=all)
```

$$y(x) = c_1 \text{BesselJ}\left(\frac{1}{5}, 3x\right) + c_2 \text{BesselY}\left(\frac{1}{5}, 3x\right)$$

✓ Solution by Mathematica

Time used: 0.031 (sec). Leaf size: 26

```
DSolve[x^2*y''[x]+x*y'[x]+(9*x^2-1/25)*y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \text{BesselJ}\left(\frac{1}{5}, 3x\right) + c_2 \text{BesselY}\left(\frac{1}{5}, 3x\right)$$

2.17 problem Problem 17

2.17.1 Solving as second order ode missing y ode	657
2.17.2 Solving as second order ode missing x ode	659
2.17.3 Maple step by step solution	661

Internal problem ID [12180]

Internal file name [OUTPUT/10832_Thursday_September_21_2023_05_47_35_AM_66257595/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 17.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_x**", "**second_order_ode_missing_y**"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_xy]]
```

$$y'' + y'^2 = 1$$

With initial conditions

$$[y(0) = 0, y'(0) = 1]$$

2.17.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x) + p(x)^2 - 1 = 0$$

Which is now solve for $p(x)$ as first order ode. Integrating both sides gives

$$\int \frac{1}{-p^2 + 1} dp = x + c_1$$
$$\operatorname{arctanh}(p) = x + c_1$$

Solving for p gives these solutions

$$p_1 = \tanh(x + c_1)$$

Initial conditions are used to solve for c_1 . Substituting $x = 0$ and $p = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = \tanh(c_1)$$

Unable to solve for constant of integration. Since $\lim_{c_1 \rightarrow \infty}$ gives $p = \tanh(x + c_1) = p = 1$ and this result satisfies the given initial condition. Since $p = y'$ then the new first order ode to solve is

$$y' = 1$$

Integrating both sides gives

$$y = \int 1 dx$$
$$= x + c_2$$

Initial conditions are used to solve for c_2 . Substituting $x = 0$ and $y = 0$ in the above solution gives an equation to solve for the constant of integration.

$$0 = c_2$$

$$c_2 = 0$$

Substituting c_2 found above in the general solution gives

$$y = x$$

Initial conditions are used to solve for the constants of integration.

Summary

The solution(s) found are the following

$$y = x \tag{1}$$




Figure 110: Solution plot

Verification of solutions

$$y = x$$

Verified OK.

2.17.2 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$p(y) \left(\frac{d}{dy} p(y) \right) + p(y)^2 = 1$$

Which is now solved as first order ode for $p(y)$. Integrating both sides gives

$$\int -\frac{p}{p^2 - 1} dp = \int dy$$

$$-\frac{\ln(p - 1)}{2} - \frac{\ln(p + 1)}{2} = y + c_1$$

The above can be written as

$$\left(-\frac{1}{2}\right) (\ln(p - 1) + \ln(p + 1)) = y + c_1$$

$$\ln(p - 1) + \ln(p + 1) = (-2)(y + c_1)$$

$$= -2y - 2c_1$$

Raising both side to exponential gives

$$e^{\ln(p-1)+\ln(p+1)} = -2c_1 e^{-2y}$$

Which simplifies to

$$p^2 - 1 = c_2 e^{-2y}$$

Unable to solve for constant of integration due to RootOf in solution.

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = \text{RootOf}(_Z^2 - c_2 e^{-2y} - 1)$$

Integrating both sides gives

$$\int \frac{1}{\text{RootOf}(_Z^2 - c_2 e^{-2y} - 1)} dy = \int dx$$

$$\int^y \frac{1}{\text{RootOf}(_Z^2 - c_2 e^{-2_a} - 1)} d_a = c_3 + x$$

Unable to solve for constant of integration due to RootOf in solution.

Initial conditions are used to solve for the constants of integration.

Looking at the above solution

$$\int^y \frac{1}{\text{RootOf}(_Z^2 - c_2 e^{-2_a} - 1)} d_a = c_3 + x \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 0$ and $x = 0$ in the above gives

$$\int^0 \frac{1}{\text{RootOf}(-Z^2 - c_2 e^{-2-a} - 1)} d_a = c_3 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \text{RootOf} \left(-Z^2 - c_2 e^{-2 \text{RootOf} \left(- \left(\int^{-Z} \frac{1}{\text{RootOf}(-Z^2 - c_2 e^{-2-a} - 1)} d_a \right) + c_3 + x \right)} - 1 \right)$$

substituting $y' = 1$ and $x = 0$ in the above gives

$$1 = \lim_{x \rightarrow 0} \text{RootOf} \left(-Z^2 - c_2 e^{-2 \text{RootOf} \left(- \left(\int^{-Z} \frac{1}{\text{RootOf}(-Z^2 - c_2 e^{-2-a} - 1)} d_a \right) + c_3 + x \right)} - 1 \right) \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_2, c_3\}$. There is no solution for the constants of integrations. This solution is removed.

Verification of solutions N/A

2.17.3 Maple step by step solution

Let's solve

$$\left[y'' + y'^2 = 1, y(0) = 0, y' \Big|_{\{x=0\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2

y''

- Make substitution $u = y'$ to reduce order of ODE

$$u'(x) + u(x)^2 = 1$$

- Separate variables

$$\frac{u'(x)}{-u(x)^2 + 1} = 1$$

- Integrate both sides with respect to x

$$\int \frac{u'(x)}{-u(x)^2 + 1} dx = \int 1 dx + c_1$$

- Evaluate integral

$$\operatorname{arctanh}(u(x)) = x + c_1$$

- Solve for $u(x)$

$$u(x) = \tanh(x + c_1)$$
- Solve 1st ODE for $u(x)$

$$u(x) = \tanh(x + c_1)$$
- Make substitution $u = y'$

$$y' = \tanh(x + c_1)$$
- Integrate both sides to solve for y

$$\int y' dx = \int \tanh(x + c_1) dx + c_2$$
- Compute integrals
$$y = \ln(\cosh(x + c_1)) + c_2$$
- Check validity of solution $y = \ln(\cosh(x + c_1)) + c_2$
 - Use initial condition $y(0) = 0$

$$0 = \ln(\cosh(c_1)) + c_2$$
 - Compute derivative of the solution
$$y' = \frac{\sinh(x+c_1)}{\cosh(x+c_1)}$$
 - Use the initial condition $y' \Big|_{\{x=0\}} = 1$

$$1 = \frac{\sinh(c_1)}{\cosh(c_1)}$$
 - Solve for c_1 and c_2
 - The solution does not satisfy the initial conditions

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying 2nd order Liouville  
trying 2nd order WeierstrassP  
trying 2nd order JacobiSN  
differential order: 2; trying a linearization to 3rd order  
trying 2nd order ODE linearizable_by_differentiation  
trying 2nd order, 2 integrating factors of the form mu(x,y)  
trying a quadrature  
checking if the LODE has constant coefficients  
<- constant coefficients successful  
<- 2nd order, 2 integrating factors of the form mu(x,y) successful`
```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 5

```
dsolve([diff(y(x),x$2)+diff(y(x),x)^2=1,y(0) = 0, D(y)(0) = 1],y(x), singsol=all)
```

$$y(x) = x$$

✓ Solution by Mathematica

Time used: 0.004 (sec). Leaf size: 6

```
DSolve[{y'[x]+y'[x]^2==1,{y[0]==0,y'[0]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x$$

2.18 problem Problem 18

2.18.1 Solving as second order ode can be made integrable ode 664

2.18.2 Solving as second order ode missing x ode 667

Internal problem ID [12181]

Internal file name [OUTPUT/10833_Thursday_September_21_2023_05_47_36_AM_66756878/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 18.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_x", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x], [_2nd_order , _reducible , _mu_x_y1]]
```

$$y'' - 3\sqrt{y} = 0$$

With initial conditions

$$[y(0) = 1, y'(0) = 2]$$

2.18.1 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' - 3\sqrt{y}y' = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' - 3\sqrt{y}y') dx = 0$$
$$\frac{y'^2}{2} - 2y^{\frac{3}{2}} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{4y^{\frac{3}{2}} + 2c_1} \quad (1)$$

$$y' = -\sqrt{4y^{\frac{3}{2}} + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{4y^{\frac{3}{2}} + 2c_1}} dy = \int dx$$

$$\int^y \frac{1}{\sqrt{4a^{\frac{3}{2}} + 2c_1}} da = x + c_2$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{4y^{\frac{3}{2}} + 2c_1}} dy = \int dx$$

$$\int^y -\frac{1}{\sqrt{4a^{\frac{3}{2}} + 2c_1}} da = c_3 + x$$

Initial conditions are used to solve for the constants of integration.

Looking at the First solution

$$\int^y \frac{1}{\sqrt{4a^{\frac{3}{2}} + 2c_1}} da = x + c_2 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$\int^1 \frac{1}{\sqrt{4a^{\frac{3}{2}} + 2c_1}} da = c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \sqrt{4\text{RootOf}\left(-\left(\int^{-z} \frac{1}{\sqrt{4a^{\frac{3}{2}} + 2c_1}} da\right) + x + c_2\right)^{\frac{3}{2}} + 2c_1}$$

substituting $y' = 2$ and $x = 0$ in the above gives

$$2 = \sqrt{4\text{RootOf}\left(-\left(\int^{-z} \frac{1}{\sqrt{4a^{\frac{3}{2}} + 2c_1}} da\right) + c_2\right)^{\frac{3}{2}} + 2c_1} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. There is no solution for the constants of integrations. This solution is removed.

Looking at the Second solution

$$\int^y -\frac{1}{\sqrt{4a^{\frac{3}{2}} + 2c_1}} da = c_3 + x \quad (2)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 0$ in the above gives

$$-\left(\int^1 \frac{1}{\sqrt{4a^{\frac{3}{2}} + 2c_1}} da\right) = c_3 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\sqrt{4\text{RootOf}\left(-\left(\int^{-z} -\frac{1}{\sqrt{4a^{\frac{3}{2}} + 2c_1}} da\right) + c_3 + x\right)^{\frac{3}{2}} + 2c_1}$$

substituting $y' = 2$ and $x = 0$ in the above gives

$$2 = -\sqrt{4\text{RootOf}\left(\int^{-z} \frac{1}{\sqrt{4a^{\frac{3}{2}} + 2c_1}} da + c_3\right)^{\frac{3}{2}} + 2c_1} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_3\}$. There is no solution for the constants of integrations. This solution is removed.

Verification of solutions N/A

2.18.2 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned}y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy}\end{aligned}$$

Hence the ode becomes

$$p(y) \left(\frac{d}{dy} p(y) \right) - 3\sqrt{y} = 0$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned}p' &= F(y, p) \\ &= f(y)g(p) \\ &= \frac{3\sqrt{y}}{p}\end{aligned}$$

Where $f(y) = 3\sqrt{y}$ and $g(p) = \frac{1}{p}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{p} dp &= 3\sqrt{y} dy \\ \int \frac{1}{p} dp &= \int 3\sqrt{y} dy \\ \frac{p^2}{2} &= 2y^{\frac{3}{2}} + c_1\end{aligned}$$

The solution is

$$\frac{p(y)^2}{2} - 2y^{\frac{3}{2}} - c_1 = 0$$

Initial conditions are used to solve for c_1 . Substituting $y = 1$ and $p = 2$ in the above solution gives an equation to solve for the constant of integration.

$$-c_1 = 0$$

$$c_1 = 0$$

Substituting c_1 found above in the general solution gives

$$\frac{p^2}{2} - 2y^{\frac{3}{2}} = 0$$

Solving for $p(y)$ from the above gives

$$p(y) = 2\sqrt{y^{\frac{3}{2}}}$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = 2\sqrt{y^{\frac{3}{2}}}$$

Integrating both sides gives

$$\int \frac{1}{2\sqrt{y^{\frac{3}{2}}}} dy = \int dx$$
$$\frac{2y}{\sqrt{y^{\frac{3}{2}}}} = x + c_2$$

Initial conditions are used to solve for c_2 . Substituting $x = 0$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$2 = c_2$$

$$c_2 = 2$$

Substituting c_2 found above in the general solution gives

$$\frac{2y}{\sqrt{y^{\frac{3}{2}}}} = x + 2$$

The above simplifies to

$$-x\sqrt{y^{\frac{3}{2}}} - 2\sqrt{y^{\frac{3}{2}}} + 2y = 0$$

Initial conditions are used to solve for the constants of integration.

Summary

The solution(s) found are the following

$$(-x - 2)\sqrt{y^{\frac{3}{2}}} + 2y = 0 \quad (1)$$

Verification of solutions

$$(-x - 2)\sqrt{y^{\frac{3}{2}}} + 2y = 0$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
*** Sublevel 2 ***
Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)-3*_a^(1/2) = 0, _b(_a), HINT
symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[_a, 3/4*_b]
```

✓ Solution by Maple

Time used: 0.281 (sec). Leaf size: 11

```
dsolve([diff(y(x),x$2)=3*sqrt(y(x)),y(0) = 1, D(y)(0) = 2],y(x), singsol=all)
```

$$y(x) = \frac{(x + 2)^4}{16}$$

✓ Solution by Mathematica

Time used: 0.1 (sec). Leaf size: 14

```
DSolve[{y'[x]==3*Sqrt[y[x]],{y[0]==1,y'[0]==2}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{16}(x+2)^4$$

2.19 problem Problem 19

2.19.1 Solving as second order linear constant coeff ode	671
2.19.2 Solving using Kovacic algorithm	676
2.19.3 Maple step by step solution	681

Internal problem ID [12182]

Internal file name [OUTPUT/10834_Thursday_September_21_2023_05_47_40_AM_79672690/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 19.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = 1 - \frac{1}{\sin(x)}$$

2.19.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = -\csc(x) + 1$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\begin{aligned} \lambda_1 &= +i \\ \lambda_2 &= -i \end{aligned}$$

Which simplifies to

$$\begin{aligned} \lambda_1 &= i \\ \lambda_2 &= -i \end{aligned}$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x)(-\csc(x) + 1)}{1} dx$$

Which simplifies to

$$u_1 = - \int (-1 + \sin(x)) dx$$

Hence

$$u_1 = x + \cos(x)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos(x)(-\csc(x) + 1)}{1} dx$$

Which simplifies to

$$u_2 = \int (-\cot(x) + \cos(x)) dx$$

Hence

$$u_2 = -\ln(\sin(x)) + \sin(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = (x + \cos(x))\cos(x) + (-\ln(\sin(x)) + \sin(x))\sin(x)$$

Which simplifies to

$$y_p(x) = 1 - \sin(x)\ln(\sin(x)) + \cos(x)x$$

Therefore the general solution is

$$\begin{aligned}y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + (1 - \sin(x) \ln(\sin(x)) + \cos(x)x)\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(x) + c_2 \sin(x) + 1 - \sin(x) \ln(\sin(x)) + \cos(x)x \quad (1)$$




Figure 111: Slope field plot

Verification of solutions

$$y = c_1 \cos(x) + c_2 \sin(x) + 1 - \sin(x) \ln(\sin(x)) + \cos(x)x$$

Verified OK.

2.19.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 77: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}y_1 &= z_1 \\ &= \cos(x)\end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned}y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x))\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x)))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(x)$$

$$y_2 = \sin(x)$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ \frac{d}{dx}(\cos(x)) & \frac{d}{dx}(\sin(x)) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{vmatrix}$$

Therefore

$$W = (\cos(x))(\cos(x)) - (\sin(x))(-\sin(x))$$

Which simplifies to

$$W = \cos(x)^2 + \sin(x)^2$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\sin(x) (-\csc(x) + 1)}{1} dx$$

Which simplifies to

$$u_1 = - \int (-1 + \sin(x)) dx$$

Hence

$$u_1 = x + \cos(x)$$

And Eq. (3) becomes

$$u_2 = \int \frac{\cos(x) (-\csc(x) + 1)}{1} dx$$

Which simplifies to

$$u_2 = \int (-\cot(x) + \cos(x)) dx$$

Hence

$$u_2 = -\ln(\sin(x)) + \sin(x)$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = (x + \cos(x)) \cos(x) + (-\ln(\sin(x)) + \sin(x)) \sin(x)$$

Which simplifies to

$$y_p(x) = 1 - \sin(x) \ln(\sin(x)) + \cos(x) x$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + (1 - \sin(x) \ln(\sin(x)) + \cos(x) x) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(x) + c_2 \sin(x) + 1 - \sin(x) \ln(\sin(x)) + \cos(x)x \quad (1)$$




Figure 112: Slope field plot

Verification of solutions

$$y = c_1 \cos(x) + c_2 \sin(x) + 1 - \sin(x) \ln(\sin(x)) + \cos(x)x$$

Verified OK.

2.19.3 Maple step by step solution

Let's solve

$$y'' + y = -\csc(x) + 1$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$
- Roots of the characteristic polynomial

$$r = (-I, I)$$
- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$
- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$
- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$
- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + y_p(x)$$
- Find a particular solution $y_p(x)$ of the ODE
 - Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x), y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x), y_2(x))} dx \right), f(x) = -\csc(x) + 1 \right]$$
 - Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$
 - Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$
 - Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\cos(x) \left(\int (-1 + \sin(x)) dx \right) + \sin(x) \left(\int (-\cot(x) + \cos(x)) dx \right)$$
 - Compute integrals

$$y_p(x) = 1 - \sin(x) \ln(\sin(x)) + \cos(x)x$$
- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + 1 - \sin(x) \ln(\sin(x)) + \cos(x)x$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 23

```
dsolve(diff(y(x),x$2)+y(x)=1-1/sin(x),y(x), singsol=all)
```

$$y(x) = -\sin(x) \ln(\sin(x)) + \cos(x)(c_1 + x) + \sin(x)c_2 + 1$$

✓ Solution by Mathematica

Time used: 0.082 (sec). Leaf size: 25

```
DSolve[y''[x]+y[x]==1-1/Sin[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow (x + c_1) \cos(x) + \sin(x)(-\log(\sin(x)) + c_2) + 1$$

2.20 problem Problem 20

2.20.1 Solving as second order integrable as is ode	685
2.20.2 Solving as second order ode missing y ode	686
2.20.3 Solving as second order ode non constant coeff transformation on B ode	687
2.20.4 Solving as type second_order_integrable_as_is (not using ABC version)	689
2.20.5 Solving using Kovacic algorithm	690
2.20.6 Solving as exact linear second order ode ode	693
2.20.7 Maple step by step solution	695

Internal problem ID [12183]

Internal file name [OUTPUT/10835_Thursday_September_21_2023_05_47_42_AM_28860862/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 20.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "exact linear second order ode", "second_order_integrable_as_is", "second_order_ode_missing_y", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_y]]
```

$$u'' + \frac{2u'}{r} = 0$$

2.20.1 Solving as second order integrable as is ode

Integrating both sides of the ODE w.r.t r gives

$$\int (u''r + 2u') dr = 0$$
$$ru' + u = c_1$$

Which is now solved for u . In canonical form the ODE is

$$u' = F(r, u)$$
$$= f(r)g(u)$$
$$= \frac{-u + c_1}{r}$$

Where $f(r) = \frac{1}{r}$ and $g(u) = -u + c_1$. Integrating both sides gives

$$\frac{1}{-u + c_1} du = \frac{1}{r} dr$$
$$\int \frac{1}{-u + c_1} du = \int \frac{1}{r} dr$$
$$-\ln(-u + c_1) = \ln(r) + c_2$$

Raising both side to exponential gives

$$\frac{1}{-u + c_1} = e^{\ln(r) + c_2}$$

Which simplifies to

$$\frac{1}{-u + c_1} = c_3 r$$

Which simplifies to

$$u = \frac{(c_3 e^{c_2} r c_1 - 1) e^{-c_2}}{c_3 r}$$

Summary

The solution(s) found are the following

$$u = \frac{(c_3 e^{c_2} r c_1 - 1) e^{-c_2}}{c_3 r} \quad (1)$$

Verification of solutions

$$u = \frac{(c_3 e^{c_2} r c_1 - 1) e^{-c_2}}{c_3 r}$$

Verified OK.

2.20.2 Solving as second order ode missing y ode

This is second order ode with missing dependent variable u . Let

$$p(r) = u'$$

Then

$$p'(r) = u''$$

Hence the ode becomes

$$p'(r)r + 2p(r) = 0$$

Which is now solve for $p(r)$ as first order ode. In canonical form the ODE is

$$\begin{aligned} p' &= F(r, p) \\ &= f(r)g(p) \\ &= -\frac{2p}{r} \end{aligned}$$

Where $f(r) = -\frac{2}{r}$ and $g(p) = p$. Integrating both sides gives

$$\begin{aligned} \frac{1}{p} dp &= -\frac{2}{r} dr \\ \int \frac{1}{p} dp &= \int -\frac{2}{r} dr \\ \ln(p) &= -2 \ln(r) + c_1 \\ p &= e^{-2 \ln(r) + c_1} \\ &= \frac{c_1}{r^2} \end{aligned}$$

Since $p = u'$ then the new first order ode to solve is

$$u' = \frac{c_1}{r^2}$$

Integrating both sides gives

$$\begin{aligned} u &= \int \frac{c_1}{r^2} dr \\ &= -\frac{c_1}{r} + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$u = -\frac{c_1}{r} + c_2 \tag{1}$$

Verification of solutions

$$u = -\frac{c_1}{r} + c_2$$

Verified OK.

2.20.3 Solving as second order ode non constant coeff transformation on B ode

Given an ode of the form

$$Au'' + Bu' + Cu = F(r)$$

This method reduces the order ode the ODE by one by applying the transformation

$$u = Bv$$

This results in

$$\begin{aligned}u' &= B'v + v'B \\u'' &= B''v + B'v' + v''B + v'B' \\&= v''B + 2v' + B' + B''v\end{aligned}$$

And now the original ode becomes

$$\begin{aligned}A(v''B + 2v'B' + B''v) + B(B'v + v'B) + CBv &= 0 \\ABv'' + (2AB' + B^2)v' + (AB'' + BB' + CB)v &= 0\end{aligned}\tag{1}$$

If the term $AB'' + BB' + CB$ is zero, then this method works and can be used to solve

$$ABv'' + (2AB' + B^2)v' = 0$$

By Using $u = v'$ which reduces the order of the above ode to one. The new ode is

$$ABu' + (2AB' + B^2)u = 0$$

The above ode is first order ode which is solved for u . Now a new ode $v' = u$ is solved for v as first order ode. Then the final solution is obtain from $u = Bv$.

This method works only if the term $AB'' + BB' + CB$ is zero. The given ODE shows that

$$\begin{aligned}A &= r \\B &= 2 \\C &= 0 \\F &= 0\end{aligned}$$

The above shows that for this ode

$$\begin{aligned} AB'' + BB' + CB &= (r)(0) + (2)(0) + (0)(2) \\ &= 0 \end{aligned}$$

Hence the ode in v given in (1) now simplifies to

$$2rv'' + (4)v' = 0$$

Now by applying $v' = u$ the above becomes

$$2ru'(r) + 4u(r) = 0$$

Which is now solved for u . In canonical form the ODE is

$$\begin{aligned} u' &= F(r, u) \\ &= f(r)g(u) \\ &= -\frac{2u}{r} \end{aligned}$$

Where $f(r) = -\frac{2}{r}$ and $g(u) = u$. Integrating both sides gives

$$\begin{aligned} \frac{1}{u} du &= -\frac{2}{r} dr \\ \int \frac{1}{u} du &= \int -\frac{2}{r} dr \\ \ln(u) &= -2 \ln(r) + c_1 \\ u &= e^{-2 \ln(r) + c_1} \\ &= \frac{c_1}{r^2} \end{aligned}$$

The ode for v now becomes

$$\begin{aligned} v' &= u \\ &= \frac{c_1}{r^2} \end{aligned}$$

Which is now solved for v . Integrating both sides gives

$$\begin{aligned} v(r) &= \int \frac{c_1}{r^2} dr \\ &= -\frac{c_1}{r} + c_2 \end{aligned}$$

Therefore the solution is

$$\begin{aligned}u(r) &= Bv \\&= (2) \left(-\frac{c_1}{r} + c_2 \right) \\&= -\frac{2c_1}{r} + 2c_2\end{aligned}$$

Summary

The solution(s) found are the following

$$u = -\frac{2c_1}{r} + 2c_2 \quad (1)$$

Verification of solutions

$$u = -\frac{2c_1}{r} + 2c_2$$

Verified OK.

2.20.4 Solving as type second_order_integrable_as_is (not using ABC version)

Writing the ode as

$$u''r + 2u' = 0$$

Integrating both sides of the ODE w.r.t r gives

$$\begin{aligned}\int (u''r + 2u') dr &= 0 \\ru' + u &= c_1\end{aligned}$$

Which is now solved for u . In canonical form the ODE is

$$\begin{aligned}u' &= F(r, u) \\&= f(r)g(u) \\&= \frac{-u + c_1}{r}\end{aligned}$$

Where $f(r) = \frac{1}{r}$ and $g(u) = -u + c_1$. Integrating both sides gives

$$\begin{aligned}\frac{1}{-u + c_1} du &= \frac{1}{r} dr \\ \int \frac{1}{-u + c_1} du &= \int \frac{1}{r} dr \\ -\ln(-u + c_1) &= \ln(r) + c_2\end{aligned}$$

Raising both side to exponential gives

$$\frac{1}{-u + c_1} = e^{\ln(r)+c_2}$$

Which simplifies to

$$\frac{1}{-u + c_1} = c_3 r$$

Which simplifies to

$$u = \frac{(c_3 e^{c_2} r c_1 - 1) e^{-c_2}}{c_3 r}$$

Summary

The solution(s) found are the following

$$u = \frac{(c_3 e^{c_2} r c_1 - 1) e^{-c_2}}{c_3 r} \quad (1)$$

Verification of solutions

$$u = \frac{(c_3 e^{c_2} r c_1 - 1) e^{-c_2}}{c_3 r}$$

Verified OK.

2.20.5 Solving using Kovacic algorithm

Writing the ode as

$$u'' r + 2u' = 0 \quad (1)$$

$$Au'' + Bu' + Cu = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= r \\ B &= 2 \\ C &= 0 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(r) = u e^{\int \frac{B}{2A} dr}$$

Then (2) becomes

$$z''(r) = r z(r) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 0$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(r) = 0 \tag{7}$$

Equation (7) is now solved. After finding $z(r)$ then u is found using the inverse transformation

$$u = z(r) e^{-\int \frac{B}{2A} dr}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 79: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of r , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(r) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in u is found from

$$\begin{aligned} u_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dr} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2}{r} dr} \\ &= z_1 e^{-\ln(r)} \\ &= z_1 \left(\frac{1}{r} \right) \end{aligned}$$

Which simplifies to

$$u_1 = \frac{1}{r}$$

The second solution u_2 to the original ode is found using reduction of order

$$u_2 = u_1 \int \frac{e^{\int -\frac{B}{A} dr}}{u_1^2} dr$$

Substituting gives

$$\begin{aligned} u_2 &= u_1 \int \frac{e^{\int -\frac{2}{r} dr}}{(u_1)^2} dr \\ &= u_1 \int \frac{e^{-2 \ln(r)}}{(u_1)^2} dr \\ &= u_1(r) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}u &= c_1 u_1 + c_2 u_2 \\ &= c_1 \left(\frac{1}{r} \right) + c_2 \left(\frac{1}{r}(r) \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$u = \frac{c_1}{r} + c_2 \quad (1)$$

Verification of solutions

$$u = \frac{c_1}{r} + c_2$$

Verified OK.

2.20.6 Solving as exact linear second order ode

An ode of the form

$$p(r) u'' + q(r) u' + r(r) u = s(r)$$

is exact if

$$p''(r) - q'(r) + r(r) = 0 \quad (1)$$

For the given ode we have

$$p(x) = r$$

$$q(x) = 2$$

$$r(x) = 0$$

$$s(x) = 0$$

Hence

$$p''(x) = 0$$

$$q'(x) = 0$$

Therefore (1) becomes

$$0 - (0) + (0) = 0$$

Hence the ode is exact. Since we now know the ode is exact, it can be written as

$$(p(r) u' + (q(r) - p'(r)) u)' = s(x)$$

Integrating gives

$$p(r) u' + (q(r) - p'(r)) u = \int s(r) dr$$

Substituting the above values for p, q, r, s gives

$$ru' + u = c_1$$

We now have a first order ode to solve which is

$$ru' + u = c_1$$

In canonical form the ODE is

$$\begin{aligned} u' &= F(r, u) \\ &= f(r)g(u) \\ &= \frac{-u + c_1}{r} \end{aligned}$$

Where $f(r) = \frac{1}{r}$ and $g(u) = -u + c_1$. Integrating both sides gives

$$\begin{aligned} \frac{1}{-u + c_1} du &= \frac{1}{r} dr \\ \int \frac{1}{-u + c_1} du &= \int \frac{1}{r} dr \\ -\ln(-u + c_1) &= \ln(r) + c_2 \end{aligned}$$

Raising both side to exponential gives

$$\frac{1}{-u + c_1} = e^{\ln(r)+c_2}$$

Which simplifies to

$$\frac{1}{-u + c_1} = c_3 r$$

Which simplifies to

$$u = \frac{(c_3 e^{c_2} r c_1 - 1) e^{-c_2}}{c_3 r}$$

Summary

The solution(s) found are the following

$$u = \frac{(c_3 e^{c_2 r} c_1 - 1) e^{-c_2}}{c_3 r} \quad (1)$$

Verification of solutions

$$u = \frac{(c_3 e^{c_2 r} c_1 - 1) e^{-c_2}}{c_3 r}$$

Verified OK.

2.20.7 Maple step by step solution

Let's solve

$$u''r + 2u' = 0$$

- Highest derivative means the order of the ODE is 2

$$u''$$

- Isolate 2nd derivative

$$u'' = -\frac{2u'}{r}$$

- Group terms with u on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$u'' + \frac{2u'}{r} = 0$$

- Multiply by denominators of the ODE

$$u''r + 2u' = 0$$

- Make a change of variables

$$t = \ln(r)$$

- Substitute the change of variables back into the ODE

- Calculate the 1st derivative of u with respect to r , using the chain rule

$$u' = \left(\frac{d}{dt}u(t)\right) t'(r)$$

- Compute derivative

$$u' = \frac{\frac{d}{dt}u(t)}{r}$$

- Calculate the 2nd derivative of u with respect to r , using the chain rule

$$u'' = \left(\frac{d^2}{dt^2}u(t)\right) t'(r)^2 + t''(r) \left(\frac{d}{dt}u(t)\right)$$

- Compute derivative

$$u'' = \frac{\frac{d^2}{dt^2}u(t)}{r^2} - \frac{\frac{d}{dt}u(t)}{r^2}$$

Substitute the change of variables back into the ODE

$$\left(\frac{\frac{d^2}{dt^2}u(t)}{r^2} - \frac{\frac{d}{dt}u(t)}{r^2} \right) r + \frac{2\left(\frac{d}{dt}u(t)\right)}{r} = 0$$

- Simplify

$$\frac{\frac{d^2}{dt^2}u(t) + \frac{d}{dt}u(t)}{r} = 0$$

- Isolate 2nd derivative

$$\frac{d^2}{dt^2}u(t) = -\frac{d}{dt}u(t)$$

- Group terms with $u(t)$ on the lhs of the ODE and the rest on the rhs of the ODE; ODE is linear

$$\frac{d^2}{dt^2}u(t) + \frac{d}{dt}u(t) = 0$$

- Characteristic polynomial of ODE

$$r^2 + r = 0$$

- Factor the characteristic polynomial

$$r(r + 1) = 0$$

- Roots of the characteristic polynomial

$$r = (-1, 0)$$

- 1st solution of the ODE

$$u_1(t) = e^{-t}$$

- 2nd solution of the ODE

$$u_2(t) = 1$$

- General solution of the ODE

$$u(t) = c_1u_1(t) + c_2u_2(t)$$

- Substitute in solutions

$$u(t) = c_1e^{-t} + c_2$$

- Change variables back using $t = \ln(r)$

$$u = \frac{c_1}{r} + c_2$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
<- LODE of Euler type successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 11

```
dsolve(diff(u(r),r$2)+2/r*diff(u(r),r)=0,u(r), singsol=all)
```

$$u(r) = c_1 + \frac{c_2}{r}$$

✓ Solution by Mathematica

Time used: 0.018 (sec). Leaf size: 15

```
DSolve[u''[r]+2/r*u'[r]==0,u[r],r,IncludeSingularSolutions -> True]
```

$$u(r) \rightarrow c_2 - \frac{c_1}{r}$$

2.21 problem Problem 30

2.21.1 Solving as second order nonlinear solved by mainardi lioville
method ode 698

Internal problem ID [12184]

Internal file name [OUTPUT/10836_Thursday_September_21_2023_05_47_43_AM_53806614/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 30.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_nonlinear_solved_by_mainardi_lioville_method**"

Maple gives the following as the ode type

```
[_Liouville, [_2nd_order, _with_linear_symmetries], [_2nd_order, _reducible, _mu_x_y1], [_2nd_order, _reducible, _mu_xy]]
```

$$yy'' + y'^2 - \frac{yy'}{\sqrt{x^2 + 1}} = 0$$

2.21.1 Solving as second order nonlinear solved by mainardi lioville method ode

The ode has the Liouville form given by

$$y'' + f(x)y' + g(y)y'^2 = 0 \tag{1A}$$

Where in this problem

$$f(x) = -\frac{1}{\sqrt{x^2 + 1}}$$
$$g(y) = \frac{1}{y}$$

Dividing through by y' then Eq (1A) becomes

$$\frac{y''}{y'} + f + gy' = 0 \tag{2A}$$

But the first term in Eq (2A) can be written as

$$\frac{y''}{y'} = \frac{d}{dx} \ln(y') \quad (3A)$$

And the last term in Eq (2A) can be written as

$$\begin{aligned} g \frac{dy}{dx} &= \left(\frac{d}{dy} \int g dy \right) \frac{dy}{dx} \\ &= \frac{d}{dx} \int g dy \end{aligned} \quad (4A)$$

Substituting (3A,4A) back into (2A) gives

$$\frac{d}{dx} \ln(y') + \frac{d}{dx} \int g dy = -f \quad (5A)$$

Integrating the above w.r.t. x gives

$$\ln(y') + \int g dy = - \int f dx + c_1$$

Where c_1 is arbitrary constant. Taking the exponential of the above gives

$$y' = c_2 e^{\int -g dy} e^{\int -f dx} \quad (6A)$$

Where c_2 is a new arbitrary constant. But since $g = \frac{1}{y}$ and $f = -\frac{1}{\sqrt{x^2+1}}$, then

$$\begin{aligned} \int -g dy &= \int -\frac{1}{y} dy \\ &= -\ln(y) \\ \int -f dx &= \int \frac{1}{\sqrt{x^2+1}} dx \\ &= \operatorname{arcsinh}(x) \end{aligned}$$

Substituting the above into Eq(6A) gives

$$y' = \frac{c_2(x + \sqrt{x^2+1})}{y}$$

Which is now solved as first order separable ode. In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= \frac{c_2(x + \sqrt{x^2+1})}{y} \end{aligned}$$

Where $f(x) = c_2(x + \sqrt{x^2 + 1})$ and $g(y) = \frac{1}{y}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= c_2(x + \sqrt{x^2 + 1}) dx \\ \int \frac{1}{y} dy &= \int c_2(x + \sqrt{x^2 + 1}) dx \\ \frac{y^2}{2} &= c_2 \left(\frac{x^2}{2} + \frac{x\sqrt{x^2 + 1}}{2} + \frac{\operatorname{arcsinh}(x)}{2} \right) + c_3\end{aligned}$$

The solution is

$$\frac{y^2}{2} - c_2 \left(\frac{x^2}{2} + \frac{x\sqrt{x^2 + 1}}{2} + \frac{\operatorname{arcsinh}(x)}{2} \right) - c_3 = 0$$

Summary

The solution(s) found are the following

$$\frac{y^2}{2} - c_2 \left(\frac{x^2}{2} + \frac{x\sqrt{x^2 + 1}}{2} + \frac{\operatorname{arcsinh}(x)}{2} \right) - c_3 = 0 \quad (1)$$

Verification of solutions

$$\frac{y^2}{2} - c_2 \left(\frac{x^2}{2} + \frac{x\sqrt{x^2 + 1}}{2} + \frac{\operatorname{arcsinh}(x)}{2} \right) - c_3 = 0$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying 2nd order Liouville  
<- 2nd_order Liouville successful`
```

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 63

```
dsolve(y(x)*diff(y(x),x$2)+diff(y(x),x)^2= y(x)*diff(y(x),x)/sqrt(1+x^2),y(x), singsol=all)
```

$$y(x) = 0$$

$$y(x) = \sqrt{c_1 x \sqrt{x^2 + 1} + c_1 x^2 + c_1 \operatorname{arcsinh}(x) + 2c_2}$$

$$y(x) = -\sqrt{c_1 x \sqrt{x^2 + 1} + c_1 x^2 + c_1 \operatorname{arcsinh}(x) + 2c_2}$$

✓ Solution by Mathematica

Time used: 60.936 (sec). Leaf size: 73

```
DSolve[y[x]*y'[x]+y'[x]^2== y[x]*y'[x]/Sqrt[1+x^2],y[x],x,IncludeSingularSolutions -> True]
```

$y(x)$

$$\rightarrow c_2 \exp \left(\int_1^x \frac{1}{-K[1]c_1 + \sqrt{K[1]^2 + 1}c_1 + K[1] + \left(K[1] - \sqrt{K[1]^2 + 1} \right) \log \left(\sqrt{K[1]^2 + 1} - K[1] \right)} dK[1] \right)$$

2.22 problem Problem 31

2.22.1 Solving as second order ode missing x ode 702

Internal problem ID [12185]

Internal file name [OUTPUT/10837_Thursday_September_21_2023_05_47_44_AM_26570493/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 31.

ODE order: 2.

ODE degree: 2.

The type(s) of ODE detected by this program : "second_order_ode_missing_x"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$yy'y'' - y'^3 - y''^2 = 0$$

2.22.1 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$\left(yp(y) - p(y) \left(\frac{d}{dy} p(y) \right) \right) p(y) \left(\frac{d}{dy} p(y) \right) - p(y)^3 = 0$$

Which is now solved as first order ode for $p(y)$. Solving the given ode for $\frac{d}{dy}p(y)$ results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$\frac{d}{dy}p(y) = \frac{y}{2} + \frac{\sqrt{y^2 - 4p(y)}}{2} \quad (1)$$

$$\frac{d}{dy}p(y) = \frac{y}{2} - \frac{\sqrt{y^2 - 4p(y)}}{2} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Writing the ode as

$$\begin{aligned} \frac{d}{dy}p(y) &= \frac{y}{2} + \frac{\sqrt{y^2 - 4p}}{2} \\ \frac{d}{dy}p(y) &= \omega(y, p) \end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_y + \omega(\eta_p - \xi_y) - \omega^2 \xi_p - \omega_y \xi - \omega_p \eta = 0 \quad (A)$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = pa_3 + ya_2 + a_1 \quad (1E)$$

$$\eta = pb_3 + yb_2 + b_1 \quad (2E)$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 + \left(\frac{y}{2} + \frac{\sqrt{y^2 - 4p}}{2} \right) (b_3 - a_2) - \left(\frac{y}{2} + \frac{\sqrt{y^2 - 4p}}{2} \right)^2 a_3 \\ - \left(\frac{1}{2} + \frac{y}{2\sqrt{y^2 - 4p}} \right) (pa_3 + ya_2 + a_1) + \frac{pb_3 + yb_2 + b_1}{\sqrt{y^2 - 4p}} = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{(y^2 - 4p)^{\frac{3}{2}} a_3 + \sqrt{y^2 - 4p} y^2 a_3 + 2y^3 a_3 + 2\sqrt{y^2 - 4p} p a_3 + 4\sqrt{y^2 - 4p} y a_2 - 2\sqrt{y^2 - 4p} y b_3 - 6p y a_3 + 4\sqrt{y^2 - 4p} a_1 + 4b_2 \sqrt{y^2 - 4p} + 8p a_2 - 4p b_3 - 2y a_1 + 4y b_2 + 4b_1}{4\sqrt{y^2 - 4p}} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} & -(y^2 - 4p)^{\frac{3}{2}} a_3 - \sqrt{y^2 - 4p} y^2 a_3 - 2y^3 a_3 - 2\sqrt{y^2 - 4p} p a_3 \\ & - 4\sqrt{y^2 - 4p} y a_2 + 2\sqrt{y^2 - 4p} y b_3 + 6p y a_3 - 4y^2 a_2 + 2y^2 b_3 \\ & - 2\sqrt{y^2 - 4p} a_1 + 4b_2 \sqrt{y^2 - 4p} + 8p a_2 - 4p b_3 - 2y a_1 + 4y b_2 + 4b_1 = 0 \end{aligned} \quad (6E)$$

Simplifying the above gives

$$\begin{aligned} & -(y^2 - 4p)^{\frac{3}{2}} a_3 - 2(y^2 - 4p) y a_3 - \sqrt{y^2 - 4p} y^2 a_3 - 2(y^2 - 4p) a_2 \\ & + 2(y^2 - 4p) b_3 - 2\sqrt{y^2 - 4p} p a_3 - 4\sqrt{y^2 - 4p} y a_2 + 2\sqrt{y^2 - 4p} y b_3 - 2p y a_3 \\ & - 2y^2 a_2 - 2\sqrt{y^2 - 4p} a_1 + 4b_2 \sqrt{y^2 - 4p} + 4p b_3 - 2y a_1 + 4y b_2 + 4b_1 = 0 \end{aligned} \quad (6E)$$

Since the PDE has radicals, simplifying gives

$$\begin{aligned} & -2\sqrt{y^2 - 4p} y^2 a_3 - 2y^3 a_3 + 2\sqrt{y^2 - 4p} p a_3 + 6p y a_3 \\ & - 4\sqrt{y^2 - 4p} y a_2 + 2\sqrt{y^2 - 4p} y b_3 - 4y^2 a_2 + 2y^2 b_3 + 8p a_2 \\ & - 4p b_3 - 2\sqrt{y^2 - 4p} a_1 + 4b_2 \sqrt{y^2 - 4p} - 2y a_1 + 4y b_2 + 4b_1 = 0 \end{aligned}$$

Looking at the above PDE shows the following are all the terms with $\{p, y\}$ in them.

$$\left\{ p, y, \sqrt{y^2 - 4p} \right\}$$

The following substitution is now made to be able to collect on all terms with $\{p, y\}$ in them

$$\left\{ p = v_1, y = v_2, \sqrt{y^2 - 4p} = v_3 \right\}$$

The above PDE (6E) now becomes

$$\begin{aligned} & -2v_3^3 a_3 - 2v_3 v_2^2 a_3 - 4v_2^2 a_2 - 4v_3 v_2 a_2 + 6v_1 v_2 a_3 + 2v_3 v_1 a_3 + 2v_2^2 b_3 \\ & + 2v_3 v_2 b_3 - 2v_2 a_1 - 2v_3 a_1 + 8v_1 a_2 + 4v_2 b_2 + 4b_2 v_3 - 4v_1 b_3 + 4b_1 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3\}$$

Equation (7E) now becomes

$$6v_1v_2a_3 + 2v_3v_1a_3 + (8a_2 - 4b_3)v_1 - 2v_2^3a_3 - 2v_3v_2^2a_3 + (-4a_2 + 2b_3)v_2^2 + (-4a_2 + 2b_3)v_2v_3 + (-2a_1 + 4b_2)v_2 + (-2a_1 + 4b_2)v_3 + 4b_1 = 0 \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} -2a_3 &= 0 \\ 2a_3 &= 0 \\ 6a_3 &= 0 \\ 4b_1 &= 0 \\ -2a_1 + 4b_2 &= 0 \\ -4a_2 + 2b_3 &= 0 \\ 8a_2 - 4b_3 &= 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 2b_2 \\ a_2 &= a_2 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= b_2 \\ b_3 &= 2a_2 \end{aligned}$$

Substituting the above solution in the anstaz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= 2 \\ \eta &= y \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(y, p) \xi \\ &= y - \left(\frac{y}{2} + \frac{\sqrt{y^2 - 4p}}{2} \right) (2) \\ &= -\sqrt{y^2 - 4p} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(y, p) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dy}{\xi} = \frac{dp}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial y} + \eta \frac{\partial}{\partial p}\right) S(y, p) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{-\sqrt{y^2 - 4p}} dy \end{aligned}$$

Which results in

$$S = \frac{\sqrt{y^2 - 4p}}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_y + \omega(y, p)S_p}{R_y + \omega(y, p)R_p} \quad (2)$$

Where in the above R_y, R_p, S_y, S_p are all partial derivatives and $\omega(y, p)$ is the right hand side of the original ode given by

$$\omega(y, p) = \frac{y}{2} + \frac{\sqrt{y^2 - 4p}}{2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_y &= 1 \\ R_p &= 0 \\ S_y &= \frac{y}{2\sqrt{y^2 - 4p}} \\ S_p &= -\frac{1}{\sqrt{y^2 - 4p}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -\frac{1}{2} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for y, p in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{1}{2}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\frac{R}{2} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to y, p coordinates. This results in

$$\frac{\sqrt{y^2 - 4p(y)}}{2} = -\frac{y}{2} + c_1$$

Which simplifies to

$$\frac{\sqrt{y^2 - 4p(y)}}{2} = -\frac{y}{2} + c_1$$

Which gives

$$p(y) = -c_1^2 + c_1 y$$

Solving equation (2)

Writing the ode as

$$\begin{aligned} \frac{d}{dy}p(y) &= \frac{y}{2} - \frac{\sqrt{y^2 - 4p}}{2} \\ \frac{d}{dy}p(y) &= \omega(y, p) \end{aligned}$$

The condition of Lie symmetry is the linearized PDE given by

$$\eta_y + \omega(\eta_p - \xi_y) - \omega^2 \xi_p - \omega_y \xi - \omega_p \eta = 0 \quad (A)$$

The type of this ode is not in the lookup table. To determine ξ, η then (A) is solved using ansatz. Making bivariate polynomials of degree 1 to use as anstaz gives

$$\xi = pa_3 + ya_2 + a_1 \quad (1E)$$

$$\eta = pb_3 + yb_2 + b_1 \quad (2E)$$

Where the unknown coefficients are

$$\{a_1, a_2, a_3, b_1, b_2, b_3\}$$

Substituting equations (1E,2E) and ω into (A) gives

$$\begin{aligned} b_2 + \left(\frac{y}{2} - \frac{\sqrt{y^2 - 4p}}{2} \right) (b_3 - a_2) - \left(\frac{y}{2} - \frac{\sqrt{y^2 - 4p}}{2} \right)^2 a_3 \\ - \left(\frac{1}{2} - \frac{y}{2\sqrt{y^2 - 4p}} \right) (pa_3 + ya_2 + a_1) - \frac{pb_3 + yb_2 + b_1}{\sqrt{y^2 - 4p}} = 0 \end{aligned} \quad (5E)$$

Putting the above in normal form gives

$$\frac{(y^2 - 4p)^{\frac{3}{2}} a_3 + \sqrt{y^2 - 4p} y^2 a_3 - 2y^3 a_3 + 2\sqrt{y^2 - 4p} pa_3 + 4\sqrt{y^2 - 4p} ya_2 - 2\sqrt{y^2 - 4p} yb_3 + 6pya_3 - 4}{4\sqrt{y^2 - 4p}} = 0$$

Setting the numerator to zero gives

$$\begin{aligned} -(y^2 - 4p)^{\frac{3}{2}} a_3 - \sqrt{y^2 - 4p} y^2 a_3 + 2y^3 a_3 - 2\sqrt{y^2 - 4p} pa_3 \\ - 4\sqrt{y^2 - 4p} ya_2 + 2\sqrt{y^2 - 4p} yb_3 - 6pya_3 + 4y^2 a_2 - 2y^2 b_3 \\ - 2\sqrt{y^2 - 4p} a_1 + 4b_2 \sqrt{y^2 - 4p} - 8pa_2 + 4pb_3 + 2ya_1 - 4yb_2 - 4b_1 = 0 \end{aligned} \quad (6E)$$

Simplifying the above gives

$$\begin{aligned} -(y^2 - 4p)^{\frac{3}{2}} a_3 + 2(y^2 - 4p) ya_3 - \sqrt{y^2 - 4p} y^2 a_3 + 2(y^2 - 4p) a_2 \\ - 2(y^2 - 4p) b_3 - 2\sqrt{y^2 - 4p} pa_3 - 4\sqrt{y^2 - 4p} ya_2 + 2\sqrt{y^2 - 4p} yb_3 + 2pya_3 \\ + 2y^2 a_2 - 2\sqrt{y^2 - 4p} a_1 + 4b_2 \sqrt{y^2 - 4p} - 4pb_3 + 2ya_1 - 4yb_2 - 4b_1 = 0 \end{aligned} \quad (6E)$$

Since the PDE has radicals, simplifying gives

$$\begin{aligned} & -2\sqrt{y^2 - 4p}y^2a_3 + 2y^3a_3 + 2\sqrt{y^2 - 4p}pa_3 - 6pya_3 \\ & - 4\sqrt{y^2 - 4p}ya_2 + 2\sqrt{y^2 - 4p}yb_3 + 4y^2a_2 - 2y^2b_3 - 8pa_2 \\ & + 4pb_3 - 2\sqrt{y^2 - 4p}a_1 + 4b_2\sqrt{y^2 - 4p} + 2ya_1 - 4yb_2 - 4b_1 = 0 \end{aligned}$$

Looking at the above PDE shows the following are all the terms with $\{p, y\}$ in them.

$$\{p, y, \sqrt{y^2 - 4p}\}$$

The following substitution is now made to be able to collect on all terms with $\{p, y\}$ in them

$$\{p = v_1, y = v_2, \sqrt{y^2 - 4p} = v_3\}$$

The above PDE (6E) now becomes

$$\begin{aligned} & 2v_2^3a_3 - 2v_3v_2^2a_3 + 4v_2^2a_2 - 4v_3v_2a_2 - 6v_1v_2a_3 + 2v_3v_1a_3 - 2v_2^2b_3 \\ & + 2v_3v_2b_3 + 2v_2a_1 - 2v_3a_1 - 8v_1a_2 - 4v_2b_2 + 4b_2v_3 + 4v_1b_3 - 4b_1 = 0 \end{aligned} \quad (7E)$$

Collecting the above on the terms v_i introduced, and these are

$$\{v_1, v_2, v_3\}$$

Equation (7E) now becomes

$$\begin{aligned} & -6v_1v_2a_3 + 2v_3v_1a_3 + (-8a_2 + 4b_3)v_1 + 2v_2^3a_3 - 2v_3v_2^2a_3 + (4a_2 - 2b_3)v_2^2 \\ & + (-4a_2 + 2b_3)v_2v_3 + (2a_1 - 4b_2)v_2 + (-2a_1 + 4b_2)v_3 - 4b_1 = 0 \end{aligned} \quad (8E)$$

Setting each coefficients in (8E) to zero gives the following equations to solve

$$\begin{aligned} & -6a_3 = 0 \\ & -2a_3 = 0 \\ & 2a_3 = 0 \\ & -4b_1 = 0 \\ & -2a_1 + 4b_2 = 0 \\ & 2a_1 - 4b_2 = 0 \\ & -8a_2 + 4b_3 = 0 \\ & -4a_2 + 2b_3 = 0 \\ & 4a_2 - 2b_3 = 0 \end{aligned}$$

Solving the above equations for the unknowns gives

$$\begin{aligned} a_1 &= 2b_2 \\ a_2 &= a_2 \\ a_3 &= 0 \\ b_1 &= 0 \\ b_2 &= b_2 \\ b_3 &= 2a_2 \end{aligned}$$

Substituting the above solution in the ansatz (1E,2E) (using 1 as arbitrary value for any unknown in the RHS) gives

$$\begin{aligned} \xi &= 2 \\ \eta &= y \end{aligned}$$

Shifting is now applied to make $\xi = 0$ in order to simplify the rest of the computation

$$\begin{aligned} \eta &= \eta - \omega(y, p) \xi \\ &= y - \left(\frac{y}{2} - \frac{\sqrt{y^2 - 4p}}{2} \right) (2) \\ &= \sqrt{y^2 - 4p} \\ \xi &= 0 \end{aligned}$$

The next step is to determine the canonical coordinates R, S . The canonical coordinates map $(y, p) \rightarrow (R, S)$ where (R, S) are the canonical coordinates which make the original ode become a quadrature and hence solved by integration.

The characteristic pde which is used to find the canonical coordinates is

$$\frac{dy}{\xi} = \frac{dp}{\eta} = dS \quad (1)$$

The above comes from the requirements that $\left(\xi \frac{\partial}{\partial y} + \eta \frac{\partial}{\partial p} \right) S(y, p) = 1$. Starting with the first pair of ode's in (1) gives an ode to solve for the independent variable R in the canonical coordinates, where $S(R)$. Since $\xi = 0$ then in this special case

$$R = y$$

S is found from

$$\begin{aligned} S &= \int \frac{1}{\eta} dy \\ &= \int \frac{1}{\sqrt{y^2 - 4p}} dy \end{aligned}$$

Which results in

$$S = -\frac{\sqrt{y^2 - 4p}}{2}$$

Now that R, S are found, we need to setup the ode in these coordinates. This is done by evaluating

$$\frac{dS}{dR} = \frac{S_y + \omega(y, p)S_p}{R_y + \omega(y, p)R_p} \quad (2)$$

Where in the above R_y, R_p, S_y, S_p are all partial derivatives and $\omega(y, p)$ is the right hand side of the original ode given by

$$\omega(y, p) = \frac{y}{2} - \frac{\sqrt{y^2 - 4p}}{2}$$

Evaluating all the partial derivatives gives

$$\begin{aligned} R_y &= 1 \\ R_p &= 0 \\ S_y &= -\frac{y}{2\sqrt{y^2 - 4p}} \\ S_p &= \frac{1}{\sqrt{y^2 - 4p}} \end{aligned}$$

Substituting all the above in (2) and simplifying gives the ode in canonical coordinates.

$$\frac{dS}{dR} = -\frac{1}{2} \quad (2A)$$

We now need to express the RHS as function of R only. This is done by solving for y, p in terms of R, S from the result obtained earlier and simplifying. This gives

$$\frac{dS}{dR} = -\frac{1}{2}$$

The above is a quadrature ode. This is the whole point of Lie symmetry method. It converts an ode, no matter how complicated it is, to one that can be solved by integration when the ode is in the canonical coordinates R, S . Integrating the above gives

$$S(R) = -\frac{R}{2} + c_1 \quad (4)$$

To complete the solution, we just need to transform (4) back to y, p coordinates. This results in

$$-\frac{\sqrt{y^2 - 4p(y)}}{2} = -\frac{y}{2} + c_1$$

Which simplifies to

$$-\frac{\sqrt{y^2 - 4p(y)}}{2} = -\frac{y}{2} + c_1$$

Which gives

$$p(y) = -c_1^2 + c_1 y$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = -c_1^2 + y c_1$$

Integrating both sides gives

$$\int \frac{1}{-c_1^2 + c_1 y} dy = \int dx$$
$$\frac{\ln(-c_1^2 + c_1 y)}{c_1} = c_3 + x$$

Raising both side to exponential gives

$$e^{\frac{\ln(-c_1^2 + c_1 y)}{c_1}} = e^{c_3 + x}$$

Which simplifies to

$$(c_1(y - c_1))^{\frac{1}{c_1}} = e^x c_4$$

Summary

The solution(s) found are the following

$$y = \frac{(e^x c_4)^{c_1}}{c_1} + c_1 \tag{1}$$

Verification of solutions

$$y = \frac{(e^x c_4)^{c_1}}{c_1} + c_1$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
*** Sublevel 2 ***
Methods for second order ODEs:
Successful isolation of d^2y/dx^2: 2 solutions were found. Trying to solve each resulting
*** Sublevel 3 ***
Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)+(1/2)*(-_a+(_a^2-4*_b(_a)))
symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[a, 2*_b]
```

✓ Solution by Maple

Time used: 7.281 (sec). Leaf size: 42

```
dsolve(y(x)*diff(y(x),x)*diff(y(x),x$2)=diff(y(x),x)^3+diff(y(x),x$2)^2,y(x), singsol=all)
```

$$y(x) = -\frac{4}{-4c_1 + x}$$

$$y(x) = c_1$$

$$y(x) = e^{-c_1(c_2+x)} - c_1$$

$$y(x) = e^{c_1(c_2+x)} + c_1$$

✓ Solution by Mathematica

Time used: 13.794 (sec). Leaf size: 119

```
DSolve[y[x]*y'[x]*y''[x]==y'[x]^3+y''[x]^2,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2} \left(e^{-\frac{1}{2}(1+e^{c_1})(x+c_2)} - 1 - e^{c_1} \right)$$

$$y(x) \rightarrow \frac{1 + e^{\frac{x+c_2}{-1+\tanh\left(\frac{c_1}{2}\right)}}}{-1 + \tanh\left(\frac{c_1}{2}\right)}$$

$$y(x) \rightarrow -\frac{1}{2} - \frac{1}{2} e^{-\frac{x}{2} - \frac{c_2}{2}}$$

$$y(x) \rightarrow \frac{1}{2} \left(-1 + e^{-\frac{x}{2} - \frac{c_2}{2}} \right)$$

2.23 problem Problem 32

2.23.1 Solving as second order linear constant coeff ode	715
2.23.2 Solving using Kovacic algorithm	719
2.23.3 Maple step by step solution	724

Internal problem ID [12186]

Internal file name [OUTPUT/10838_Thursday_September_21_2023_05_47_44_AM_48181889/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 32.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$x'' + 9x = t \sin(3t)$$

2.23.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ax''(t) + Bx'(t) + Cx(t) = f(t)$$

Where $A = 1, B = 0, C = 9, f(t) = t \sin(3t)$. Let the solution be

$$x = x_h + x_p$$

Where x_h is the solution to the homogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = 0$, and x_p is a particular solution to the non-homogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = f(t)$. x_h is the solution to

$$x'' + 9x = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ax''(t) + Bx'(t) + Cx(t) = 0$$

Where in the above $A = 1, B = 0, C = 9$. Let the solution be $x = e^{\lambda t}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda t} + 9 e^{\lambda t} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda t}$ gives

$$\lambda^2 + 9 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 9$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(9)} \\ &= \pm 3i \end{aligned}$$

Hence

$$\lambda_1 = +3i$$

$$\lambda_2 = -3i$$

Which simplifies to

$$\lambda_1 = 3i$$

$$\lambda_2 = -3i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 3$. Therefore the final solution, when using Euler relation, can be written as

$$x = e^{\alpha t} (c_1 \cos(\beta t) + c_2 \sin(\beta t))$$

Which becomes

$$x = e^0 (c_1 \cos(3t) + c_2 \sin(3t))$$

Or

$$x = c_1 \cos(3t) + c_2 \sin(3t)$$

Therefore the homogeneous solution x_h is

$$x_h = c_1 \cos(3t) + c_2 \sin(3t)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$t \sin(3t)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{t \cos(3t), t \sin(3t), \cos(3t), \sin(3t)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(3t), \sin(3t)\}$$

Since $\cos(3t)$ is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{t \cos(3t), t \sin(3t), t^2 \cos(3t), t^2 \sin(3t)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$x_p = A_1 t \cos(3t) + A_2 t \sin(3t) + A_3 t^2 \cos(3t) + A_4 t^2 \sin(3t)$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution x_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} -6A_1 \sin(3t) + 6A_2 \cos(3t) + 2A_3 \cos(3t) - 12A_3 t \sin(3t) \\ + 2A_4 \sin(3t) + 12A_4 t \cos(3t) = t \sin(3t) \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = \frac{1}{36}, A_3 = -\frac{1}{12}, A_4 = 0 \right]$$

Substituting the above back in the above trial solution x_p , gives the particular solution

$$x_p = \frac{t \sin(3t)}{36} - \frac{t^2 \cos(3t)}{12}$$

Therefore the general solution is

$$\begin{aligned}x &= x_h + x_p \\ &= (c_1 \cos(3t) + c_2 \sin(3t)) + \left(\frac{t \sin(3t)}{36} - \frac{t^2 \cos(3t)}{12} \right)\end{aligned}$$

Summary

The solution(s) found are the following

$$x = c_1 \cos(3t) + c_2 \sin(3t) + \frac{t \sin(3t)}{36} - \frac{t^2 \cos(3t)}{12} \quad (1)$$




Figure 113: Slope field plot

Verification of solutions

$$x = c_1 \cos(3t) + c_2 \sin(3t) + \frac{t \sin(3t)}{36} - \frac{t^2 \cos(3t)}{12}$$

Verified OK.

2.23.2 Solving using Kovacic algorithm

Writing the ode as

$$x'' + 9x = 0 \quad (1)$$

$$Ax'' + Bx' + Cx = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 0 \\ C &= 9 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = xe^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-9}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -9 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = -9z(t) \quad (7)$$

Equation (7) is now solved. After finding $z(t)$ then x is found using the inverse transformation

$$x = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 81: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -9$ is not a function of t , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(t) = \cos(3t)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in x is found from

$$x_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}x_1 &= z_1 \\ &= \cos(3t)\end{aligned}$$

Which simplifies to

$$x_1 = \cos(3t)$$

The second solution x_2 to the original ode is found using reduction of order

$$x_2 = x_1 \int \frac{e^{\int -\frac{B}{A} dt}}{x_1^2} dt$$

Since $B = 0$ then the above becomes

$$\begin{aligned}x_2 &= x_1 \int \frac{1}{x_1^2} dt \\ &= \cos(3t) \int \frac{1}{\cos(3t)^2} dt \\ &= \cos(3t) \left(\frac{\tan(3t)}{3} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}x &= c_1 x_1 + c_2 x_2 \\ &= c_1(\cos(3t)) + c_2 \left(\cos(3t) \left(\frac{\tan(3t)}{3} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$x = x_h + x_p$$

Where x_h is the solution to the homogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = 0$, and x_p is a particular solution to the nonhomogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = f(t)$. x_h is the solution to

$$x'' + 9x = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$x_h = c_1 \cos(3t) + \frac{c_2 \sin(3t)}{3}$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$t \sin(3t)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{t \cos(3t), t \sin(3t), \cos(3t), \sin(3t)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ \frac{\sin(3t)}{3}, \cos(3t) \right\}$$

Since $\cos(3t)$ is duplicated in the UC_set, then this basis is multiplied by extra t . The UC_set becomes

$$[\{t \cos(3t), t \sin(3t), t^2 \cos(3t), t^2 \sin(3t)\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$x_p = A_1 t \cos(3t) + A_2 t \sin(3t) + A_3 t^2 \cos(3t) + A_4 t^2 \sin(3t)$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution x_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$\begin{aligned} -6A_1 \sin(3t) + 6A_2 \cos(3t) + 2A_3 \cos(3t) - 12A_3 t \sin(3t) \\ + 2A_4 \sin(3t) + 12A_4 t \cos(3t) = t \sin(3t) \end{aligned}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = \frac{1}{36}, A_3 = -\frac{1}{12}, A_4 = 0 \right]$$

Substituting the above back in the above trial solution x_p , gives the particular solution

$$x_p = \frac{t \sin(3t)}{36} - \frac{t^2 \cos(3t)}{12}$$

Therefore the general solution is

$$\begin{aligned} x &= x_h + x_p \\ &= \left(c_1 \cos(3t) + \frac{c_2 \sin(3t)}{3} \right) + \left(\frac{t \sin(3t)}{36} - \frac{t^2 \cos(3t)}{12} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$x = c_1 \cos(3t) + \frac{c_2 \sin(3t)}{3} + \frac{t \sin(3t)}{36} - \frac{t^2 \cos(3t)}{12} \quad (1)$$




Figure 114: Slope field plot

Verification of solutions

$$x = c_1 \cos(3t) + \frac{c_2 \sin(3t)}{3} + \frac{t \sin(3t)}{36} - \frac{t^2 \cos(3t)}{12}$$

Verified OK.

2.23.3 Maple step by step solution

Let's solve

$$x'' + 9x = t \sin(3t)$$

- Highest derivative means the order of the ODE is 2

$$x''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 9 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-36})}{2}$$

- Roots of the characteristic polynomial

$$r = (-3I, 3I)$$

- 1st solution of the homogeneous ODE

$$x_1(t) = \cos(3t)$$

- 2nd solution of the homogeneous ODE

$$x_2(t) = \sin(3t)$$

- General solution of the ODE

$$x = c_1 x_1(t) + c_2 x_2(t) + x_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$x = c_1 \cos(3t) + c_2 \sin(3t) + x_p(t)$$

- Find a particular solution $x_p(t)$ of the ODE

- Use variation of parameters to find x_p here $f(t)$ is the forcing function

$$\left[x_p(t) = -x_1(t) \left(\int \frac{x_2(t)f(t)}{W(x_1(t),x_2(t))} dt \right) + x_2(t) \left(\int \frac{x_1(t)f(t)}{W(x_1(t),x_2(t))} dt \right), f(t) = t \sin(3t) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(x_1(t), x_2(t)) = \begin{bmatrix} \cos(3t) & \sin(3t) \\ -3 \sin(3t) & 3 \cos(3t) \end{bmatrix}$$

- Compute Wronskian

$$W(x_1(t), x_2(t)) = 3$$

- Substitute functions into equation for $x_p(t)$

$$x_p(t) = -\frac{\cos(3t) \left(\int \sin(3t)^2 t dt \right)}{3} + \frac{\sin(3t) \left(\int \sin(6t) t dt \right)}{6}$$

- Compute integrals

$$x_p(t) = \frac{t \sin(3t)}{36} - \frac{t^2 \cos(3t)}{12}$$

- Substitute particular solution into general solution to ODE

$$x = c_1 \cos(3t) + c_2 \sin(3t) + \frac{t \sin(3t)}{36} - \frac{t^2 \cos(3t)}{12}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 31

```
dsolve(diff(x(t),t$2)+9*x(t)=t*sin(3*t),x(t), singsol=all)
```

$$x(t) = \frac{(-3t^2 + 36c_1) \cos(3t)}{36} + \frac{\sin(3t)(t + 36c_2)}{36}$$

✓ Solution by Mathematica

Time used: 0.223 (sec). Leaf size: 38

```
DSolve[x''[t]+9*x[t]==t*Sin[3*t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \left(-\frac{t^2}{12} + \frac{1}{216} + c_1 \right) \cos(3t) + \frac{1}{36}(t + 36c_2) \sin(3t)$$

2.24 problem Problem 33

2.24.1 Solving as second order linear constant coeff ode	726
2.24.2 Solving as linear second order ode solved by an integrating factor ode	731
2.24.3 Solving using Kovacic algorithm	732
2.24.4 Maple step by step solution	738

Internal problem ID [12187]

Internal file name [OUTPUT/10839_Thursday_September_21_2023_05_47_48_AM_46806992/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 33.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + 2y' + y = \sinh(x)$$

2.24.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 2, C = 1, f(x) = \sinh(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 2y' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 2, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + 2\lambda e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 2\lambda + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 2, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(2)^2 - (4)(1)(1)} \\ &= -1 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 1$. Therefore the solution is

$$y = c_1 e^{-x} + c_2 x e^{-x} \quad (1)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 e^{-x} + x e^{-x} c_2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} y_1 &= e^{-x} \\ y_2 &= x e^{-x} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-x} & x e^{-x} \\ \frac{d}{dx}(e^{-x}) & \frac{d}{dx}(x e^{-x}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-x} & x e^{-x} \\ -e^{-x} & e^{-x} - x e^{-x} \end{vmatrix}$$

Therefore

$$W = (e^{-x})(e^{-x} - x e^{-x}) - (x e^{-x})(-e^{-x})$$

Which simplifies to

$$W = e^{-2x}$$

Which simplifies to

$$W = e^{-2x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x e^{-x} \sinh(x)}{e^{-2x}} dx$$

Which simplifies to

$$u_1 = - \int \sinh(x) x e^x dx$$

Hence

$$u_1 = -\frac{\sinh(x) x \cosh(x)}{2} + \frac{x^2}{4} + \frac{\cosh(x)^2}{4} - \frac{x \cosh(x)^2}{2} + \frac{\cosh(x) \sinh(x)}{4} + \frac{x}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-x} \sinh(x)}{e^{-2x}} dx$$

Which simplifies to

$$u_2 = \int \sinh(x) e^x dx$$

Hence

$$u_2 = \frac{\cosh(x) \sinh(x)}{2} - \frac{x}{2} + \frac{\cosh(x)^2}{2}$$

Which simplifies to

$$u_1 = \frac{(-2x+1) \cosh(x)^2}{4} + \frac{(-2x+1) \sinh(x) \cosh(x)}{4} + \frac{x^2}{4} + \frac{x}{4}$$

$$u_2 = -\frac{x}{2} + \frac{\sinh(2x)}{4} + \frac{\cosh(2x)}{4} + \frac{1}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(\frac{(-2x+1) \cosh(x)^2}{4} + \frac{(-2x+1) \sinh(x) \cosh(x)}{4} + \frac{x^2}{4} + \frac{x}{4} \right) e^{-x}$$

$$+ \left(-\frac{x}{2} + \frac{\sinh(2x)}{4} + \frac{\cosh(2x)}{4} + \frac{1}{4} \right) x e^{-x}$$

Which simplifies to

$$y_p(x) = \frac{(\cosh(x) \sinh(x) + \cosh(x)^2 - x^2 + x) e^{-x}}{4}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= (c_1 e^{-x} + x e^{-x} c_2) + \left(\frac{(\cosh(x) \sinh(x) + \cosh(x)^2 - x^2 + x) e^{-x}}{4} \right)$$

Which simplifies to

$$y = e^{-x}(c_2x + c_1) + \frac{(\cosh(x) \sinh(x) + \cosh(x)^2 - x^2 + x) e^{-x}}{4}$$

Summary

The solution(s) found are the following

$$y = e^{-x}(c_2x + c_1) + \frac{(\cosh(x) \sinh(x) + \cosh(x)^2 - x^2 + x) e^{-x}}{4} \quad (1)$$




Figure 115: Slope field plot

Verification of solutions

$$y = e^{-x}(c_2x + c_1) + \frac{(\cosh(x) \sinh(x) + \cosh(x)^2 - x^2 + x) e^{-x}}{4}$$

Verified OK.

2.24.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$y'' + p(x)y' + \frac{(p(x)^2 + p'(x))y}{2} = f(x)$$

Where $p(x) = 2$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int 2 dx} \\ &= e^x \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned} (M(x)y)'' &= \sinh(x)e^x \\ (e^x y)'' &= \sinh(x)e^x \end{aligned}$$

Integrating once gives

$$(e^x y)' = -\frac{x}{2} + \frac{\sinh(2x)}{4} + \frac{\cosh(2x)}{4} + \frac{1}{4} + c_1$$

Integrating again gives

$$(e^x y) = -\frac{x^2}{4} + c_1 x + \frac{x}{4} + \frac{\sinh(2x)}{8} + \frac{\cosh(2x)}{8} + c_2$$

Hence the solution is

$$y = \frac{-\frac{x^2}{4} + c_1 x + \frac{x}{4} + \frac{\sinh(2x)}{8} + \frac{\cosh(2x)}{8} + c_2}{e^x}$$

Or

$$y = e^{-x} c_1 x - \frac{x^2 e^{-x}}{4} + \frac{\cosh(x)^2 e^{-x}}{4} + \frac{\cosh(x) e^{-x} \sinh(x)}{4} + c_2 e^{-x} + \frac{x e^{-x}}{4} - \frac{e^{-x}}{8}$$

Summary

The solution(s) found are the following

$$y = e^{-x} c_1 x - \frac{x^2 e^{-x}}{4} + \frac{\cosh(x)^2 e^{-x}}{4} + \frac{\cosh(x) e^{-x} \sinh(x)}{4} + c_2 e^{-x} + \frac{x e^{-x}}{4} - \frac{e^{-x}}{8} \quad (1)$$




Figure 116: Slope field plot

Verification of solutions

$$y = e^{-x}c_1x - \frac{x^2e^{-x}}{4} + \frac{\cosh(x)^2 e^{-x}}{4} + \frac{\cosh(x) e^{-x} \sinh(x)}{4} + c_2e^{-x} + \frac{x e^{-x}}{4} - \frac{e^{-x}}{8}$$

Verified OK.

2.24.3 Solving using Kovacic algorithm

Writing the ode as

$$y'' + 2y' + y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 2 \\ C &= 1 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$r = \frac{s}{t} \quad (5)$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$s = 0$$

$$t = 1$$

Therefore eq. (4) becomes

$$z''(x) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 83: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{2}{1} dx} \\ &= z_1 e^{-x} \\ &= z_1 (e^{-x}) \end{aligned}$$

Which simplifies to

$$y_1 = e^{-x}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned} y_2 &= y_1 \int \frac{e^{\int -\frac{2}{1} dx}}{(y_1)^2} dx \\ &= y_1 \int \frac{e^{-2x}}{(y_1)^2} dx \\ &= y_1(x) \end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (e^{-x}) + c_2 (e^{-x}(x))\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + 2y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 e^{-x} + x e^{-x} c_2$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned}y_1 &= e^{-x} \\ y_2 &= x e^{-x}\end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-x} & x e^{-x} \\ \frac{d}{dx}(e^{-x}) & \frac{d}{dx}(x e^{-x}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-x} & x e^{-x} \\ -e^{-x} & e^{-x} - x e^{-x} \end{vmatrix}$$

Therefore

$$W = (e^{-x})(e^{-x} - x e^{-x}) - (x e^{-x})(-e^{-x})$$

Which simplifies to

$$W = e^{-2x}$$

Which simplifies to

$$W = e^{-2x}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{x e^{-x} \sinh(x)}{e^{-2x}} dx$$

Which simplifies to

$$u_1 = - \int \sinh(x) x e^x dx$$

Hence

$$u_1 = - \frac{\sinh(x) x \cosh(x)}{2} + \frac{x^2}{4} + \frac{\cosh(x)^2}{4} - \frac{x \cosh(x)^2}{2} + \frac{\cosh(x) \sinh(x)}{4} + \frac{x}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-x} \sinh(x)}{e^{-2x}} dx$$

Which simplifies to

$$u_2 = \int \sinh(x) e^x dx$$

Hence

$$u_2 = \frac{\cosh(x) \sinh(x)}{2} - \frac{x}{2} + \frac{\cosh(x)^2}{2}$$

Which simplifies to

$$u_1 = \frac{(-2x+1)\cosh(x)^2}{4} + \frac{(-2x+1)\sinh(x)\cosh(x)}{4} + \frac{x^2}{4} + \frac{x}{4}$$

$$u_2 = -\frac{x}{2} + \frac{\sinh(2x)}{4} + \frac{\cosh(2x)}{4} + \frac{1}{4}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(\frac{(-2x+1)\cosh(x)^2}{4} + \frac{(-2x+1)\sinh(x)\cosh(x)}{4} + \frac{x^2}{4} + \frac{x}{4} \right) e^{-x}$$

$$+ \left(-\frac{x}{2} + \frac{\sinh(2x)}{4} + \frac{\cosh(2x)}{4} + \frac{1}{4} \right) x e^{-x}$$

Which simplifies to

$$y_p(x) = \frac{(\cosh(x)\sinh(x) + \cosh(x)^2 - x^2 + x) e^{-x}}{4}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= (c_1 e^{-x} + x e^{-x} c_2) + \left(\frac{(\cosh(x)\sinh(x) + \cosh(x)^2 - x^2 + x) e^{-x}}{4} \right)$$

Which simplifies to

$$y = e^{-x}(c_2 x + c_1) + \frac{(\cosh(x)\sinh(x) + \cosh(x)^2 - x^2 + x) e^{-x}}{4}$$

Summary

The solution(s) found are the following

$$y = e^{-x}(c_2x + c_1) + \frac{(\cosh(x)\sinh(x) + \cosh(x)^2 - x^2 + x)e^{-x}}{4} \quad (1)$$




Figure 117: Slope field plot

Verification of solutions

$$y = e^{-x}(c_2x + c_1) + \frac{(\cosh(x)\sinh(x) + \cosh(x)^2 - x^2 + x)e^{-x}}{4}$$

Verified OK.

2.24.4 Maple step by step solution

Let's solve

$$y'' + 2y' + y = \sinh(x)$$

- Highest derivative means the order of the ODE is 2
- y''
- Characteristic polynomial of homogeneous ODE

$$r^2 + 2r + 1 = 0$$

- Factor the characteristic polynomial

$$(r + 1)^2 = 0$$

- Root of the characteristic polynomial

$$r = -1$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^{-x}$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x e^{-x}$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 e^{-x} + x e^{-x} c_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sinh(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^{-x} & x e^{-x} \\ -e^{-x} & e^{-x} - x e^{-x} \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{-2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = e^{-x} \left(- \left(\int \sinh(x) x e^x dx \right) + x \left(\int \sinh(x) e^x dx \right) \right)$$

- Compute integrals

$$y_p(x) = \frac{e^{-x}(-2x^2+2x+1+\sinh(2x)+\cosh(2x))}{8}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 e^{-x} + x e^{-x} c_2 + \frac{e^{-x}(-2x^2+2x+1+\sinh(2x)+\cosh(2x))}{8}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 32

```
dsolve(diff(y(x),x$2)+2*diff(y(x),x)+y(x)=sinh(x),y(x), singsol=all)
```

$$y(x) = \frac{(-2x^2 + (8c_1 + 2)x + 8c_2 + 1)e^{-x}}{8} + \frac{e^x}{8}$$

✓ Solution by Mathematica

Time used: 0.103 (sec). Leaf size: 34

```
DSolve[y''[x]+2*y'[x]+y[x]==Sinh[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{8}e^{-x}(-2x^2 + e^{2x} + 8c_2x + 8c_1)$$

2.25 problem Problem 34

2.25.1 Maple step by step solution 743

Internal problem ID [12188]

Internal file name [OUTPUT/10840_Thursday_September_21_2023_05_47_51_AM_70798317/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 34.

ODE order: 3.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_3rd_order , _with_linear_symmetries]]
```

$$y''' - y = e^x$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y''' - y = 0$$

The characteristic equation is

$$\lambda^3 - 1 = 0$$

The roots of the above equation are

$$\lambda_1 = 1$$

$$\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

$$\lambda_3 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

Therefore the homogeneous solution is

$$y_h(x) = e^x c_1 + e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_2 + e^{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_3$$

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned} y_1 &= e^x \\ y_2 &= e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} \\ y_3 &= e^{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} \end{aligned}$$

Now the particular solution to the given ODE is found

$$y''' - y = e^x$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^x, e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x}, e^{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} \right\}$$

Since e^x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x e^x\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x e^x$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$3A_1 e^x = e^x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{3} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x e^x}{3}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= \left(e^x c_1 + e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_2 + e^{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_3 \right) + \left(\frac{x e^x}{3} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = e^x c_1 + e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_2 + e^{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_3 + \frac{x e^x}{3} \quad (1)$$

Verification of solutions

$$y = e^x c_1 + e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} c_2 + e^{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)x} c_3 + \frac{x e^x}{3}$$

Verified OK.

2.25.1 Maple step by step solution

Let's solve

$$y''' - y = e^x$$

- Highest derivative means the order of the ODE is 3

$$y'''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $y_1(x)$

$$y_1(x) = y$$

- Define new variable $y_2(x)$

$$y_2(x) = y'$$

- Define new variable $y_3(x)$

$$y_3(x) = y''$$

- Isolate for $y_3'(x)$ using original ODE

$$y_3'(x) = e^x + y_1(x)$$

Convert linear ODE into a system of first order ODEs

$$[y_2(x) = y_1'(x), y_3(x) = y_2'(x), y_3'(x) = e^x + y_1(x)]$$

- Define vector

$$\vec{y}(x) = \begin{bmatrix} y_1(x) \\ y_2(x) \\ y_3(x) \end{bmatrix}$$

- System to solve

$$\vec{y}'(x) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \cdot \vec{y}(x) + \begin{bmatrix} 0 \\ 0 \\ e^x \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(x) = \begin{bmatrix} 0 \\ 0 \\ e^x \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{y}'(x) = A \cdot \vec{y}(x) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\begin{array}{c} \left[\begin{array}{c} 1 \\ 1 \\ 1 \end{array} \right] \\ 1, \left[\begin{array}{c} 1 \\ 1 \\ 1 \end{array} \right] \end{array} \right], \left[\begin{array}{c} -\frac{1}{2} - \frac{i\sqrt{3}}{2}, \left[\begin{array}{c} \frac{1}{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{array} \right] \end{array} \right], \left[\begin{array}{c} -\frac{1}{2} + \frac{i\sqrt{3}}{2}, \left[\begin{array}{c} \frac{1}{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} + \frac{i\sqrt{3}}{2}} \\ 1 \end{array} \right] \end{array} \right]$$

- Consider eigenpair

$$\left[\begin{array}{c} \left[\begin{array}{c} 1 \\ 1 \\ 1 \end{array} \right] \\ 1, \left[\begin{array}{c} 1 \\ 1 \\ 1 \end{array} \right] \end{array} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{y}_1 = e^x \cdot \left[\begin{array}{c} 1 \\ 1 \\ 1 \end{array} \right]$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[\begin{array}{c} -\frac{1}{2} - \frac{i\sqrt{3}}{2}, \left[\begin{array}{c} \frac{1}{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{array} \right] \end{array} \right]$$

- Solution from eigenpair

$$e^{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)x} \cdot \left[\begin{array}{c} \frac{1}{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{array} \right]$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{-\frac{x}{2}} \cdot \left(\cos\left(\frac{\sqrt{3}x}{2}\right) - i \sin\left(\frac{\sqrt{3}x}{2}\right) \right) \cdot \left[\begin{array}{c} \frac{1}{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{array} \right]$$

- Simplify expression

$$e^{-\frac{x}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{3}x}{2}\right) - I \sin\left(\frac{\sqrt{3}x}{2}\right)}{\left(-\frac{1}{2} - \frac{I\sqrt{3}}{2}\right)^2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right) - I \sin\left(\frac{\sqrt{3}x}{2}\right)}{-\frac{1}{2} - \frac{I\sqrt{3}}{2}} \\ \cos\left(\frac{\sqrt{3}x}{2}\right) - I \sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{y}_2(x) = e^{-\frac{x}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \\ \cos\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}, \vec{y}_3(x) = e^{-\frac{x}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ \frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \\ -\sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{y}_p(x)$
 $\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + \vec{y}_p(x)$

□ Fundamental matrix

- Let $\phi(x)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(x) = \begin{bmatrix} e^x & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \right) & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \right) \\ e^x & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \right) & e^{-\frac{x}{2}} \left(\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \right) \\ e^x & e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & -e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix}$$

- The fundamental matrix, $\Phi(x)$ is a normalized version of $\phi(x)$ satisfying $\Phi(0) = I$ where I is the identity matrix

$$\Phi(x) = \phi(x) \cdot \frac{1}{\phi(0)}$$

- Substitute the value of $\phi(x)$ and $\phi(0)$

$$\Phi(x) = \begin{bmatrix} e^x & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \right) & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \right) \\ e^x & e^{-\frac{x}{2}} \left(-\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} \right) & e^{-\frac{x}{2}} \left(\frac{\cos\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}x}{2}\right)}{2} \right) \\ e^x & e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) & -e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) \end{bmatrix} \cdot \begin{bmatrix} 1 & -\frac{1}{2} & -\frac{\sqrt{3}}{2} \\ 1 & -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ 1 & 1 & 0 \end{bmatrix}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(x) = \begin{bmatrix} \frac{e^x}{3} + \frac{2e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} & \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} + \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{3} & \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} \\ \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} - \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{3} & \frac{e^x}{3} + \frac{2e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} & \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} \\ \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} + \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{3} & \frac{e^x}{3} - \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} - \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{3} & \frac{e^x}{3} + \frac{2e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} \end{bmatrix}$$

□

Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(x)$ and solve for $\vec{v}(x)$

$$\vec{y}_p(x) = \Phi(x) \cdot \vec{v}(x)$$

- Take the derivative of the particular solution

$$\vec{y}'_p(x) = \Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(x)$

$$A \cdot \Phi(x) \cdot \vec{v}(x) + \Phi(x) \cdot \vec{v}'(x) = A \cdot \Phi(x) \cdot \vec{v}(x) + \vec{f}(x)$$

- Cancel like terms

$$\Phi(x) \cdot \vec{v}'(x) = \vec{f}(x)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(x) = \frac{1}{\Phi(x)} \cdot \vec{f}(x)$$

- Integrate to solve for $\vec{v}(x)$

$$\vec{v}(x) = \int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(x)$ into the equation for the particular solution

$$\vec{y}_p(x) = \Phi(x) \cdot \left(\int_0^x \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{y}_p(x) = \begin{bmatrix} \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} + \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{9} + \frac{e^x(x-1)}{3} \\ -\frac{2e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{9} + \frac{x e^x}{3} \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} + \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{9} + \frac{e^x(x+1)}{3} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{y}(x) = c_1 \vec{y}_1 + c_2 \vec{y}_2(x) + c_3 \vec{y}_3(x) + \begin{bmatrix} \frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} + \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{9} + \frac{e^x(x-1)}{3} \\ -\frac{2e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{9} + \frac{x e^x}{3} \\ -\frac{e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{3} + \frac{e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)\sqrt{3}}{9} + \frac{e^x(x+1)}{3} \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$y = -\frac{(c_3\sqrt{3}+c_2-\frac{2}{3})e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right)}{2} - \frac{((c_2-\frac{2}{9})\sqrt{3}-c_3)e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right)}{2} + \frac{e^x(x+3c_1-1)}{3}$$

Maple trace

```

`Methods for third order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 3; linear nonhomogeneous with symmetry [0,1]
trying high order linear exact nonhomogeneous
trying differential order: 3; missing the dependent variable
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 40

```
dsolve(diff(y(x),x$3)-y(x)=exp(x),y(x), singsol=all)
```

$$y(x) = c_2 e^{-\frac{x}{2}} \cos\left(\frac{\sqrt{3}x}{2}\right) + c_3 e^{-\frac{x}{2}} \sin\left(\frac{\sqrt{3}x}{2}\right) + \frac{e^x(x+3c_1)}{3}$$

✓ Solution by Mathematica

Time used: 0.726 (sec). Leaf size: 62

```
DSolve[y'''[x]-y[x]==Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{3} e^{-x/2} \left(e^{3x/2} (x-1+3c_1) + 3c_2 \cos\left(\frac{\sqrt{3}x}{2}\right) + 3c_3 \sin\left(\frac{\sqrt{3}x}{2}\right) \right)$$

2.26 problem Problem 35

2.26.1 Solving as second order linear constant coeff ode	749
2.26.2 Solving using Kovacic algorithm	753
2.26.3 Maple step by step solution	758

Internal problem ID [12189]

Internal file name [OUTPUT/10841_Thursday_September_21_2023_05_47_51_AM_89062283/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 35.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' - 2y' + 2y = x e^x \cos(x)$$

2.26.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = -2, C = 2, f(x) = x e^x \cos(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + 2y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = -2, C = 2$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} - 2\lambda e^{\lambda x} + 2 e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 - 2\lambda + 2 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = -2, C = 2$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{2}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{-2^2 - (4)(1)(2)} \\ &= 1 \pm i \end{aligned}$$

Hence

$$\lambda_1 = 1 + i$$

$$\lambda_2 = 1 - i$$

Which simplifies to

$$\lambda_1 = 1 + i$$

$$\lambda_2 = 1 - i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 1$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^x (c_1 \cos(x) + c_2 \sin(x))$$

Therefore the homogeneous solution y_h is

$$y_h = e^x (c_1 \cos(x) + c_2 \sin(x))$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x e^x \cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^x \cos(x), e^x \sin(x), x e^x \cos(x), x e^x \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x \cos(x), e^x \sin(x)\}$$

Since $e^x \cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x e^x \cos(x), x e^x \sin(x), \cos(x) e^x x^2, e^x \sin(x) x^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x e^x \cos(x) + A_2 x e^x \sin(x) + A_3 \cos(x) e^x x^2 + A_4 e^x \sin(x) x^2$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_3 \cos(x) e^x + 2A_4 e^x \sin(x) - 2A_1 e^x \sin(x) + 2A_2 e^x \cos(x) - 4A_3 \sin(x) e^x x + 4A_4 e^x \cos(x) x = x e^x \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{4}, A_2 = 0, A_3 = 0, A_4 = \frac{1}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x e^x \cos(x)}{4} + \frac{e^x \sin(x) x^2}{4}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= (e^x(c_1 \cos(x) + c_2 \sin(x))) + \left(\frac{x e^x \cos(x)}{4} + \frac{e^x \sin(x) x^2}{4} \right)$$

Summary

The solution(s) found are the following

$$y = e^x(c_1 \cos(x) + c_2 \sin(x)) + \frac{x e^x \cos(x)}{4} + \frac{e^x \sin(x) x^2}{4} \quad (1)$$




Figure 118: Slope field plot

Verification of solutions

$$y = e^x(c_1 \cos(x) + c_2 \sin(x)) + \frac{x e^x \cos(x)}{4} + \frac{e^x \sin(x) x^2}{4}$$

Verified OK.

2.26.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' - 2y' + 2y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= -2 \\ C &= 2 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -1 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x) \quad (7)$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 86: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - 0 \\ &= 0 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$\begin{aligned} y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\ &= z_1 e^{-\int \frac{1}{2} \frac{-2}{1} dx} \end{aligned}$$

$$\begin{aligned}
&= z_1 e^x \\
&= z_1 (e^x)
\end{aligned}$$

Which simplifies to

$$y_1 = e^x \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}
y_2 &= y_1 \int \frac{e^{\int -\frac{-2}{1} dx}}{(y_1)^2} dx \\
&= y_1 \int \frac{e^{2x}}{(y_1)^2} dx \\
&= y_1 (\tan(x))
\end{aligned}$$

Therefore the solution is

$$\begin{aligned}
y &= c_1 y_1 + c_2 y_2 \\
&= c_1 (e^x \cos(x)) + c_2 (e^x \cos(x) (\tan(x)))
\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' - 2y' + 2y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = e^x \cos(x) c_1 + e^x \sin(x) c_2$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x e^x \cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^x \cos(x), e^x \sin(x), x e^x \cos(x), x e^x \sin(x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{e^x \cos(x), e^x \sin(x)\}$$

Since $e^x \cos(x)$ is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x e^x \cos(x), x e^x \sin(x), \cos(x) e^x x^2, e^x \sin(x) x^2\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1 x e^x \cos(x) + A_2 x e^x \sin(x) + A_3 \cos(x) e^x x^2 + A_4 e^x \sin(x) x^2$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$2A_4 e^x \sin(x) + 2A_3 \cos(x) e^x + 4A_4 e^x \cos(x) x - 4A_3 \sin(x) e^x x - 2A_1 e^x \sin(x) + 2A_2 e^x \cos(x) = x e^x \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{4}, A_2 = 0, A_3 = 0, A_4 = \frac{1}{4} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{x e^x \cos(x)}{4} + \frac{e^x \sin(x) x^2}{4}$$

Therefore the general solution is

$$y = y_h + y_p$$

$$= (e^x \cos(x) c_1 + e^x \sin(x) c_2) + \left(\frac{x e^x \cos(x)}{4} + \frac{e^x \sin(x) x^2}{4} \right)$$

Which simplifies to

$$y = e^x(c_1 \cos(x) + c_2 \sin(x)) + \frac{x e^x \cos(x)}{4} + \frac{e^x \sin(x) x^2}{4}$$

Summary

The solution(s) found are the following

$$y = e^x(c_1 \cos(x) + c_2 \sin(x)) + \frac{x e^x \cos(x)}{4} + \frac{e^x \sin(x) x^2}{4} \quad (1)$$




Figure 119: Slope field plot

Verification of solutions

$$y = e^x(c_1 \cos(x) + c_2 \sin(x)) + \frac{x e^x \cos(x)}{4} + \frac{e^x \sin(x) x^2}{4}$$

Verified OK.

2.26.3 Maple step by step solution

Let's solve

$$y'' - 2y' + 2y = x e^x \cos(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 - 2r + 2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{2 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (1 - i, 1 + i)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = e^x \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = e^x \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = e^x \cos(x) c_1 + e^x \sin(x) c_2 + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = x e^x \cos(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} e^x \cos(x) & e^x \sin(x) \\ e^x \cos(x) - e^x \sin(x) & e^x \cos(x) + e^x \sin(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = e^{2x}$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{e^x(\cos(x)(\int \sin(2x)xdx) - 2\sin(x)(\int \cos(x)^2xdx))}{2}$$

- Compute integrals

$$y_p(x) = \frac{(\sin(x)x^2 + \cos(x)x - \sin(x))e^x}{4}$$

- Substitute particular solution into general solution to ODE

$$y = e^x \cos(x) c_1 + e^x \sin(x) c_2 + \frac{(\sin(x)x^2 + \cos(x)x - \sin(x))e^x}{4}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 28

```
dsolve(diff(y(x),x$2)-2*diff(y(x),x) +2*y(x)=x*exp(x)*cos(x),y(x), singsol=all)
```

$$y(x) = \frac{e^x((x^2 + 4c_2 - 1) \sin(x) + \cos(x)(4c_1 + x))}{4}$$

✓ Solution by Mathematica

Time used: 0.074 (sec). Leaf size: 37

```
DSolve[y''[x]-2*y'[x] +2*y[x]==x*Exp[x]*Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{8}e^x((2x^2 - 1 + 8c_1) \sin(x) + 2(x + 4c_2) \cos(x))$$

2.27 problem Problem 36

2.27.1 Solving using Kovacic algorithm 760

Internal problem ID [12190]

Internal file name [OUTPUT/10842_Thursday_September_21_2023_05_47_55_AM_26792915/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 36.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : **"kovacic"**

Maple gives the following as the ode type

```
[[_2nd_order , _with_linear_symmetries]]
```

$$(x^2 - 1)y'' - 6y = 1$$

2.27.1 Solving using Kovacic algorithm

Writing the ode as

$$(x^2 - 1)y'' - 6y = 0 \tag{1}$$

$$Ay'' + By' + Cy = 0 \tag{2}$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= x^2 - 1 \\ B &= 0 \\ C &= -6 \end{aligned} \tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$r = \frac{s}{t} \tag{5}$$

$$= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{6}{x^2 - 1} \tag{6}$$

Comparing the above to (5) shows that

$$s = 6$$

$$t = x^2 - 1$$

Therefore eq. (4) becomes

$$z''(x) = \left(\frac{6}{x^2 - 1} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 88: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = x^2 - 1$. There is a pole at $x = 1$ of order 1. There is a pole at $x = -1$ of order 1. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 1. For the pole at $x = 1$ of order 1 then

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= 1 \\ \alpha_c^- &= 1 \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ . which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = \frac{6}{x^2 - 1}$$

Since the $\gcd(s, t) = 1$. This gives $b = 6$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = 3 \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = -2 \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = \frac{6}{x^2 - 1}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
1	1	0	0	1

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	3	-2

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω . Trying $\alpha_\infty^+ = 3$ then

$$\begin{aligned} d &= \alpha_\infty^+ - (\alpha_{c_1}^-) \\ &= 3 - (1) \\ &= 2 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c) [\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty) [\sqrt{r}]_\infty$$

Substituting the above values in the above results in

$$\begin{aligned} \omega &= \left((-) [\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^-}{x - c_1} \right) + (+) [\sqrt{r}]_\infty \\ &= \frac{1}{x - 1} + (0) \\ &= \frac{1}{x - 1} \\ &= \frac{1}{x - 1} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 2$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r) p = 0 \quad (1A)$$

Let

$$p(x) = x^2 + a_1 x + a_0 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$(2) + 2\left(\frac{1}{x-1}\right)(2x+a_1) + \left(\left(-\frac{1}{(x-1)^2}\right) + \left(\frac{1}{x-1}\right)^2 - \left(\frac{6}{x^2-1}\right)\right) = 0$$

$$\frac{(-4a_1+4)x - 6a_0 + 2a_1 - 2}{x^2-1} = 0$$

Solving for the coefficients a_i in the above using method of undetermined coefficients gives

$$\{a_0 = 0, a_1 = 1\}$$

Substituting these coefficients in $p(x)$ in eq. (2A) results in

$$p(x) = x^2 + x$$

Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= (x^2 + x) e^{\int \frac{1}{x-1} dx} \\ &= (x^2 + x) x - 1 \\ &= x^3 - x \end{aligned}$$

The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned} y_1 &= z_1 \\ &= x^3 - x \end{aligned}$$

Which simplifies to

$$y_1 = x^3 - x$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= x^3 - x \int \frac{1}{(x^3 - x)^2} dx \\ &= x^3 - x \left(-\frac{1}{4x - 4} - \frac{3 \ln(x - 1)}{4} - \frac{1}{4 + 4x} + \frac{3 \ln(x + 1)}{4} - \frac{1}{x} \right) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (x^3 - x) + c_2 \left(x^3 - x \left(-\frac{1}{4x - 4} - \frac{3 \ln(x - 1)}{4} - \frac{1}{4 + 4x} + \frac{3 \ln(x + 1)}{4} - \frac{1}{x} \right) \right) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$(x^2 - 1)y'' - 6y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 (x^3 - x) + c_2 \left(\frac{(3x^3 - 3x) \ln(x + 1)}{4} + \frac{(-3x^3 + 3x) \ln(x - 1)}{4} - \frac{3x^2}{2} + 1 \right)$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = x^3 - x$$

$$y_2 = \frac{(3x^3 - 3x) \ln(x+1)}{4} + \frac{(-3x^3 + 3x) \ln(x-1)}{4} - \frac{3x^2}{2} + 1$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} x^3 - x & \frac{(3x^3-3x) \ln(x+1)}{4} + \frac{(-3x^3+3x) \ln(x-1)}{4} - \frac{3x^2}{2} + 1 \\ \frac{d}{dx}(x^3 - x) & \frac{d}{dx} \left(\frac{(3x^3-3x) \ln(x+1)}{4} + \frac{(-3x^3+3x) \ln(x-1)}{4} - \frac{3x^2}{2} + 1 \right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} x^3 - x & \frac{(3x^3-3x) \ln(x+1)}{4} + \frac{(-3x^3+3x) \ln(x-1)}{4} - \frac{3x^2}{2} + 1 \\ 3x^2 - 1 & \frac{(9x^2-3) \ln(x+1)}{4} + \frac{3x^3-3x}{4+4x} + \frac{(-9x^2+3) \ln(x-1)}{4} + \frac{-3x^3+3x}{4x-4} - 3x \end{vmatrix}$$

Therefore

$$W = (x^3 - x) \left(\frac{(9x^2 - 3) \ln(x+1)}{4} + \frac{3x^3 - 3x}{4 + 4x} + \frac{(-9x^2 + 3) \ln(x-1)}{4} + \frac{-3x^3 + 3x}{4x - 4} - 3x \right) - \left(\frac{(3x^3 - 3x) \ln(x+1)}{4} + \frac{(-3x^3 + 3x) \ln(x-1)}{4} - \frac{3x^2}{2} + 1 \right) (3x^2 - 1)$$

Which simplifies to

$$W = 1$$

Which simplifies to

$$W = 1$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{\frac{(3x^3-3x)\ln(x+1)}{4} + \frac{(-3x^3+3x)\ln(x-1)}{4} - \frac{3x^2}{2} + 1}{x^2 - 1} dx$$

Which simplifies to

$$u_1 = - \int \frac{(3x^3 - 3x) \ln(x + 1) + (-3x^3 + 3x) \ln(x - 1) - 6x^2 + 4}{4x^2 - 4} dx$$

Hence

$$u_1 = \frac{3x}{4} + \frac{\ln(x-1)}{4} - \frac{\ln(x+1)}{4} + \frac{3(x-1)^2 \ln(x-1)}{8} \\ + \frac{3(x-1) \ln(x-1)}{4} - \frac{3(x+1)^2 \ln(x+1)}{8} + \frac{3(x+1) \ln(x+1)}{4}$$

And Eq. (3) becomes

$$u_2 = \int \frac{x^3 - x}{x^2 - 1} dx$$

Which simplifies to

$$u_2 = \int x dx$$

Hence

$$u_2 = \frac{x^2}{2}$$

Which simplifies to

$$u_1 = \frac{3 \ln(x-1) x^2}{8} - \frac{3 \ln(x+1) x^2}{8} - \frac{\ln(x-1)}{8} + \frac{\ln(x+1)}{8} + \frac{3x}{4} \\ u_2 = \frac{x^2}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \left(\frac{3 \ln(x-1)x^2}{8} - \frac{3 \ln(x+1)x^2}{8} - \frac{\ln(x-1)}{8} + \frac{\ln(x+1)}{8} + \frac{3x}{4} \right) (x^3 - x) + \frac{x^2 \left(\frac{(3x^3-3x)\ln(x+1)}{4} + \frac{(-3x^3+3x)\ln(x-1)}{4} - \frac{3x^2}{2} + 1 \right)}{2}$$

Which simplifies to

$$y_p(x) = \frac{x((x^2-1)\ln(x+1) - \ln(x-1)x^2 - 2x + \ln(x-1))}{8}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1(x^3 - x) + c_2 \left(\frac{(3x^3 - 3x)\ln(x+1)}{4} + \frac{(-3x^3 + 3x)\ln(x-1)}{4} - \frac{3x^2}{2} + 1 \right) \right) + \left(\frac{x((x^2-1)\ln(x+1) - \ln(x-1)x^2 - 2x + \ln(x-1))}{8} \right)$$

Which simplifies to

$$y = \frac{3c_2(x^3 - x)\ln(x+1)}{4} + \frac{3(-x^3 + x)c_2\ln(x-1)}{4} + c_1x^3 - \frac{3c_2x^2}{2} - c_1x + c_2 + \frac{x((x^2-1)\ln(x+1) - \ln(x-1)x^2 - 2x + \ln(x-1))}{8}$$

Summary

The solution(s) found are the following

$$y = \frac{3c_2(x^3 - x)\ln(x+1)}{4} + \frac{3(-x^3 + x)c_2\ln(x-1)}{4} + c_1x^3 - \frac{3c_2x^2}{2} - c_1x + c_2 + \frac{x((x^2-1)\ln(x+1) - \ln(x-1)x^2 - 2x + \ln(x-1))}{8} \quad (1)$$

Verification of solutions

$$y = \frac{3c_2(x^3 - x)\ln(x+1)}{4} + \frac{3(-x^3 + x)c_2\ln(x-1)}{4} + c_1x^3 - \frac{3c_2x^2}{2} - c_1x + c_2 + \frac{x((x^2-1)\ln(x+1) - \ln(x-1)x^2 - 2x + \ln(x-1))}{8}$$

Verified OK.

Maple trace Kovacic algorithm successful

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    checking if the LODE is of Euler type
    trying a symmetry of the form [xi=0, eta=F(x)]
    checking if the LODE is missing y
-> Trying a Liouvillian solution using Kovacics algorithm
    A Liouvillian solution exists
    Reducible group (found an exponential solution)
    Group is reducible, not completely reducible
<- Kovacics algorithm successful
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 52

```
dsolve((x^2-1)*diff(y(x),x$2)-6*y(x)=1,y(x), singsol=all)
```

$$y(x) = -\frac{1}{6} + \frac{3(x^3 - x)c_1 \ln(-1 + x)}{4} + \frac{3c_1(-x^3 + x) \ln(1 + x)}{4} + c_2 x^3 + \frac{3c_1 x^2}{2} - c_2 x - c_1$$

✓ Solution by Mathematica

Time used: 0.086 (sec). Leaf size: 67

```
DSolve[(x^2-1)*y''[x]-6*y[x]==1,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{12} \left(-9c_2 x(x^2 - 1) \log(1 - x) + 9c_2 x(x^2 - 1) \log(x + 1) \right. \\ \left. + 2(6c_1 x^3 - 9c_2 x^2 - 6c_1 x - 1 + 6c_2) \right)$$

2.28 problem Problem 40(a)

Internal problem ID [12191]

Internal file name [OUTPUT/10843_Thursday_September_21_2023_05_47_56_AM_22119359/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 40(a).

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]
```

Unable to solve or complete the solution.

$$mx'' - f(x) = 0$$

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
`, `-> Computing symmetries using: way = exp_sym
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)-f(_a)/m = 0, _b(_a)` *** Suble
  Methods for first order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying 1st order linear
  trying Bernoulli
  <- Bernoulli successful
<- differential order: 2; canonical coordinates successful
<- differential order 2; missing variables successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 62

```
dsolve(m*diff(x(t),t$2)=f(x(t)),x(t), singsol=all)
```

$$m \left(\int^{x(t)} \frac{1}{\sqrt{m(c_1 m + 2(\int f(_b) d_b))}} d_b \right) - t - c_2 = 0$$
$$-m \left(\int^{x(t)} \frac{1}{\sqrt{m(c_1 m + 2(\int f(_b) d_b))}} d_b \right) - t - c_2 = 0$$

✓ Solution by Mathematica

Time used: 0.049 (sec). Leaf size: 44

```
DSolve[m*x'[t]==f[x[t]],x[t],t,IncludeSingularSolutions -> True]
```

$$\text{Solve} \left[\int_1^{x(t)} \frac{1}{\sqrt{c_1 + 2 \int_1^{K[2]} \frac{f(K[1])}{m} dK[1]}} dK[2]^2 = (t + c_2)^2, x(t) \right]$$

2.29 problem Problem 40(b)

2.29.1 Solving as second order ode missing y ode 773

2.29.2 Solving as second order ode missing x ode 774

Internal problem ID [12192]

Internal file name [OUTPUT/10844_Thursday_September_21_2023_05_47_56_AM_69214506/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 40(b).

ODE order: 2.

ODE degree: 0.

The type(s) of ODE detected by this program : "second_order_ode_missing_x", "second_order_ode_missing_y"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_x]]
```

$$mx'' - f(x') = 0$$

2.29.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable x . Let

$$p(t) = x'$$

Then

$$p'(t) = x''$$

Hence the ode becomes

$$mp'(t) - f(p(t)) = 0$$

Which is now solve for $p(t)$ as first order ode. Integrating both sides gives

$$\int \frac{m}{f(p)} dp = \int dt$$
$$\int^{p(t)} \frac{m}{f(a)} da = t + c_1$$

For solution (1) found earlier, since $p = x'$ then we now have a new first order ode to solve which is

$$\int^{x'} \frac{m}{f(-a)} d_{-a} = t + c_1$$

Integrating both sides gives

$$\begin{aligned} x &= \int \text{RootOf} \left(- \left(\int^{-Z} \frac{m}{f(-a)} d_{-a} \right) + t + c_1 \right) dt \\ &= \int \text{RootOf} \left(- \left(\int^{-Z} \frac{m}{f(-a)} d_{-a} \right) + t + c_1 \right) dt + c_2 \end{aligned}$$

Summary

The solution(s) found are the following

$$x = \int \text{RootOf} \left(- \left(\int^{-Z} \frac{m}{f(-a)} d_{-a} \right) + t + c_1 \right) dt + c_2 \quad (1)$$

Verification of solutions

$$x = \int \text{RootOf} \left(- \left(\int^{-Z} \frac{m}{f(-a)} d_{-a} \right) + t + c_1 \right) dt + c_2$$

Verified OK.

2.29.2 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable x an independent variable. Using

$$x' = p(x)$$

Then

$$\begin{aligned} x'' &= \frac{dp}{dt} \\ &= \frac{dx}{dt} \frac{dp}{dx} \\ &= p \frac{dp}{dx} \end{aligned}$$

Hence the ode becomes

$$mp(x) \left(\frac{d}{dx} p(x) \right) = f(p(x))$$

Which is now solved as first order ode for $p(x)$. Integrating both sides gives

$$\int \frac{mp}{f(p)} dp = \int dx$$

$$\int^{p(x)} \frac{m_a}{f(a)} da = x + c_1$$

For solution (1) found earlier, since $p = x'$ then we now have a new first order ode to solve which is

$$\int^{x'} \frac{m_a}{f(a)} da = x + c_1$$

Integrating both sides gives

$$\int \frac{1}{\text{RootOf}\left(-\left(\int^{-Z} \frac{m_a}{f(a)} da\right) + x + c_1\right)} dx = \int dt$$

$$\int^x \frac{1}{\text{RootOf}\left(-\left(\int^{-Z} \frac{m_a}{f(a)} da\right) + a + c_1\right)} da = t + c_2$$

Summary

The solution(s) found are the following

$$\int^x \frac{1}{\text{RootOf}\left(-\left(\int^{-Z} \frac{m_a}{f(a)} da\right) + a + c_1\right)} da = t + c_2 \quad (1)$$

Verification of solutions

$$\int^x \frac{1}{\text{RootOf}\left(-\left(\int^{-Z} \frac{m_a}{f(a)} da\right) + a + c_1\right)} da = t + c_2$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying 2nd order Liouville  
trying 2nd order WeierstrassP  
trying 2nd order JacobiSN  
differential order: 2; trying a linearization to 3rd order  
trying 2nd order ODE linearizable_by_differentiation  
trying 2nd order, 2 integrating factors of the form mu(x,y)  
trying differential order: 2; missing variables  
, ` -> Computing symmetries using: way = 3  
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = f(_b(_a))/m, _b(_a), HINT = [[1, 0]]`  
symmetry methods on request  
, `1st order, trying reduction of order with given symmetries:` [1, 0]
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 23

```
dsolve(m*diff(x(t),t$2)=f(diff(x(t),t)),x(t), singsol=all)
```

$$x(t) = \int \text{RootOf} \left(t - m \left(\int^{-Z} \frac{1}{f(_f)} d_f \right) + c_1 \right) dt + c_2$$

✓ Solution by Mathematica

Time used: 2.257 (sec). Leaf size: 39

```
DSolve[m*x''[t]==f[x'[t]],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \int_1^t \text{InverseFunction} \left[\int_1^{\#1} \frac{1}{f(K[1])} dK[1] \& \right] \left[c_1 + \frac{K[2]}{m} \right] dK[2] + c_2$$

2.30 problem Problem 41

Internal problem ID [12193]

Internal file name [OUTPUT/10845_Thursday_September_21_2023_05_47_57_AM_43712024/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 41.

ODE order: 6.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_y]]
```

$$y^{(6)} - 3y^{(5)} + 3y'''' - y''' = x$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y^{(6)} - 3y^{(5)} + 3y'''' - y''' = 0$$

The characteristic equation is

$$\lambda^6 - 3\lambda^5 + 3\lambda^4 - \lambda^3 = 0$$

The roots of the above equation are

$$\lambda_1 = 0$$

$$\lambda_2 = 0$$

$$\lambda_3 = 0$$

$$\lambda_4 = 1$$

$$\lambda_5 = 1$$

$$\lambda_6 = 1$$

Therefore the homogeneous solution is

$$y_h(x) = c_3x^2 + c_2x + c_1 + e^x c_4 + x e^x c_5 + x^2 e^x c_6$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = 1$$

$$y_2 = x$$

$$y_3 = x^2$$

$$y_4 = e^x$$

$$y_5 = x e^x$$

$$y_6 = x^2 e^x$$

Now the particular solution to the given ODE is found

$$y^{(6)} - 3y^{(5)} + 3y'''' - y''' = x$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x, x^2, x e^x, x^2 e^x, e^x\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x, x^2\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^2, x^3\}]$$

Since x^2 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{x^3, x^4\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_2x^4 + A_1x^3$$

The unknowns $\{A_1, A_2\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-24xA_2 - 6A_1 + 72A_2 = x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = -\frac{1}{2}, A_2 = -\frac{1}{24} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{1}{24}x^4 - \frac{1}{2}x^3$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_3x^2 + c_2x + c_1 + e^x c_4 + x e^x c_5 + x^2 e^x c_6) + \left(-\frac{1}{24}x^4 - \frac{1}{2}x^3 \right) \end{aligned}$$

Which simplifies to

$$y = (c_6x^2 + c_5x + c_4) e^x + c_3x^2 + c_2x + c_1 - \frac{x^4}{24} - \frac{x^3}{2}$$

Summary

The solution(s) found are the following

$$y = (c_6x^2 + c_5x + c_4) e^x + c_3x^2 + c_2x + c_1 - \frac{x^4}{24} - \frac{x^3}{2} \quad (1)$$

Verification of solutions

$$y = (c_6x^2 + c_5x + c_4) e^x + c_3x^2 + c_2x + c_1 - \frac{x^4}{24} - \frac{x^3}{2}$$

Verified OK.

Maple trace

```
`Methods for high order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 6; linear nonhomogeneous with symmetry [0,1]
-> Calling odsolve with the ODE`, diff(diff(diff(_b(_a), _a), _a), _a) = 3*(diff(diff(_b(_a)
  Methods for third order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying high order exact linear fully integrable
  trying differential order: 3; linear nonhomogeneous with symmetry [0,1]
  trying high order linear exact nonhomogeneous
  trying differential order: 3; missing the dependent variable
  checking if the LODE has constant coefficients
  <- constant coefficients successful
<- differential order: 6; linear nonhomogeneous with symmetry [0,1] successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 48

```
dsolve(diff(y(x),x$6)-3*diff(y(x),x$5)+3*diff(y(x),x$4)-diff(y(x),x$3)=x,y(x), singsol=all)
```

$$y(x) = (c_3x^2 + (c_2 - 6c_3)x + c_1 - 3c_2 + 12c_3)e^x - \frac{x^4}{24} - \frac{x^3}{2} + \frac{c_4x^2}{2} + c_5x + c_6$$

✓ Solution by Mathematica

Time used: 0.256 (sec). Leaf size: 61

```
DSolve[y''''''[x]-3*y''''''[x]+3*y''''''[x]-y''''[x]==x,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow -\frac{x^4}{24} - \frac{x^3}{2} + c_6x^2 + c_3e^x(x^2 - 6x + 12) + c_5x + c_1e^x + c_2e^x(x - 3) + c_4$$

2.31 problem Problem 42

Internal problem ID [12194]

Internal file name [OUTPUT/10846_Thursday_September_21_2023_05_47_57_AM_65265185/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 42.

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _linear , _nonhomogeneous]]
```

$$x'''' + 2x'' + x = \cos(t)$$

This is higher order nonhomogeneous ODE. Let the solution be

$$x = x_h + x_p$$

Where x_h is the solution to the homogeneous ODE And x_p is a particular solution to the nonhomogeneous ODE. x_h is the solution to

$$x'''' + 2x'' + x = 0$$

The characteristic equation is

$$\lambda^4 + 2\lambda^2 + 1 = 0$$

The roots of the above equation are

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

$$\lambda_3 = i$$

$$\lambda_4 = -i$$

Therefore the homogeneous solution is

$$x_h(t) = e^{it}c_1 + te^{it}c_2 + e^{-it}c_3 + te^{-it}c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned}x_1 &= e^{it} \\x_2 &= e^{it}t \\x_3 &= e^{-it} \\x_4 &= e^{-it}t\end{aligned}$$

Now the particular solution to the given ODE is found

$$x'''' + 2x'' + x = \cos(t)$$

Let the particular solution be

$$x_p = U_1x_1 + U_2x_2 + U_3x_3 + U_4x_4$$

Where x_i are the basis solutions found above for the homogeneous solution x_h and $U_i(t)$ are functions to be determined as follows

$$U_i = (-1)^{n-i} \int \frac{F(t)W_i(t)}{aW(t)} dt$$

Where $W(t)$ is the Wronskian and $W_i(t)$ is the Wronskian that results after deleting the last row and the i -th column of the determinant and n is the order of the ODE or equivalently, the number of basis solutions, and a is the coefficient of the leading derivative in the ODE, and $F(t)$ is the RHS of the ODE. Therefore, the first step is to find the Wronskian $W(t)$. This is given by

$$W(t) = \begin{vmatrix} x_1 & x_2 & x_3 & x_4 \\ x_1' & x_2' & x_3' & x_4' \\ x_1'' & x_2'' & x_3'' & x_4'' \\ x_1''' & x_2''' & x_3''' & x_4''' \end{vmatrix}$$

Substituting the fundamental set of solutions x_i found above in the Wronskian gives

$$W = \begin{bmatrix} e^{it} & e^{it}t & e^{-it} & e^{-it}t \\ ie^{it} & e^{it}(it+1) & -ie^{-it} & e^{-it}(-it+1) \\ -e^{it} & e^{it}(2i-t) & -e^{-it} & e^{-it}(-2i-t) \\ -ie^{it} & -e^{it}(it+3) & ie^{-it} & e^{-it}(it-3) \end{bmatrix}$$

$$|W| = 16e^{2it}e^{-2it}$$

The determinant simplifies to

$$|W| = 16$$

Now we determine W_i for each U_i .

$$\begin{aligned} W_1(t) &= \det \begin{bmatrix} e^{it} & e^{-it} & e^{-it}t \\ e^{it}(it+1) & -ie^{-it} & e^{-it}(-it+1) \\ e^{it}(2i-t) & -e^{-it} & e^{-it}(-2i-t) \end{bmatrix} \\ &= -4e^{-it}(-i+t) \end{aligned}$$

$$\begin{aligned} W_2(t) &= \det \begin{bmatrix} e^{it} & e^{-it} & e^{-it}t \\ ie^{it} & -ie^{-it} & e^{-it}(-it+1) \\ -e^{it} & -e^{-it} & e^{-it}(-2i-t) \end{bmatrix} \\ &= -4e^{-it} \end{aligned}$$

$$\begin{aligned} W_3(t) &= \det \begin{bmatrix} e^{it} & e^{it}t & e^{-it}t \\ ie^{it} & e^{it}(it+1) & e^{-it}(-it+1) \\ -e^{it} & e^{it}(2i-t) & e^{-it}(-2i-t) \end{bmatrix} \\ &= -4e^{it}(i+t) \end{aligned}$$

$$\begin{aligned} W_4(t) &= \det \begin{bmatrix} e^{it} & e^{it}t & e^{-it} \\ ie^{it} & e^{it}(it+1) & -ie^{-it} \\ -e^{it} & e^{it}(2i-t) & -e^{-it} \end{bmatrix} \\ &= -4e^{it} \end{aligned}$$

Now we are ready to evaluate each $U_i(t)$.

$$\begin{aligned} U_1 &= (-1)^{4-1} \int \frac{F(t)W_1(t)}{aW(t)} dt \\ &= (-1)^3 \int \frac{(\cos(t))(-4e^{-it}(-i+t))}{(1)(16)} dt \\ &= - \int \frac{-4 \cos(t) e^{-it}(-i+t)}{16} dt \\ &= - \int \left(-\frac{\cos(t) e^{-it}(-i+t)}{4} \right) dt \\ &= - \left(\int -\frac{\cos(t) e^{-it}(-i+t)}{4} dt \right) \end{aligned}$$

$$\begin{aligned}
U_2 &= (-1)^{4-2} \int \frac{F(t)W_2(t)}{aW(t)} dt \\
&= (-1)^2 \int \frac{(\cos(t))(-4e^{-it})}{(1)(16)} dt \\
&= \int \frac{-4 \cos(t) e^{-it}}{16} dt \\
&= \int \left(-\frac{\cos(t) e^{-it}}{4} \right) dt \\
&= \int -\frac{\cos(t) e^{-it}}{4} dt
\end{aligned}$$

$$\begin{aligned}
U_3 &= (-1)^{4-3} \int \frac{F(t)W_3(t)}{aW(t)} dt \\
&= (-1)^1 \int \frac{(\cos(t))(-4e^{it}(i+t))}{(1)(16)} dt \\
&= - \int \frac{-4 \cos(t) e^{it}(i+t)}{16} dt \\
&= - \int \left(-\frac{\cos(t) e^{it}(i+t)}{4} \right) dt \\
&= \frac{t^2}{16} + \frac{it}{8} - \frac{i(3i+2t)e^{2it}}{32}
\end{aligned}$$

$$\begin{aligned}
U_4 &= (-1)^{4-4} \int \frac{F(t)W_4(t)}{aW(t)} dt \\
&= (-1)^0 \int \frac{(\cos(t))(-4e^{it})}{(1)(16)} dt \\
&= \int \frac{-4 \cos(t) e^{it}}{16} dt \\
&= \int \left(-\frac{\cos(t) e^{it}}{4} \right) dt \\
&= -\frac{t}{8} + \frac{ie^{2it}}{16}
\end{aligned}$$

Now that all the U_i functions have been determined, the particular solution is found from

$$x_p = U_1x_1 + U_2x_2 + U_3x_3 + U_4x_4$$

Hence

$$\begin{aligned}
 x_p &= \left(- \left(\int - \frac{\cos(t) e^{-it} (-i + t)}{4} dt \right) \right) (e^{it}) \\
 &+ \left(\int - \frac{\cos(t) e^{-it}}{4} dt \right) (e^{it}t) \\
 &+ \left(\frac{t^2}{16} + \frac{it}{8} - \frac{i(3i + 2t) e^{2it}}{32} \right) (e^{-it}) \\
 &+ \left(-\frac{t}{8} + \frac{ie^{2it}}{16} \right) (e^{-it}t)
 \end{aligned}$$

Therefore the particular solution is

$$x_p = \frac{\cos(t)(-4t^2 + 2it + 5)}{32} - \frac{\sin(t)(i - 6t)}{32}$$

Which simplifies to

$$x_p = \frac{\cos(t)(-4t^2 + 2it + 5)}{32} - \frac{\sin(t)(i - 6t)}{32}$$

Therefore the general solution is

$$\begin{aligned}
 x &= x_h + x_p \\
 &= (e^{it}c_1 + te^{it}c_2 + e^{-it}c_3 + te^{-it}c_4) + \left(\frac{\cos(t)(-4t^2 + 2it + 5)}{32} - \frac{\sin(t)(i - 6t)}{32} \right)
 \end{aligned}$$

Which simplifies to

$$x = (c_2t + c_1)e^{it} + (c_4t + c_3)e^{-it} + \frac{\cos(t)(-4t^2 + 2it + 5)}{32} - \frac{\sin(t)(i - 6t)}{32}$$

Summary

The solution(s) found are the following

$$x = (c_2t + c_1)e^{it} + (c_4t + c_3)e^{-it} + \frac{\cos(t)(-4t^2 + 2it + 5)}{32} - \frac{\sin(t)(i - 6t)}{32} \quad (1)$$

Verification of solutions

$$x = (c_2t + c_1)e^{it} + (c_4t + c_3)e^{-it} + \frac{\cos(t)(-4t^2 + 2it + 5)}{32} - \frac{\sin(t)(i - 6t)}{32}$$

Verified OK.

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 4; linear nonhomogeneous with symmetry [0,1]  
trying high order linear exact nonhomogeneous  
trying differential order: 4; missing the dependent variable  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 33

```
dsolve(diff(x(t),t$4)+2*diff(x(t),t$2)+x(t)=cos(t),x(t), singsol=all)
```

$$x(t) = \frac{(8c_3t - t^2 + 8c_1 + 2) \cos(t)}{8} + \left(\left(c_4 + \frac{3}{8} \right) t + c_2 \right) \sin(t)$$

✓ Solution by Mathematica

Time used: 0.071 (sec). Leaf size: 43

```
DSolve[x''''[t]+2*x''[t]+x[t]==Cos[t],x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow \left(-\frac{t^2}{8} + c_2t + \frac{5}{16} + c_1 \right) \cos(t) + \frac{1}{4}(t + 4c_4t + 4c_3) \sin(t)$$

2.32 problem Problem 43

- 2.32.1 Solving as second order change of variable on x method 2 ode . 787
- 2.32.2 Solving as second order change of variable on x method 1 ode . 793
- 2.32.3 Solving using Kovacic algorithm 797

Internal problem ID [12195]

Internal file name [OUTPUT/10847_Thursday_September_21_2023_05_47_58_AM_83090191/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 43.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_change_of_variable_on_x_method_1", "second_order_change_of_variable_on_x_method_2"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$(x + 1)^2 y'' + (x + 1) y' + y = 2 \cos(\ln(x + 1))$$

2.32.1 Solving as second order change of variable on x method 2 ode

This is second order non-homogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$(x + 1)^2 y'' + (x + 1) y' + y = 0$$

In normal form the ode

$$(x + 1)^2 y'' + (x + 1) y' + y = 0 \tag{1}$$

Becomes

$$y'' + p(x)y' + q(x)y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x+1}$$
$$q(x) = \frac{1}{(x+1)^2}$$

Applying change of variables $\tau = g(x)$ to (2) gives

$$\frac{d^2}{d\tau^2}y(\tau) + p_1\left(\frac{d}{d\tau}y(\tau)\right) + q_1y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $p_1 = 0$. Eq (4) simplifies to

$$\tau''(x) + p(x)\tau'(x) = 0$$

This ode is solved resulting in

$$\begin{aligned} \tau &= \int e^{-(\int p(x)dx)} dx \\ &= \int e^{-(\int \frac{1}{x+1} dx)} dx \\ &= \int e^{-\ln(x+1)} dx \\ &= \int \frac{1}{x+1} dx \\ &= \ln(x+1) \end{aligned} \quad (6)$$

Using (6) to evaluate q_1 from (5) gives

$$\begin{aligned} q_1(\tau) &= \frac{q(x)}{\tau'(x)^2} \\ &= \frac{1}{(x+1)^2} \\ &= \frac{1}{(x+1)^2} \\ &= 1 \end{aligned} \quad (7)$$

Substituting the above in (3) and noting that now $p_1 = 0$ results in

$$\begin{aligned}\frac{d^2}{d\tau^2}y(\tau) + q_1y(\tau) &= 0 \\ \frac{d^2}{d\tau^2}y(\tau) + y(\tau) &= 0\end{aligned}$$

The above ode is now solved for $y(\tau)$. This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(\tau) + By'(\tau) + Cy(\tau) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y(\tau) = e^{\lambda\tau}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda\tau} + e^{\lambda\tau} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda\tau}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned}\lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i\end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y(\tau) = e^{\alpha\tau}(c_1 \cos(\beta\tau) + c_2 \sin(\beta\tau))$$

Which becomes

$$y(\tau) = e^0(c_1 \cos(\tau) + c_2 \sin(\tau))$$

Or

$$y(\tau) = c_1 \cos(\tau) + c_2 \sin(\tau)$$

The above solution is now transformed back to y using (6) which results in

$$y = c_1 \cos(\ln(x+1)) + c_2 \sin(\ln(x+1))$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(\ln(x+1)) + c_2 \sin(\ln(x+1))$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(\ln(x+1))$$

$$y_2 = \sin(\ln(x+1))$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(\ln(x+1)) & \sin(\ln(x+1)) \\ \frac{d}{dx}(\cos(\ln(x+1))) & \frac{d}{dx}(\sin(\ln(x+1))) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(\ln(x+1)) & \sin(\ln(x+1)) \\ -\frac{\sin(\ln(x+1))}{x+1} & \frac{\cos(\ln(x+1))}{x+1} \end{vmatrix}$$

Therefore

$$W = (\cos(\ln(x+1))) \left(\frac{\cos(\ln(x+1))}{x+1} \right) - (\sin(\ln(x+1))) \left(-\frac{\sin(\ln(x+1))}{x+1} \right)$$

Which simplifies to

$$W = \frac{\cos(\ln(x+1))^2 + \sin(\ln(x+1))^2}{x+1}$$

Which simplifies to

$$W = \frac{1}{x+1}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{2 \sin(\ln(x+1)) \cos(\ln(x+1))}{x+1} dx$$

Which simplifies to

$$u_1 = - \int \frac{\sin(2 \ln(x+1))}{x+1} dx$$

Hence

$$u_1 = \frac{\cos(2 \ln(x+1))}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{2 \cos(\ln(x+1))^2}{x+1} dx$$

Which simplifies to

$$u_2 = \int \frac{2 \cos(\ln(x+1))^2}{x+1} dx$$

Hence

$$u_2 = \sin(\ln(x+1)) \cos(\ln(x+1)) + \ln(x+1)$$

Which simplifies to

$$u_1 = \frac{\cos(2 \ln(x+1))}{2}$$
$$u_2 = \ln(x+1) + \frac{\sin(2 \ln(x+1))}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\cos(2 \ln(x+1)) \cos(\ln(x+1))}{2} + \left(\ln(x+1) + \frac{\sin(2 \ln(x+1))}{2} \right) \sin(\ln(x+1))$$

Which simplifies to

$$y_p(x) = \frac{\cos(\ln(x+1))}{2} + \sin(\ln(x+1)) \ln(x+1)$$

Therefore the general solution is

$$y = y_h + y_p$$
$$= (c_1 \cos(\ln(x+1)) + c_2 \sin(\ln(x+1))) + \left(\frac{\cos(\ln(x+1))}{2} + \sin(\ln(x+1)) \ln(x+1) \right)$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(\ln(x+1)) + c_2 \sin(\ln(x+1)) + \frac{\cos(\ln(x+1))}{2} + \sin(\ln(x+1)) \ln(x+1)$$

Verification of solutions

$$y = c_1 \cos(\ln(x+1)) + c_2 \sin(\ln(x+1)) + \frac{\cos(\ln(x+1))}{2} + \sin(\ln(x+1)) \ln(x+1)$$

Verified OK.

2.32.2 Solving as second order change of variable on x method 1 ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = (x + 1)^2$, $B = x + 1$, $C = 1$, $f(x) = 2 \cos(\ln(x + 1))$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. Solving for y_h from

$$(x + 1)^2 y'' + (x + 1) y' + y = 0$$

In normal form the ode

$$(x + 1)^2 y'' + (x + 1) y' + y = 0 \quad (1)$$

Becomes

$$y'' + p(x) y' + q(x) y = 0 \quad (2)$$

Where

$$p(x) = \frac{1}{x + 1}$$

$$q(x) = \frac{1}{(x + 1)^2}$$

Applying change of variables $\tau = g(x)$ to (2) results

$$\frac{d^2}{d\tau^2} y(\tau) + p_1 \left(\frac{d}{d\tau} y(\tau) \right) + q_1 y(\tau) = 0 \quad (3)$$

Where τ is the new independent variable, and

$$p_1(\tau) = \frac{\tau''(x) + p(x) \tau'(x)}{\tau'(x)^2} \quad (4)$$

$$q_1(\tau) = \frac{q(x)}{\tau'(x)^2} \quad (5)$$

Let $q_1 = c^2$ where c is some constant. Therefore from (5)

$$\tau' = \frac{1}{c} \sqrt{q}$$

$$= \frac{\sqrt{\frac{1}{(x+1)^2}}}{c} \quad (6)$$

$$\tau'' = -\frac{1}{c \sqrt{\frac{1}{(x+1)^2}} (x + 1)^3}$$

Substituting the above into (4) results in

$$\begin{aligned}
 p_1(\tau) &= \frac{\tau''(x) + p(x)\tau'(x)}{\tau'(x)^2} \\
 &= \frac{-\frac{1}{c\sqrt{\frac{1}{(x+1)^2}}(x+1)^3} + \frac{1}{x+1}\frac{\sqrt{\frac{1}{(x+1)^2}}}{c}}{\left(\frac{\sqrt{\frac{1}{(x+1)^2}}}{c}\right)^2} \\
 &= 0
 \end{aligned}$$

Therefore ode (3) now becomes

$$\begin{aligned}
 y(\tau)'' + p_1y(\tau)' + q_1y(\tau) &= 0 \\
 \frac{d^2}{d\tau^2}y(\tau) + c^2y(\tau) &= 0 \tag{7}
 \end{aligned}$$

The above ode is now solved for $y(\tau)$. Since the ode is now constant coefficients, it can be easily solved to give

$$y(\tau) = c_1 \cos(c\tau) + c_2 \sin(c\tau)$$

Now from (6)

$$\begin{aligned}
 \tau &= \int \frac{1}{c}\sqrt{q} dx \\
 &= \frac{\int \sqrt{\frac{1}{(x+1)^2}} dx}{c} \\
 &= \frac{\sqrt{\frac{1}{(x+1)^2}}(x+1) \ln(x+1)}{c}
 \end{aligned}$$

Substituting the above into the solution obtained gives

$$y = c_1 \cos(\ln(x+1)) + c_2 \sin(\ln(x+1))$$

Now the particular solution to this ODE is found

$$(x+1)^2 y'' + (x+1)y' + y = 2 \cos(\ln(x+1))$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of

parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \quad (1)$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = \cos(\ln(x+1))$$

$$y_2 = \sin(\ln(x+1))$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{y_2 f(x)}{aW(x)} \quad (2)$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \quad (3)$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} \cos(\ln(x+1)) & \sin(\ln(x+1)) \\ \frac{d}{dx}(\cos(\ln(x+1))) & \frac{d}{dx}(\sin(\ln(x+1))) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} \cos(\ln(x+1)) & \sin(\ln(x+1)) \\ -\frac{\sin(\ln(x+1))}{x+1} & \frac{\cos(\ln(x+1))}{x+1} \end{vmatrix}$$

Therefore

$$W = (\cos(\ln(x+1))) \left(\frac{\cos(\ln(x+1))}{x+1} \right) - (\sin(\ln(x+1))) \left(-\frac{\sin(\ln(x+1))}{x+1} \right)$$

Which simplifies to

$$W = \frac{\cos(\ln(x+1))^2 + \sin(\ln(x+1))^2}{x+1}$$

Which simplifies to

$$W = \frac{1}{x+1}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{2 \sin(\ln(x+1)) \cos(\ln(x+1))}{x+1} dx$$

Which simplifies to

$$u_1 = - \int \frac{\sin(2 \ln(x+1))}{x+1} dx$$

Hence

$$u_1 = \frac{\cos(2 \ln(x+1))}{2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{2 \cos(\ln(x+1))^2}{x+1} dx$$

Which simplifies to

$$u_2 = \int \frac{2 \cos(\ln(x+1))^2}{x+1} dx$$

Hence

$$u_2 = \sin(\ln(x+1)) \cos(\ln(x+1)) + \ln(x+1)$$

Which simplifies to

$$u_1 = \frac{\cos(2 \ln(x+1))}{2}$$
$$u_2 = \ln(x+1) + \frac{\sin(2 \ln(x+1))}{2}$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = \frac{\cos(2 \ln(x+1)) \cos(\ln(x+1))}{2} + \left(\ln(x+1) + \frac{\sin(2 \ln(x+1))}{2} \right) \sin(\ln(x+1))$$

Which simplifies to

$$y_p(x) = \frac{\cos(\ln(x+1))}{2} + \sin(\ln(x+1)) \ln(x+1)$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(\ln(x+1)) + c_2 \sin(\ln(x+1))) + \left(\frac{\cos(\ln(x+1))}{2} + \sin(\ln(x+1)) \ln(x+1) \right) \\ &= c_1 \cos(\ln(x+1)) + c_2 \sin(\ln(x+1)) + \frac{\cos(\ln(x+1))}{2} + \sin(\ln(x+1)) \ln(x+1) \end{aligned}$$

Which simplifies to

$$y = \frac{(2c_1 + 1) \cos(\ln(x+1))}{2} + \sin(\ln(x+1)) (c_2 + \ln(x+1))$$

Summary

The solution(s) found are the following

$$y = \frac{(2c_1 + 1) \cos(\ln(x+1))}{2} + \sin(\ln(x+1)) (c_2 + \ln(x+1)) \quad (1)$$

Verification of solutions

$$y = \frac{(2c_1 + 1) \cos(\ln(x+1))}{2} + \sin(\ln(x+1)) (c_2 + \ln(x+1))$$

Verified OK.

2.32.3 Solving using Kovacic algorithm

Writing the ode as

$$(x+1)^2 y'' + (x+1) y' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= (x+1)^2 \\ B &= x+1 \\ C &= 1 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x) \tag{4}$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-5}{4(x+1)^2} \tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= -5 \\ t &= 4(x+1)^2 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = \left(-\frac{5}{4(x+1)^2} \right) z(x) \tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 89: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 2 - 0 \\ &= 2 \end{aligned}$$

The poles of r in eq. (7) and the order of each pole are determined by solving for the roots of $t = 4(x+1)^2$. There is a pole at $x = -1$ of order 2. Since there is no odd order pole larger than 2 and the order at ∞ is 2 then the necessary conditions for case one are met. Since there is a pole of order 2 then necessary conditions for case two are met. Since pole order is not larger than 2 and the order at ∞ is 2 then the necessary conditions for case three are met. Therefore

$$L = [1, 2, 4, 6, 12]$$

Attempting to find a solution using case $n = 1$.

Looking at poles of order 2. The partial fractions decomposition of r is

$$r = -\frac{5}{4(x+1)^2}$$

For the pole at $x = -1$ let b be the coefficient of $\frac{1}{(x+1)^2}$ in the partial fractions decom-

position of r given above. Therefore $b = -\frac{5}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_c &= 0 \\ \alpha_c^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + i \\ \alpha_c^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - i \end{aligned}$$

Since the order of r at ∞ is 2 then $[\sqrt{r}]_\infty = 0$. Let b be the coefficient of $\frac{1}{x^2}$ in the Laurent series expansion of r at ∞ , which can be found by dividing the leading coefficient of s by the leading coefficient of t from

$$r = \frac{s}{t} = -\frac{5}{4(x+1)^2}$$

Since the $\gcd(s, t) = 1$. This gives $b = -\frac{5}{4}$. Hence

$$\begin{aligned} [\sqrt{r}]_\infty &= 0 \\ \alpha_\infty^+ &= \frac{1}{2} + \sqrt{1 + 4b} = \frac{1}{2} + i \\ \alpha_\infty^- &= \frac{1}{2} - \sqrt{1 + 4b} = \frac{1}{2} - i \end{aligned}$$

The following table summarizes the findings so far for poles and for the order of r at ∞ where r is

$$r = -\frac{5}{4(x+1)^2}$$

pole c location	pole order	$[\sqrt{r}]_c$	α_c^+	α_c^-
-1	2	0	$\frac{1}{2} + i$	$\frac{1}{2} - i$

Order of r at ∞	$[\sqrt{r}]_\infty$	α_∞^+	α_∞^-
2	0	$\frac{1}{2} + i$	$\frac{1}{2} - i$

Now that the all $[\sqrt{r}]_c$ and its associated α_c^\pm have been determined for all the poles in the set Γ and $[\sqrt{r}]_\infty$ and its associated α_∞^\pm have also been found, the next step is to determine possible non negative integer d from these using

$$d = \alpha_\infty^{s(\infty)} - \sum_{c \in \Gamma} \alpha_c^{s(c)}$$

Where $s(c)$ is either $+$ or $-$ and $s(\infty)$ is the sign of α_∞^\pm . This is done by trial over all set of families $s = (s(c))_{c \in \Gamma \cup \infty}$ until such d is found to work in finding candidate ω .

Trying $\alpha_{\infty}^{-} = \frac{1}{2} - i$ then

$$\begin{aligned} d &= \alpha_{\infty}^{-} - (\alpha_{c_1}^{-}) \\ &= \frac{1}{2} - i - \left(\frac{1}{2} - i\right) \\ &= 0 \end{aligned}$$

Since d an integer and $d \geq 0$ then it can be used to find ω using

$$\omega = \sum_{c \in \Gamma} \left(s(c)[\sqrt{r}]_c + \frac{\alpha_c^{s(c)}}{x - c} \right) + s(\infty)[\sqrt{r}]_{\infty}$$

The above gives

$$\begin{aligned} \omega &= \left((-)[\sqrt{r}]_{c_1} + \frac{\alpha_{c_1}^{-}}{x - c_1} \right) + (-)[\sqrt{r}]_{\infty} \\ &= \frac{\frac{1}{2} - i}{x + 1} + (-)(0) \\ &= \frac{\frac{1}{2} - i}{x + 1} \\ &= \frac{\frac{1}{2} - i}{x + 1} \end{aligned}$$

Now that ω is determined, the next step is find a corresponding minimal polynomial $p(x)$ of degree $d = 0$ to solve the ode. The polynomial $p(x)$ needs to satisfy the equation

$$p'' + 2\omega p' + (\omega' + \omega^2 - r)p = 0 \quad (1A)$$

Let

$$p(x) = 1 \quad (2A)$$

Substituting the above in eq. (1A) gives

$$\begin{aligned} (0) + 2\left(\frac{\frac{1}{2} - i}{x + 1}\right)(0) + \left(\left(\frac{-\frac{1}{2} + i}{(x + 1)^2}\right) + \left(\frac{\frac{1}{2} - i}{x + 1}\right)^2 - \left(-\frac{5}{4(x + 1)^2}\right)\right) &= 0 \\ 0 &= 0 \end{aligned}$$

The equation is satisfied since both sides are zero. Therefore the first solution to the ode $z'' = rz$ is

$$\begin{aligned} z_1(x) &= pe^{\int \omega dx} \\ &= e^{\int \frac{\frac{1}{2} - i}{x + 1} dx} \\ &= (x + 1)^{\frac{1}{2} - i} \end{aligned}$$

The first solution to the original ode in y is found from

$$\begin{aligned}y_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx} \\&= z_1 e^{-\int \frac{1}{2} \frac{x+1}{(x+1)^2} dx} \\&= z_1 e^{-\frac{\ln(x+1)}{2}} \\&= z_1 \left(\frac{1}{\sqrt{x+1}} \right)\end{aligned}$$

Which simplifies to

$$y_1 = (x+1)^{-i}$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Substituting gives

$$\begin{aligned}y_2 &= y_1 \int \frac{e^{\int -\frac{x+1}{(x+1)^2} dx}}{(y_1)^2} dx \\&= y_1 \int \frac{e^{-\ln(x+1)}}{(y_1)^2} dx \\&= y_1 \left(-\frac{i(x+1)^{2i}}{2} \right)\end{aligned}$$

Therefore the solution is

$$\begin{aligned}y &= c_1 y_1 + c_2 y_2 \\&= c_1 \left((x+1)^{-i} \right) + c_2 \left((x+1)^{-i} \left(-\frac{i(x+1)^{2i}}{2} \right) \right)\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$(x + 1)^2 y'' + (x + 1) y' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1(x + 1)^{-i} - \frac{ic_2(x + 1)^i}{2}$$

The particular solution y_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on x as well. Let

$$y_p(x) = u_1 y_1 + u_2 y_2 \tag{1}$$

Where u_1, u_2 to be determined, and y_1, y_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$y_1 = (x + 1)^{-i}$$

$$y_2 = -\frac{i(x + 1)^i}{2}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = -\int \frac{y_2 f(x)}{aW(x)} \tag{2}$$

$$u_2 = \int \frac{y_1 f(x)}{aW(x)} \tag{3}$$

Where $W(x)$ is the Wronskian and a is the coefficient in front of y'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} (x + 1)^{-i} & -\frac{i(x+1)^i}{2} \\ \frac{d}{dx}((x + 1)^{-i}) & \frac{d}{dx}\left(-\frac{i(x+1)^i}{2}\right) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} (x+1)^{-i} & -\frac{i(x+1)^i}{2} \\ -\frac{i(x+1)^{-i}}{x+1} & \frac{(x+1)^i}{2x+2} \end{vmatrix}$$

Therefore

$$W = ((x+1)^{-i}) \left(\frac{(x+1)^i}{2x+2} \right) - \left(-\frac{i(x+1)^i}{2} \right) \left(-\frac{i(x+1)^{-i}}{x+1} \right)$$

Which simplifies to

$$W = \frac{(x+1)^{-i} (x+1)^i}{x+1}$$

Which simplifies to

$$W = \frac{1}{x+1}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{-i(x+1)^i \cos(\ln(x+1))}{x+1} dx$$

Which simplifies to

$$u_1 = - \int -i(x+1)^{-1+i} \cos(\ln(x+1)) dx$$

Hence

$$u_1 = - \left(\int_0^x -i(\alpha+1)^{-1+i} \cos(\ln(\alpha+1)) d\alpha \right)$$

And Eq. (3) becomes

$$u_2 = \int \frac{2(x+1)^{-i} \cos(\ln(x+1))}{x+1} dx$$

Which simplifies to

$$u_2 = \int 2(x+1)^{-1-i} \cos(\ln(x+1)) dx$$

Hence

$$u_2 = \int_0^x 2(\alpha + 1)^{-1-i} \cos(\ln(\alpha + 1)) d\alpha$$

Therefore the particular solution, from equation (1) is

$$y_p(x) = -\left(\int_0^x -i(\alpha + 1)^{-1+i} \cos(\ln(\alpha + 1)) d\alpha\right) (x + 1)^{-i} - \frac{i\left(\int_0^x 2(\alpha + 1)^{-1-i} \cos(\ln(\alpha + 1)) d\alpha\right) (x + 1)^i}{2}$$

Which simplifies to

$$y_p(x) = -i\left(\left(\int_0^x (\alpha + 1)^{-1-i} \cos(\ln(\alpha + 1)) d\alpha\right) (x + 1)^i - \left(\int_0^x (\alpha + 1)^{-1+i} \cos(\ln(\alpha + 1)) d\alpha\right) (x + 1)^{-i}\right)$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1(x + 1)^{-i} - \frac{ic_2(x + 1)^i}{2}\right) + \left(-i\left(\left(\int_0^x (\alpha + 1)^{-1-i} \cos(\ln(\alpha + 1)) d\alpha\right) (x + 1)^i - \left(\int_0^x (\alpha + 1)^{-1+i} \cos(\ln(\alpha + 1)) d\alpha\right) (x + 1)^{-i}\right)\right)$$

Summary

The solution(s) found are the following

$$y = c_1(x + 1)^{-i} - \frac{ic_2(x + 1)^i}{2} - i\left(\left(\int_0^x (\alpha + 1)^{-1-i} \cos(\ln(\alpha + 1)) d\alpha\right) (x + 1)^i - \left(\int_0^x (\alpha + 1)^{-1+i} \cos(\ln(\alpha + 1)) d\alpha\right) (x + 1)^{-i}\right) \quad (1)$$

Verification of solutions

$$y = c_1(x + 1)^{-i} - \frac{ic_2(x + 1)^i}{2} - i\left(\left(\int_0^x (\alpha + 1)^{-1-i} \cos(\ln(\alpha + 1)) d\alpha\right) (x + 1)^i - \left(\int_0^x (\alpha + 1)^{-1+i} \cos(\ln(\alpha + 1)) d\alpha\right) (x + 1)^{-i}\right)$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    checking if the LODE is of Euler type  
    <- LODE of Euler type successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 24

```
dsolve((1+x)^2*diff(y(x),x$2)+(1+x)*diff(y(x),x)+y(x)=2*cos(ln(1+x)),y(x), singsol=all)
```

$$y(x) = (c_2 + \ln(1+x)) \sin(\ln(1+x)) + \cos(\ln(1+x)) c_1$$

✓ Solution by Mathematica

Time used: 0.192 (sec). Leaf size: 31

```
DSolve[(1+x)^2*y''[x]+(1+x)*y'[x]+y[x]==2*Cos[Log[1+x]],y[x],x,IncludeSingularSolutions -> T
```

$$y(x) \rightarrow \left(\frac{1}{2} + c_1\right) \cos(\log(x+1)) + (\log(x+1) + c_2) \sin(\log(x+1))$$

2.33 problem Problem 47

Internal problem ID [12196]

Internal file name [OUTPUT/10848_Thursday_September_21_2023_05_48_01_AM_14234251/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 47.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "reduction_of_order", "second_order_change_of_variable_on_y_method_2", "second_order_ode_non_constant_coeff_transformation_on_B"

Maple gives the following as the ode type

```
[[_2nd_order, _with_linear_symmetries], [_2nd_order, _linear, `
  _with_symmetry_[0,F(x)]`]]
```

$$x^3 y'' - y'x + y = 0$$

Given that one solution of the ode is

$$y_1 = x$$

Given one basis solution $y_1(x)$, then the second basis solution is given by

$$y_2(x) = y_1 \left(\int \frac{e^{-\int p dx}}{y_1^2} dx \right)$$

Where $p(x)$ is the coefficient of y' when the ode is written in the normal form

$$y'' + p(x)y' + q(x)y = f(x)$$

Looking at the ode to solve shows that

$$p(x) = -\frac{1}{x^2}$$

Therefore

$$y_2(x) = x \left(\int \frac{e^{-\left(\int -\frac{1}{x^2} dx\right)}}{x^2} dx \right)$$

$$y_2(x) = x \int \frac{e^{-\frac{1}{x}}}{x^2} dx$$

$$y_2(x) = x \left(\int \frac{e^{-\frac{1}{x}}}{x^2} dx \right)$$

$$y_2(x) = e^{-\frac{1}{x}} x$$

Hence the solution is

$$\begin{aligned} y &= c_1 y_1(x) + c_2 y_2(x) \\ &= c_1 x + c_2 e^{-\frac{1}{x}} x \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 x + c_2 e^{-\frac{1}{x}} x \quad (1)$$

Verification of solutions

$$y = c_1 x + c_2 e^{-\frac{1}{x}} x$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
checking if the LODE has constant coefficients  
checking if the LODE is of Euler type  
trying a symmetry of the form [xi=0, eta=F(x)]  
<- linear_1 successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 16

```
dsolve([x^3*diff(y(x),x$2)-x*diff(y(x),x)+y(x)=0,x],singsol=all)
```

$$y(x) = \left(e^{-\frac{1}{x}} c_1 + c_2 \right) x$$

✓ Solution by Mathematica

Time used: 0.095 (sec). Leaf size: 20

```
DSolve[x^3*y''[x]-x*y'[x]+y[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow x(c_2 e^{-1/x} + c_1)$$

2.34 problem Problem 49

2.34.1 Maple step by step solution 813

Internal problem ID [12197]

Internal file name [OUTPUT/10849_Thursday_September_21_2023_05_48_02_AM_3993535/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 49.

ODE order: 4.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _linear , _nonhomogeneous]]
```

$$x'''' + x = t^3$$

This is higher order nonhomogeneous ODE. Let the solution be

$$x = x_h + x_p$$

Where x_h is the solution to the homogeneous ODE And x_p is a particular solution to the nonhomogeneous ODE. x_h is the solution to

$$x'''' + x = 0$$

The characteristic equation is

$$\lambda^4 + 1 = 0$$

The roots of the above equation are

$$\begin{aligned}\lambda_1 &= \frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2} \\ \lambda_2 &= -\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2} \\ \lambda_3 &= -\frac{\sqrt{2}}{2} - \frac{i\sqrt{2}}{2} \\ \lambda_4 &= \frac{\sqrt{2}}{2} - \frac{i\sqrt{2}}{2}\end{aligned}$$

Therefore the homogeneous solution is

$$x_h(t) = e^{\left(\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right)t} c_1 + e^{\left(-\frac{\sqrt{2}}{2} - \frac{i\sqrt{2}}{2}\right)t} c_2 + e^{\left(\frac{\sqrt{2}}{2} - \frac{i\sqrt{2}}{2}\right)t} c_3 + e^{\left(-\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right)t} c_4$$

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned}x_1 &= e^{\left(\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right)t} \\ x_2 &= e^{\left(-\frac{\sqrt{2}}{2} - \frac{i\sqrt{2}}{2}\right)t} \\ x_3 &= e^{\left(\frac{\sqrt{2}}{2} - \frac{i\sqrt{2}}{2}\right)t} \\ x_4 &= e^{\left(-\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right)t}\end{aligned}$$

Now the particular solution to the given ODE is found

$$x'''' + x = t^3$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$t^3$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{1, t, t^2, t^3\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^{\left(-\frac{\sqrt{2}}{2} - \frac{i\sqrt{2}}{2}\right)t}, e^{\left(-\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right)t}, e^{\left(\frac{\sqrt{2}}{2} - \frac{i\sqrt{2}}{2}\right)t}, e^{\left(\frac{\sqrt{2}}{2} + \frac{i\sqrt{2}}{2}\right)t} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$x_p = A_4 t^3 + A_3 t^2 + A_2 t + A_1$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution x_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$A_4 t^3 + A_3 t^2 + A_2 t + A_1 = t^3$$

Solving for the unknowns by comparing coefficients results in

$$[A_1 = 0, A_2 = 0, A_3 = 0, A_4 = 1]$$

Substituting the above back in the above trial solution x_p , gives the particular solution

$$x_p = t^3$$

Therefore the general solution is

$$\begin{aligned} x &= x_h + x_p \\ &= \left(e^{\left(\frac{\sqrt{2} + i\sqrt{2}}{2}\right)t} c_1 + e^{\left(-\frac{\sqrt{2} - i\sqrt{2}}{2}\right)t} c_2 + e^{\left(\frac{\sqrt{2} - i\sqrt{2}}{2}\right)t} c_3 + e^{\left(-\frac{\sqrt{2} + i\sqrt{2}}{2}\right)t} c_4 \right) + (t^3) \end{aligned}$$

Which simplifies to

$$x = e^{\left(\frac{1}{2} + \frac{i}{2}\right)\sqrt{2}t} c_1 + e^{\left(-\frac{1}{2} - \frac{i}{2}\right)\sqrt{2}t} c_2 + e^{\left(\frac{1}{2} - \frac{i}{2}\right)\sqrt{2}t} c_3 + e^{\left(-\frac{1}{2} + \frac{i}{2}\right)\sqrt{2}t} c_4 + t^3$$

Summary

The solution(s) found are the following

$$x = e^{\left(\frac{1}{2} + \frac{i}{2}\right)\sqrt{2}t} c_1 + e^{\left(-\frac{1}{2} - \frac{i}{2}\right)\sqrt{2}t} c_2 + e^{\left(\frac{1}{2} - \frac{i}{2}\right)\sqrt{2}t} c_3 + e^{\left(-\frac{1}{2} + \frac{i}{2}\right)\sqrt{2}t} c_4 + t^3 \quad (1)$$

Verification of solutions

$$x = e^{\left(\frac{1}{2} + \frac{i}{2}\right)\sqrt{2}t} c_1 + e^{\left(-\frac{1}{2} - \frac{i}{2}\right)\sqrt{2}t} c_2 + e^{\left(\frac{1}{2} - \frac{i}{2}\right)\sqrt{2}t} c_3 + e^{\left(-\frac{1}{2} + \frac{i}{2}\right)\sqrt{2}t} c_4 + t^3$$

Verified OK.

2.34.1 Maple step by step solution

Let's solve

$$x'''' + x = t^3$$

- Highest derivative means the order of the ODE is 4

$$x''''$$

- Convert linear ODE into a system of first order ODEs

- Define new variable $x_1(t)$

$$x_1(t) = x$$

- Define new variable $x_2(t)$

$$x_2(t) = x'$$

- Define new variable $x_3(t)$

$$x_3(t) = x''$$

- Define new variable $x_4(t)$

$$x_4(t) = x'''$$

- Isolate for $x_4'(t)$ using original ODE

$$x_4'(t) = t^3 - x_1(t)$$

Convert linear ODE into a system of first order ODEs

$$[x_2(t) = x_1'(t), x_3(t) = x_2'(t), x_4(t) = x_3'(t), x_4'(t) = t^3 - x_1(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x_1(t) \\ x_2(t) \\ x_3(t) \\ x_4(t) \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \\ 0 \\ t^3 \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(t) = \begin{bmatrix} 0 \\ 0 \\ 0 \\ t^3 \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[\begin{array}{c} -\frac{\sqrt{2}}{2} - \frac{1\sqrt{2}}{2}, \\ \frac{1}{\left(-\frac{\sqrt{2}}{2} - \frac{1\sqrt{2}}{2}\right)^3} \\ \frac{1}{\left(-\frac{\sqrt{2}}{2} - \frac{1\sqrt{2}}{2}\right)^2} \\ \frac{1}{-\frac{\sqrt{2}}{2} - \frac{1\sqrt{2}}{2}} \\ 1 \end{array} \right], \left[\begin{array}{c} -\frac{\sqrt{2}}{2} + \frac{1\sqrt{2}}{2}, \\ \frac{1}{\left(-\frac{\sqrt{2}}{2} + \frac{1\sqrt{2}}{2}\right)^3} \\ \frac{1}{\left(-\frac{\sqrt{2}}{2} + \frac{1\sqrt{2}}{2}\right)^2} \\ \frac{1}{-\frac{\sqrt{2}}{2} + \frac{1\sqrt{2}}{2}} \\ 1 \end{array} \right], \left[\begin{array}{c} \frac{\sqrt{2}}{2} - \frac{1\sqrt{2}}{2}, \\ \frac{1}{\left(\frac{\sqrt{2}}{2} - \frac{1\sqrt{2}}{2}\right)^3} \\ \frac{1}{\left(\frac{\sqrt{2}}{2} - \frac{1\sqrt{2}}{2}\right)^2} \\ \frac{1}{\frac{\sqrt{2}}{2} - \frac{1\sqrt{2}}{2}} \\ 1 \end{array} \right] \right]$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[\begin{array}{c} -\frac{\sqrt{2}}{2} - \frac{1\sqrt{2}}{2}, \\ \frac{1}{\left(-\frac{\sqrt{2}}{2} - \frac{1\sqrt{2}}{2}\right)^3} \\ \frac{1}{\left(-\frac{\sqrt{2}}{2} - \frac{1\sqrt{2}}{2}\right)^2} \\ \frac{1}{-\frac{\sqrt{2}}{2} - \frac{1\sqrt{2}}{2}} \\ 1 \end{array} \right]$$

- Solution from eigenpair

$$e^{(-\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2})t} \cdot \begin{bmatrix} \frac{1}{(-\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2})^3} \\ \frac{1}{(-\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2})^2} \\ \frac{1}{-\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{-\frac{\sqrt{2}t}{2}} \cdot \left(\cos\left(\frac{\sqrt{2}t}{2}\right) - I \sin\left(\frac{\sqrt{2}t}{2}\right) \right) \cdot \begin{bmatrix} \frac{1}{(-\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2})^3} \\ \frac{1}{(-\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2})^2} \\ \frac{1}{-\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{-\frac{\sqrt{2}t}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{2}t}{2}\right) - I \sin\left(\frac{\sqrt{2}t}{2}\right)}{(-\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2})^3} \\ \frac{\cos\left(\frac{\sqrt{2}t}{2}\right) - I \sin\left(\frac{\sqrt{2}t}{2}\right)}{(-\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2})^2} \\ \frac{\cos\left(\frac{\sqrt{2}t}{2}\right) - I \sin\left(\frac{\sqrt{2}t}{2}\right)}{-\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}} \\ \cos\left(\frac{\sqrt{2}t}{2}\right) - I \sin\left(\frac{\sqrt{2}t}{2}\right) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{x}_1(t) = e^{-\frac{\sqrt{2}t}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} + \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \\ -\sin\left(\frac{\sqrt{2}t}{2}\right) \\ -\frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} + \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \\ \cos\left(\frac{\sqrt{2}t}{2}\right) \end{bmatrix}, \vec{x}_2(t) = e^{-\frac{\sqrt{2}t}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} - \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \\ -\cos\left(\frac{\sqrt{2}t}{2}\right) \\ \frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} + \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \\ -\sin\left(\frac{\sqrt{2}t}{2}\right) \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\begin{bmatrix} \frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}, & \begin{bmatrix} \frac{1}{\left(\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}\right)^3} \\ \frac{1}{\left(\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}\right)^2} \\ \frac{1}{\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}} \\ 1 \end{bmatrix} \end{bmatrix}$$

- Solution from eigenpair

$$e^{\left(\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}\right)t} \cdot \begin{bmatrix} \frac{1}{\left(\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}\right)^3} \\ \frac{1}{\left(\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}\right)^2} \\ \frac{1}{\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{\frac{\sqrt{2}t}{2}} \cdot \left(\cos\left(\frac{\sqrt{2}t}{2}\right) - I \sin\left(\frac{\sqrt{2}t}{2}\right) \right) \cdot \begin{bmatrix} \frac{1}{\left(\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}\right)^3} \\ \frac{1}{\left(\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}\right)^2} \\ \frac{1}{\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{\frac{\sqrt{2}t}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{2}t}{2}\right) - I \sin\left(\frac{\sqrt{2}t}{2}\right)}{\left(\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}\right)^3} \\ \frac{\cos\left(\frac{\sqrt{2}t}{2}\right) - I \sin\left(\frac{\sqrt{2}t}{2}\right)}{\left(\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}\right)^2} \\ \frac{\cos\left(\frac{\sqrt{2}t}{2}\right) - I \sin\left(\frac{\sqrt{2}t}{2}\right)}{\frac{\sqrt{2}}{2} - \frac{I\sqrt{2}}{2}} \\ \cos\left(\frac{\sqrt{2}t}{2}\right) - I \sin\left(\frac{\sqrt{2}t}{2}\right) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{x}_3(t) = e^{\frac{\sqrt{2}t}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} + \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \\ \sin\left(\frac{\sqrt{2}t}{2}\right) \\ \frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} + \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \\ \cos\left(\frac{\sqrt{2}t}{2}\right) \end{bmatrix}, \vec{x}_4(t) = e^{\frac{\sqrt{2}t}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} + \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \\ \cos\left(\frac{\sqrt{2}t}{2}\right) \\ \frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} - \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \\ -\sin\left(\frac{\sqrt{2}t}{2}\right) \end{bmatrix}$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{x}_p(t)$
 $\vec{x}(t) = c_1\vec{x}_1(t) + c_2\vec{x}_2(t) + c_3\vec{x}_3(t) + c_4\vec{x}_4(t) + \vec{x}_p(t)$

□ Fundamental matrix

- Let $\phi(t)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(t) = \begin{bmatrix} e^{-\frac{\sqrt{2}t}{2}} \left(\frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} + \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \right) & e^{-\frac{\sqrt{2}t}{2}} \left(\frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} - \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \right) & e^{\frac{\sqrt{2}t}{2}} \left(-\frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \right) \\ -e^{-\frac{\sqrt{2}t}{2}} \sin\left(\frac{\sqrt{2}t}{2}\right) & -e^{-\frac{\sqrt{2}t}{2}} \cos\left(\frac{\sqrt{2}t}{2}\right) & e^{\frac{\sqrt{2}t}{2}} \sin\left(\frac{\sqrt{2}t}{2}\right) \\ e^{-\frac{\sqrt{2}t}{2}} \left(-\frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} + \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \right) & e^{-\frac{\sqrt{2}t}{2}} \left(\frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} + \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \right) & e^{\frac{\sqrt{2}t}{2}} \left(\frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \right) \\ e^{-\frac{\sqrt{2}t}{2}} \cos\left(\frac{\sqrt{2}t}{2}\right) & -e^{-\frac{\sqrt{2}t}{2}} \sin\left(\frac{\sqrt{2}t}{2}\right) & e^{\frac{\sqrt{2}t}{2}} \cos\left(\frac{\sqrt{2}t}{2}\right) \end{bmatrix}$$

- The fundamental matrix, $\Phi(t)$ is a normalized version of $\phi(t)$ satisfying $\Phi(0) = I$ where I is the identity matrix
 $\Phi(t) = \phi(t) \cdot \frac{1}{\phi(0)}$

- Substitute the value of $\phi(t)$ and $\phi(0)$

$$\Phi(t) = \begin{bmatrix} e^{-\frac{\sqrt{2}t}{2}} \left(\frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} + \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \right) & e^{-\frac{\sqrt{2}t}{2}} \left(\frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} - \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \right) & e^{\frac{\sqrt{2}t}{2}} \left(-\frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \right) \\ -e^{-\frac{\sqrt{2}t}{2}} \sin\left(\frac{\sqrt{2}t}{2}\right) & -e^{-\frac{\sqrt{2}t}{2}} \cos\left(\frac{\sqrt{2}t}{2}\right) & e^{\frac{\sqrt{2}t}{2}} \sin\left(\frac{\sqrt{2}t}{2}\right) \\ e^{-\frac{\sqrt{2}t}{2}} \left(-\frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} + \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \right) & e^{-\frac{\sqrt{2}t}{2}} \left(\frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} + \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \right) & e^{\frac{\sqrt{2}t}{2}} \left(\frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \right) \\ e^{-\frac{\sqrt{2}t}{2}} \cos\left(\frac{\sqrt{2}t}{2}\right) & -e^{-\frac{\sqrt{2}t}{2}} \sin\left(\frac{\sqrt{2}t}{2}\right) & e^{\frac{\sqrt{2}t}{2}} \cos\left(\frac{\sqrt{2}t}{2}\right) \end{bmatrix}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(t) = \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\left(e^{-\frac{\sqrt{2}t}{2}} + e^{\frac{\sqrt{2}t}{2}}\right)}{2} & -\frac{\left(\left(e^{-\frac{\sqrt{2}t}{2}} - e^{\frac{\sqrt{2}t}{2}}\right)\cos\left(\frac{\sqrt{2}t}{2}\right) - \sin\left(\frac{\sqrt{2}t}{2}\right)\right)\sqrt{2}}{4} \\ -\frac{\left(\left(e^{-\frac{\sqrt{2}t}{2}} - e^{\frac{\sqrt{2}t}{2}}\right)\cos\left(\frac{\sqrt{2}t}{2}\right) + \sin\left(\frac{\sqrt{2}t}{2}\right)\left(e^{-\frac{\sqrt{2}t}{2}} + e^{\frac{\sqrt{2}t}{2}}\right)\right)\sqrt{2}}{4} & \frac{\cos\left(\frac{\sqrt{2}t}{2}\right)\left(e^{-\frac{\sqrt{2}t}{2}} + e^{\frac{\sqrt{2}t}{2}}\right)}{2} \\ \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\left(e^{-\frac{\sqrt{2}t}{2}} - e^{\frac{\sqrt{2}t}{2}}\right)}{2} & -\frac{\left(\left(e^{-\frac{\sqrt{2}t}{2}} - e^{\frac{\sqrt{2}t}{2}}\right)\cos\left(\frac{\sqrt{2}t}{2}\right) + \sin\left(\frac{\sqrt{2}t}{2}\right)\right)\sqrt{2}}{4} \\ \frac{\left(\left(e^{-\frac{\sqrt{2}t}{2}} - e^{\frac{\sqrt{2}t}{2}}\right)\cos\left(\frac{\sqrt{2}t}{2}\right) - \sin\left(\frac{\sqrt{2}t}{2}\right)\left(e^{-\frac{\sqrt{2}t}{2}} + e^{\frac{\sqrt{2}t}{2}}\right)\right)\sqrt{2}}{4} & \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\left(e^{-\frac{\sqrt{2}t}{2}} - e^{\frac{\sqrt{2}t}{2}}\right)}{2} \end{bmatrix}$$

- Find a particular solution of the system of ODEs using variation of parameters
- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(t)$ and solve for $\vec{v}(t)$

$$\vec{x}_p(t) = \Phi(t) \cdot \vec{v}(t)$$
 - Take the derivative of the particular solution
$$\vec{x}'_p(t) = \Phi'(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t)$$
 - Substitute particular solution and its derivative into the system of ODEs
$$\Phi'(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t) = A \cdot \Phi(t) \cdot \vec{v}(t) + \vec{f}(t)$$
 - The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(t)$

$$A \cdot \Phi(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t) = A \cdot \Phi(t) \cdot \vec{v}(t) + \vec{f}(t)$$
 - Cancel like terms
$$\Phi(t) \cdot \vec{v}'(t) = \vec{f}(t)$$
 - Multiply by the inverse of the fundamental matrix
$$\vec{v}'(t) = \frac{1}{\Phi(t)} \cdot \vec{f}(t)$$
 - Integrate to solve for $\vec{v}(t)$

$$\vec{v}(t) = \int_0^t \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$
 - Plug $\vec{v}(t)$ into the equation for the particular solution
$$\vec{x}_p(t) = \Phi(t) \cdot \left(\int_0^t \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$
 - Plug in the fundamental matrix and the forcing function and compute

$$\vec{x}_p(t) = \begin{bmatrix} -\frac{3\sqrt{2}\left(e^{-\frac{\sqrt{2}t}{2}} - e^{\frac{\sqrt{2}t}{2}}\right)\cos\left(\frac{\sqrt{2}t}{2}\right)}{2} + t^3 - \frac{3e^{-\frac{\sqrt{2}t}{2}}\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} - \frac{3e^{\frac{\sqrt{2}t}{2}}\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \\ \left(3e^{-\frac{\sqrt{2}t}{2}} - 3e^{\frac{\sqrt{2}t}{2}}\right)\sin\left(\frac{\sqrt{2}t}{2}\right) + 3t^2 \\ \frac{3\sqrt{2}\left(e^{-\frac{\sqrt{2}t}{2}} - e^{\frac{\sqrt{2}t}{2}}\right)\cos\left(\frac{\sqrt{2}t}{2}\right)}{2} - \frac{3e^{-\frac{\sqrt{2}t}{2}}\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} - \frac{3e^{\frac{\sqrt{2}t}{2}}\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} + 6t \\ 6 + \left(-3e^{-\frac{\sqrt{2}t}{2}} - 3e^{\frac{\sqrt{2}t}{2}}\right)\cos\left(\frac{\sqrt{2}t}{2}\right) \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{x}(t) = c_1\vec{x}_1(t) + c_2\vec{x}_2(t) + c_3\vec{x}_3(t) + c_4\vec{x}_4(t) + \begin{bmatrix} -\frac{3\sqrt{2}\left(e^{-\frac{\sqrt{2}t}{2}} - e^{\frac{\sqrt{2}t}{2}}\right)\cos\left(\frac{\sqrt{2}t}{2}\right)}{2} + t^3 - \frac{3e^{-\frac{\sqrt{2}t}{2}}\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \\ \left(3e^{-\frac{\sqrt{2}t}{2}} - 3e^{\frac{\sqrt{2}t}{2}}\right)\sin\left(\frac{\sqrt{2}t}{2}\right) \\ \frac{3\sqrt{2}\left(e^{-\frac{\sqrt{2}t}{2}} - e^{\frac{\sqrt{2}t}{2}}\right)\cos\left(\frac{\sqrt{2}t}{2}\right)}{2} - \frac{3e^{-\frac{\sqrt{2}t}{2}}\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}}{2} \\ 6 + \left(-3e^{-\frac{\sqrt{2}t}{2}} - 3e^{\frac{\sqrt{2}t}{2}}\right)\cos\left(\frac{\sqrt{2}t}{2}\right) \end{bmatrix}$$

- First component of the vector is the solution to the ODE

$$x = \frac{\sqrt{2}\left((c_1+c_2-3)e^{-\frac{\sqrt{2}t}{2}} - e^{\frac{\sqrt{2}t}{2}}(c_3-c_4-3)\right)\cos\left(\frac{\sqrt{2}t}{2}\right)}{2} + \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}(c_1-c_2-3)e^{-\frac{\sqrt{2}t}{2}}}{2} + \frac{\sin\left(\frac{\sqrt{2}t}{2}\right)\sqrt{2}(c_3+c_4-3)e^{\frac{\sqrt{2}t}{2}}}{2}$$

Maple trace

```

`Methods for high order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 4; linear nonhomogeneous with symmetry [0,1]
trying high order linear exact nonhomogeneous
trying differential order: 4; missing the dependent variable
checking if the LODE has constant coefficients
<- constant coefficients successful`

```

✓ Solution by Maple

Time used: 0.015 (sec). Leaf size: 67

```
dsolve(diff(x(t),t$4)+x(t)=t^3,x(t), singsol=all)
```

$$x(t) = \left(c_2 e^{-\frac{\sqrt{2}t}{2}} + c_4 e^{\frac{\sqrt{2}t}{2}} \right) \sin\left(\frac{\sqrt{2}t}{2}\right) + t^3 + c_1 e^{-\frac{\sqrt{2}t}{2}} \cos\left(\frac{\sqrt{2}t}{2}\right) + c_3 e^{\frac{\sqrt{2}t}{2}} \cos\left(\frac{\sqrt{2}t}{2}\right)$$

✓ Solution by Mathematica

Time used: 0.008 (sec). Leaf size: 78

```
DSolve[x''''[t]+x[t]==t^3,x[t],t,IncludeSingularSolutions -> True]
```

$$x(t) \rightarrow e^{-\frac{t}{\sqrt{2}}} \left(e^{\frac{t}{\sqrt{2}}} t^3 + (c_1 e^{\sqrt{2}t} + c_2) \cos\left(\frac{t}{\sqrt{2}}\right) + (c_4 e^{\sqrt{2}t} + c_3) \sin\left(\frac{t}{\sqrt{2}}\right) \right)$$

2.35 problem Problem 50

2.35.1 Solving as second order ode missing y ode	821
2.35.2 Solving using Kovacic algorithm	825
2.35.3 Maple step by step solution	828

Internal problem ID [12198]

Internal file name [OUTPUT/10850_Thursday_September_21_2023_05_48_02_AM_89497818/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 50.

ODE order: 2.

ODE degree: 3.

The type(s) of ODE detected by this program : "**second_order_ode_high_degree**", "**second_order_ode_missing_y**"

Maple gives the following as the ode type

```
[[_2nd_order , _quadrature]]
```

$$y''' + y'' = x - 1$$

2.35.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$(p'(x)^2 + 1)p'(x) - x + 1 = 0$$

Which is now solve for $p(x)$ as first order ode. Solving the given ode for $p'(x)$ results in 3 differential equations to solve. Each one of these will generate a solution. The

equations generated are

$$p'(x) = \frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}}{6} - \frac{2}{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} \quad (1)$$

$$p'(x) = -\frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}}{12} + \frac{1}{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} + \frac{i\sqrt{3}}{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} \quad (2)$$

$$p'(x) = -\frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}}{12} + \frac{1}{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} - \frac{i\sqrt{3}}{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} \quad (3)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\begin{aligned} p(x) &= \int \frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} - 12}{6(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx \\ &= \int \frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} - 12}{6(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx + c_1 \end{aligned}$$

Solving equation (2)

Integrating both sides gives

$$\begin{aligned} p(x) &= \int \frac{i\sqrt{3}(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} - (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}}{12(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx \\ &= \int \frac{i\sqrt{3}(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} - (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}}{12(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx \end{aligned}$$

Solving equation (3)

Integrating both sides gives

$$\begin{aligned}
 p(x) &= \int -\frac{i\sqrt{3}(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} + (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}}}{12(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx \\
 &= \int -\frac{i\sqrt{3}(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} + (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}}}{12(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx
 \end{aligned}$$

For solution (1) found earlier, since $p = y'$ then the new first order ode to solve is

$$y' = \int \frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} - 12}{6(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx + c_1$$

Integrating both sides gives

$$\begin{aligned}
 y &= \int \int \frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} - 12}{6(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx + c_1 dx \\
 &= \int \left(\int \frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} - 12}{6(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx + c_1 \right) dx + c_4
 \end{aligned}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = \int \frac{i\sqrt{3}(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} - (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}}}{12(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx$$

Integrating both sides gives

$$\begin{aligned}
 y &= \int \int \frac{i\sqrt{3}(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} - (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}}}{12(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx \\
 &= \int \left(\int \frac{i\sqrt{3}(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} - (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}}}{12(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx \right) dx
 \end{aligned}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = \int -\frac{i\sqrt{3}(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} + (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}}}{12(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx$$

Integrating both sides gives

$$\begin{aligned}
 y &= \int \int -\frac{i\sqrt{3}(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} + (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}}}{12(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx \\
 &= \int \left(\int -\frac{i\sqrt{3}(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} + (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}}}{12(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx \right) dx
 \end{aligned}$$

Summary

The solution(s) found are the following

$$y = \int \left(\int \frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} - 12}{6(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx + c_1 \right) dx + c_4 \quad (1)$$

$$y \quad (2)$$

$$\begin{aligned}
 &= \int \left(\int \frac{i\sqrt{3}(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} - (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}}}{12(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} \right. \\
 &\quad \left. + c_2 \right) dx + c_5
 \end{aligned}$$

$$y = \int \left(\int \quad (3)$$

$$\begin{aligned}
 &\frac{i\sqrt{3}(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} + (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}}}{12(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} \\
 &\quad \left. + c_3 \right) dx + c_6
 \end{aligned}$$

Verification of solutions

$$y = \int \left(\int \frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} - 12}{6(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx + c_1 \right) dx + c_4$$

Verified OK.

$$y = \int \left(\int \frac{i\sqrt{3}(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} - (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}}}{12(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} + c_2 \right) dx + c_5$$

Verified OK.

$$y = \int \left(\int \frac{i\sqrt{3}(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} + (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}}}{12(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} + c_3 \right) dx + c_6$$

Verified OK.

2.35.2 Solving using Kovacic algorithm

Solving for y'' from the ode gives

$$y'' = \frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}}{6} - \frac{2}{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} \quad (1)$$

$$\begin{aligned}
y'' = & -\frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}}{12} \\
& + \frac{1}{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} \\
& - \frac{i\sqrt{3} \left(\frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}}{6} + \frac{2}{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} \right)}{2}
\end{aligned} \tag{2}$$

$$\begin{aligned}
y'' = & -\frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}}{12} \\
& + \frac{1}{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} \\
& + \frac{i\sqrt{3} \left(\frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}}{6} + \frac{2}{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} \right)}{2}
\end{aligned} \tag{3}$$

Now each ode is solved. Integrating once gives

$$y' = \int \frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} - 12}{6(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx + c_1$$

Integrating again gives

$$y = \int \left(\int \frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} - 12}{6(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx \right) dx + c_1x + c_2$$

Integrating once gives

$$y' = \int -\frac{(1 + i\sqrt{3})(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} - 12}{12(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx + c_3$$

Integrating again gives

$$y = \int \left(\int -\frac{(1 + i\sqrt{3})(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} - 12}{12(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx \right) dx + c_3x + c_4$$

Integrating once gives

$$y' = \int \frac{(i\sqrt{3} - 1) (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} + 12}{12 (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx + c_5$$

Integrating again gives

$$y = \int \left(\int \frac{(i\sqrt{3} - 1) (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} + 12}{12 (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx \right) dx + c_5x + c_6$$

Summary

The solution(s) found are the following

$$y = \int \int \frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} - 12}{6 (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx dx + c_1x + c_2 \quad (1)$$

$$y = \int \int -\frac{(1 + i\sqrt{3}) (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} - 12}{12 (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx dx + c_3x + c_4 \quad (2)$$

$$y = \int \int \frac{(i\sqrt{3} - 1) (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} + 12}{12 (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx dx + c_5x + c_6 \quad (3)$$

Verification of solutions

$$y = \int \int \frac{(-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} - 12}{6 (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx dx + c_1x + c_2$$

Verified OK.

$$y = \int \int -\frac{(1 + i\sqrt{3}) (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} - 12}{12 (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx dx + c_3x + c_4$$

Verified OK.

$$y = \int \int \frac{(i\sqrt{3} - 1) (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{2}{3}} + 12i\sqrt{3} + 12}{12 (-108 + 108x + 12\sqrt{81x^2 - 162x + 93})^{\frac{1}{3}}} dx dx + c_5x + c_6$$

Verified OK.

2.35.3 Maple step by step solution

Let's solve

$$(y''^2 + 1) y'' = x - 1$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Isolate 2nd derivative

$$y'' = \frac{(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{2}{3}}-12}{6(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{1}{3}}}$$

- Characteristic polynomial of homogeneous ODE

$$r^2 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{0})}{2}$$

- Roots of the characteristic polynomial

$$r = 0$$

- 1st solution of the homogeneous ODE

$$y_1(x) = 1$$

- Repeated root, multiply $y_1(x)$ by x to ensure linear independence

$$y_2(x) = x$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 + c_2 x + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right) \right], f(x) = \frac{(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{2}{3}}-12}{6(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{1}{3}}}$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\left(\int \frac{x \left((-108+108x+12\sqrt{81x^2-162x+93})^{\frac{2}{3}} - 12 \right)}{(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{1}{3}}} dx\right)}{6} + \frac{x \left(\int \frac{(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{2}{3}} - 12}{(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{1}{3}}} dx\right)}{6}$$

- Compute integrals

$$y_p(x) = -\frac{\left(\int \frac{x \left((-108+108x+12\sqrt{81x^2-162x+93})^{\frac{2}{3}} - 12 \right)}{(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{1}{3}}} dx\right)}{6} + \frac{x \left(\int \frac{(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{2}{3}} - 12}{(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{1}{3}}} dx\right)}{6}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 + c_2x - \frac{\left(\int \frac{x \left((-108+108x+12\sqrt{81x^2-162x+93})^{\frac{2}{3}} - 12 \right)}{(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{1}{3}}} dx\right)}{6} + \frac{x \left(\int \frac{(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{2}{3}} - 12}{(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{1}{3}}} dx\right)}{6}$$

Maple trace

```
`Methods for second order ODEs:
Successful isolation of  $d^2y/dx^2$ : 3 solutions were found. Trying to solve each resulting ODE
  *** Sublevel 2 ***
  Methods for second order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  <- quadrature successful
-----
* Tackling next ODE.
  *** Sublevel 2 ***
  Methods for second order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  <- quadrature successful
-----
* Tackling next ODE.
  *** Sublevel 2 ***
  Methods for second order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  <- quadrature successful`
```

✓ Solution by Maple

Time used: 0.031 (sec). Leaf size: 226

```
dsolve(diff(y(x),x$2)^3+diff(y(x),x$2)+1=x,y(x), singsol=all)
```

$$y(x) = \frac{\left(\int \int \frac{(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{2}{3}}-12}{(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{1}{3}}} dx dx \right)}{6} + c_1x + c_2$$
$$y(x) = \frac{\left(\int \int \frac{i\sqrt{3}(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{2}{3}}+12i\sqrt{3}+(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{2}{3}}-12}{(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{1}{3}}} dx dx \right)}{12} + c_1x + c_2$$
$$y(x) = \frac{\left(\int \int \frac{(i\sqrt{3}-1)(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{2}{3}}+12i\sqrt{3}+12}{(-108+108x+12\sqrt{81x^2-162x+93})^{\frac{1}{3}}} dx dx \right)}{12} + c_1x + c_2$$

✗ Solution by Mathematica

Time used: 0.0 (sec). Leaf size: 0

```
DSolve[y''[x]^3+y'[x]+1==x,y[x],x,IncludeSingularSolutions -> True]
```

Timed out

2.36 problem Problem 51

2.36.1 Solving as second order linear constant coeff ode	832
2.36.2 Solving as linear second order ode solved by an integrating factor ode	837
2.36.3 Solving using Kovacic algorithm	839
2.36.4 Maple step by step solution	845

Internal problem ID [12199]

Internal file name [OUTPUT/10851_Thursday_September_21_2023_05_48_04_AM_60851899/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 51.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff", "linear_second_order_ode_solved_by_an_integrating_factor"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$x'' + 10x' + 25x = 2^t + t e^{-5t}$$

2.36.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ax''(t) + Bx'(t) + Cx(t) = f(t)$$

Where $A = 1, B = 10, C = 25, f(t) = 2^t + t e^{-5t}$. Let the solution be

$$x = x_h + x_p$$

Where x_h is the solution to the homogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = 0$, and x_p is a particular solution to the non-homogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = f(t)$. x_h is the solution to

$$x'' + 10x' + 25x = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ax''(t) + Bx'(t) + Cx(t) = 0$$

Where in the above $A = 1, B = 10, C = 25$. Let the solution be $x = e^{\lambda t}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda t} + 10\lambda e^{\lambda t} + 25 e^{\lambda t} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda t}$ gives

$$\lambda^2 + 10\lambda + 25 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 10, C = 25$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{-10}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{(10)^2 - (4)(1)(25)} \\ &= -5 \end{aligned}$$

Hence this is the case of a double root $\lambda_{1,2} = 5$. Therefore the solution is

$$x = c_1 e^{-5t} + c_2 t e^{-5t} \quad (1)$$

Therefore the homogeneous solution x_h is

$$x_h = c_1 e^{-5t} + c_2 t e^{-5t}$$

The particular solution x_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on t as well. Let

$$x_p(t) = u_1 x_1 + u_2 x_2 \quad (1)$$

Where u_1, u_2 to be determined, and x_1, x_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$\begin{aligned} x_1 &= e^{-5t} \\ x_2 &= t e^{-5t} \end{aligned}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{x_2 f(t)}{aW(t)} \quad (2)$$

$$u_2 = \int \frac{x_1 f(t)}{aW(t)} \quad (3)$$

Where $W(t)$ is the Wronskian and a is the coefficient in front of x'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} x_1 & x_2 \\ x_1' & x_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-5t} & t e^{-5t} \\ \frac{d}{dt}(e^{-5t}) & \frac{d}{dt}(t e^{-5t}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-5t} & t e^{-5t} \\ -5 e^{-5t} & e^{-5t} - 5t e^{-5t} \end{vmatrix}$$

Therefore

$$W = (e^{-5t})(e^{-5t} - 5t e^{-5t}) - (t e^{-5t})(-5 e^{-5t})$$

Which simplifies to

$$W = e^{-10t}$$

Which simplifies to

$$W = e^{-10t}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{t e^{-5t}(2^t + t e^{-5t})}{e^{-10t}} dt$$

Which simplifies to

$$u_1 = - \int t(2^t e^{5t} + t) dt$$

Hence

$$u_1 = -\frac{t^3}{3} - \frac{t e^{5t} e^{t \ln(2)}}{\ln(2) + 5} + \frac{e^{5t} e^{t \ln(2)}}{(\ln(2) + 5)^2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-5t}(2^t + t e^{-5t})}{e^{-10t}} dt$$

Which simplifies to

$$u_2 = \int (2^t e^{5t} + t) dt$$

Hence

$$u_2 = \frac{t^2}{2} + \frac{e^{5t} e^{t \ln(2)}}{\ln(2) + 5}$$

Which simplifies to

$$u_1 = \frac{-3 \cdot 2^t (t \ln(2) + 5t - 1) e^{5t} - t^3 (\ln(2) + 5)^2}{3 (\ln(2) + 5)^2}$$
$$u_2 = \frac{2^{1+t} e^{5t} + t^2 (\ln(2) + 5)}{10 + 2 \ln(2)}$$

Therefore the particular solution, from equation (1) is

$$x_p(t) = \frac{(-3 \cdot 2^t (t \ln(2) + 5t - 1) e^{5t} - t^3 (\ln(2) + 5)^2) e^{-5t}}{3 (\ln(2) + 5)^2} + \frac{(2^{1+t} e^{5t} + t^2 (\ln(2) + 5)) t e^{-5t}}{10 + 2 \ln(2)}$$

Which simplifies to

$$x_p(t) = \frac{t^3 (\ln(2) + 5)^2 e^{-5t} + 6 \cdot 2^t}{6 (\ln(2) + 5)^2}$$

Therefore the general solution is

$$x = x_h + x_p$$
$$= (c_1 e^{-5t} + c_2 t e^{-5t}) + \left(\frac{t^3 (\ln(2) + 5)^2 e^{-5t} + 6 \cdot 2^t}{6 (\ln(2) + 5)^2} \right)$$

Which simplifies to

$$x = e^{-5t}(c_2t + c_1) + \frac{t^3(\ln(2) + 5)^2 e^{-5t} + 6 \cdot 2^t}{6(\ln(2) + 5)^2}$$

Summary

The solution(s) found are the following

$$x = e^{-5t}(c_2t + c_1) + \frac{t^3(\ln(2) + 5)^2 e^{-5t} + 6 \cdot 2^t}{6(\ln(2) + 5)^2} \quad (1)$$




Figure 120: Slope field plot

Verification of solutions

$$x = e^{-5t}(c_2t + c_1) + \frac{t^3(\ln(2) + 5)^2 e^{-5t} + 6 \cdot 2^t}{6(\ln(2) + 5)^2}$$

Verified OK.

2.36.2 Solving as linear second order ode solved by an integrating factor ode

The ode satisfies this form

$$x'' + p(t)x' + \frac{(p(t)^2 + p'(t))x}{2} = f(t)$$

Where $p(t) = 10$. Therefore, there is an integrating factor given by

$$\begin{aligned} M(x) &= e^{\frac{1}{2} \int p dx} \\ &= e^{\int 10 dx} \\ &= e^{5t} \end{aligned}$$

Multiplying both sides of the ODE by the integrating factor $M(x)$ makes the left side of the ODE a complete differential

$$\begin{aligned} (M(x)x)'' &= e^{5t}(2^t + te^{-5t}) \\ (e^{5t}x)'' &= e^{5t}(2^t + te^{-5t}) \end{aligned}$$

Integrating once gives

$$(e^{5t}x)' = \frac{2^{1+t}e^{5t} + t^2(\ln(2) + 5)}{10 + 2\ln(2)} + c_1$$

Integrating again gives

$$(e^{5t}x) = \frac{6 \cdot 2^t e^{5t} + t(\ln(2) + 5)^2 (t^2 + 6c_1)}{6(\ln(2) + 5)^2} + c_2$$

Hence the solution is

$$x = \frac{\frac{6 \cdot 2^t e^{5t} + t(\ln(2) + 5)^2 (t^2 + 6c_1)}{6(\ln(2) + 5)^2} + c_2}{e^{5t}}$$

Or

$$\begin{aligned} x &= \left(\frac{\ln(2)^2 t e^{-5t}}{(\ln(2) + 5)^2} + \frac{10 \ln(2) t e^{-5t}}{(\ln(2) + 5)^2} + \frac{25 t e^{-5t}}{(\ln(2) + 5)^2} \right) c_1 + \frac{2^t}{(\ln(2) + 5)^2} \\ &+ \frac{\ln(2)^2 t^3 e^{-5t}}{6(\ln(2) + 5)^2} + \frac{5 \ln(2) t^3 e^{-5t}}{3(\ln(2) + 5)^2} + \frac{25 t^3 e^{-5t}}{6(\ln(2) + 5)^2} + c_2 e^{-5t} \end{aligned}$$

Summary

The solution(s) found are the following

$$x = \left(\frac{\ln(2)^2 t e^{-5t}}{(\ln(2) + 5)^2} + \frac{10 \ln(2) t e^{-5t}}{(\ln(2) + 5)^2} + \frac{25 t e^{-5t}}{(\ln(2) + 5)^2} \right) c_1 + \frac{2^t}{(\ln(2) + 5)^2} \quad (1)$$
$$+ \frac{\ln(2)^2 t^3 e^{-5t}}{6 (\ln(2) + 5)^2} + \frac{5 \ln(2) t^3 e^{-5t}}{3 (\ln(2) + 5)^2} + \frac{25 t^3 e^{-5t}}{6 (\ln(2) + 5)^2} + c_2 e^{-5t}$$




Figure 121: Slope field plot

Verification of solutions

$$x = \left(\frac{\ln(2)^2 t e^{-5t}}{(\ln(2) + 5)^2} + \frac{10 \ln(2) t e^{-5t}}{(\ln(2) + 5)^2} + \frac{25 t e^{-5t}}{(\ln(2) + 5)^2} \right) c_1 + \frac{2^t}{(\ln(2) + 5)^2}$$
$$+ \frac{\ln(2)^2 t^3 e^{-5t}}{6 (\ln(2) + 5)^2} + \frac{5 \ln(2) t^3 e^{-5t}}{3 (\ln(2) + 5)^2} + \frac{25 t^3 e^{-5t}}{6 (\ln(2) + 5)^2} + c_2 e^{-5t}$$

Verified OK.

2.36.3 Solving using Kovacic algorithm

Writing the ode as

$$x'' + 10x' + 25x = 0 \quad (1)$$

$$Ax'' + Bx' + Cx = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned} A &= 1 \\ B &= 10 \\ C &= 25 \end{aligned} \quad (3)$$

Applying the Liouville transformation on the dependent variable gives

$$z(t) = xe^{\int \frac{B}{2A} dt}$$

Then (2) becomes

$$z''(t) = rz(t) \quad (4)$$

Where r is given by

$$\begin{aligned} r &= \frac{s}{t} \\ &= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2} \end{aligned} \quad (5)$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{0}{1} \quad (6)$$

Comparing the above to (5) shows that

$$\begin{aligned} s &= 0 \\ t &= 1 \end{aligned}$$

Therefore eq. (4) becomes

$$z''(t) = 0 \quad (7)$$

Equation (7) is now solved. After finding $z(t)$ then x is found using the inverse transformation

$$x = z(t) e^{-\int \frac{B}{2A} dt}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 92: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned} O(\infty) &= \deg(t) - \deg(s) \\ &= 0 - -\infty \\ &= \infty \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is *infinity* then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = 0$ is not a function of t , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(t) = 1$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in x is found from

$$\begin{aligned} x_1 &= z_1 e^{\int -\frac{1}{2} \frac{B}{A} dt} \\ &= z_1 e^{-\int \frac{1}{2} \frac{10}{1} dt} \end{aligned}$$

$$\begin{aligned}
&= z_1 e^{-5t} \\
&= z_1 (e^{-5t})
\end{aligned}$$

Which simplifies to

$$x_1 = e^{-5t}$$

The second solution x_2 to the original ode is found using reduction of order

$$x_2 = x_1 \int \frac{e^{\int -\frac{B}{A} dt}}{x_1^2} dt$$

Substituting gives

$$\begin{aligned}
x_2 &= x_1 \int \frac{e^{\int -\frac{10}{1} dt}}{(x_1)^2} dt \\
&= x_1 \int \frac{e^{-10t}}{(x_1)^2} dt \\
&= x_1(t)
\end{aligned}$$

Therefore the solution is

$$\begin{aligned}
x &= c_1 x_1 + c_2 x_2 \\
&= c_1 (e^{-5t}) + c_2 (e^{-5t}(t))
\end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$x = x_h + x_p$$

Where x_h is the solution to the homogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = 0$, and x_p is a particular solution to the nonhomogeneous ODE $Ax''(t) + Bx'(t) + Cx(t) = f(t)$. x_h is the solution to

$$x'' + 10x' + 25x = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$x_h = c_1 e^{-5t} + c_2 t e^{-5t}$$

The particular solution x_p can be found using either the method of undetermined coefficients, or the method of variation of parameters. The method of variation of parameters will be used as it is more general and can be used when the coefficients of the ODE depend on t as well. Let

$$x_p(t) = u_1x_1 + u_2x_2 \quad (1)$$

Where u_1, u_2 to be determined, and x_1, x_2 are the two basis solutions (the two linearly independent solutions of the homogeneous ODE) found earlier when solving the homogeneous ODE as

$$x_1 = e^{-5t}$$

$$x_2 = te^{-5t}$$

In the Variation of parameters u_1, u_2 are found using

$$u_1 = - \int \frac{x_2 f(t)}{aW(t)} \quad (2)$$

$$u_2 = \int \frac{x_1 f(t)}{aW(t)} \quad (3)$$

Where $W(t)$ is the Wronskian and a is the coefficient in front of x'' in the given ODE.

The Wronskian is given by $W = \begin{vmatrix} x_1 & x_2 \\ x_1' & x_2' \end{vmatrix}$. Hence

$$W = \begin{vmatrix} e^{-5t} & te^{-5t} \\ \frac{d}{dt}(e^{-5t}) & \frac{d}{dt}(te^{-5t}) \end{vmatrix}$$

Which gives

$$W = \begin{vmatrix} e^{-5t} & te^{-5t} \\ -5e^{-5t} & e^{-5t} - 5te^{-5t} \end{vmatrix}$$

Therefore

$$W = (e^{-5t})(e^{-5t} - 5te^{-5t}) - (te^{-5t})(-5e^{-5t})$$

Which simplifies to

$$W = e^{-10t}$$

Which simplifies to

$$W = e^{-10t}$$

Therefore Eq. (2) becomes

$$u_1 = - \int \frac{t e^{-5t}(2^t + t e^{-5t})}{e^{-10t}} dt$$

Which simplifies to

$$u_1 = - \int t(2^t e^{5t} + t) dt$$

Hence

$$u_1 = -\frac{t^3}{3} - \frac{t e^{5t} e^{t \ln(2)}}{\ln(2) + 5} + \frac{e^{5t} e^{t \ln(2)}}{(\ln(2) + 5)^2}$$

And Eq. (3) becomes

$$u_2 = \int \frac{e^{-5t}(2^t + t e^{-5t})}{e^{-10t}} dt$$

Which simplifies to

$$u_2 = \int (2^t e^{5t} + t) dt$$

Hence

$$u_2 = \frac{t^2}{2} + \frac{e^{5t} e^{t \ln(2)}}{\ln(2) + 5}$$

Which simplifies to

$$u_1 = \frac{-3 \cdot 2^t (t \ln(2) + 5t - 1) e^{5t} - t^3 (\ln(2) + 5)^2}{3 (\ln(2) + 5)^2}$$
$$u_2 = \frac{2^{1+t} e^{5t} + t^2 (\ln(2) + 5)}{10 + 2 \ln(2)}$$

Therefore the particular solution, from equation (1) is

$$x_p(t) = \frac{(-3 \cdot 2^t (t \ln(2) + 5t - 1) e^{5t} - t^3 (\ln(2) + 5)^2) e^{-5t}}{3 (\ln(2) + 5)^2} + \frac{(2^{1+t} e^{5t} + t^2 (\ln(2) + 5)) t e^{-5t}}{10 + 2 \ln(2)}$$

Which simplifies to

$$x_p(t) = \frac{t^3(\ln(2) + 5)^2 e^{-5t} + 6 \cdot 2^t}{6(\ln(2) + 5)^2}$$

Therefore the general solution is

$$\begin{aligned} x &= x_h + x_p \\ &= (c_1 e^{-5t} + c_2 t e^{-5t}) + \left(\frac{t^3(\ln(2) + 5)^2 e^{-5t} + 6 \cdot 2^t}{6(\ln(2) + 5)^2} \right) \end{aligned}$$

Which simplifies to

$$x = e^{-5t}(c_2 t + c_1) + \frac{t^3(\ln(2) + 5)^2 e^{-5t} + 6 \cdot 2^t}{6(\ln(2) + 5)^2}$$

Summary

The solution(s) found are the following

$$x = e^{-5t}(c_2 t + c_1) + \frac{t^3(\ln(2) + 5)^2 e^{-5t} + 6 \cdot 2^t}{6(\ln(2) + 5)^2} \quad (1)$$




Figure 122: Slope field plot

Verification of solutions

$$x = e^{-5t}(c_2t + c_1) + \frac{t^3(\ln(2) + 5)^2 e^{-5t} + 6 \cdot 2^t}{6(\ln(2) + 5)^2}$$

Verified OK.

2.36.4 Maple step by step solution

Let's solve

$$x'' + 10x' + 25x = 2^t + t e^{-5t}$$

- Highest derivative means the order of the ODE is 2

$$x''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 10r + 25 = 0$$

- Factor the characteristic polynomial

$$(r + 5)^2 = 0$$

- Root of the characteristic polynomial

$$r = -5$$

- 1st solution of the homogeneous ODE

$$x_1(t) = e^{-5t}$$

- Repeated root, multiply $x_1(t)$ by t to ensure linear independence

$$x_2(t) = t e^{-5t}$$

- General solution of the ODE

$$x = c_1 x_1(t) + c_2 x_2(t) + x_p(t)$$

- Substitute in solutions of the homogeneous ODE

$$x = c_1 e^{-5t} + c_2 t e^{-5t} + x_p(t)$$

- Find a particular solution $x_p(t)$ of the ODE

- Use variation of parameters to find x_p here $f(t)$ is the forcing function

$$\left[x_p(t) = -x_1(t) \left(\int \frac{x_2(t)f(t)}{W(x_1(t),x_2(t))} dt \right) + x_2(t) \left(\int \frac{x_1(t)f(t)}{W(x_1(t),x_2(t))} dt \right), f(t) = 2^t + t e^{-5t} \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(x_1(t), x_2(t)) = \begin{bmatrix} e^{-5t} & t e^{-5t} \\ -5 e^{-5t} & e^{-5t} - 5t e^{-5t} \end{bmatrix}$$

- Compute Wronskian

$$W(x_1(t), x_2(t)) = e^{-10t}$$

- Substitute functions into equation for $x_p(t)$

$$x_p(t) = e^{-5t} \left(- \left(\int t(2^t e^{5t} + t) dt \right) + \left(\int (2^t e^{5t} + t) dt \right) t \right)$$

- Compute integrals

$$x_p(t) = \frac{t^3(\ln(2)+5)^2 e^{-5t} + 6 \cdot 2^t}{6(\ln(2)+5)^2}$$

- Substitute particular solution into general solution to ODE

$$x = c_1 e^{-5t} + c_2 t e^{-5t} + \frac{t^3(\ln(2)+5)^2 e^{-5t} + 6 \cdot 2^t}{6(\ln(2)+5)^2}$$

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]
trying a double symmetry of the form [xi=0, eta=F(x)]
-> Try solving first the homogeneous part of the ODE
    checking if the LODE has constant coefficients
    <- constant coefficients successful
<- solving first the homogeneous part of the ODE successful`

```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 40

```
dsolve(diff(x(t),t$2)+10*diff(x(t),t)+25*x(t)=2^t+t*exp(-5*t),x(t), singsol=all)
```

$$x(t) = \frac{(\ln(2) + 5)^2 (t^3 + 6c_1 t + 6c_2) e^{-5t} + 6 \cdot 2^t}{6(\ln(2) + 5)^2}$$

✓ Solution by Mathematica

Time used: 0.341 (sec). Leaf size: 72

```
DSolve[x''[t]+10*x'[t]+25*x[t]==2^t+t*Exp[-5*t],x[t],t,IncludeSingularSolutions -> True]
```

$x(t)$

$$\rightarrow \frac{e^{-5t}(t^3(25 + \log^2(2) + \log(1024)) + 3 \cdot 2^{t+1}e^{5t} + c_2 t(150 + 6 \log^2(2) + \log(1152921504606846976)) + c_1}{6(5 + \log(2))^2}$$

2.37 problem Problem 52

2.37.1 Solving as second order nonlinear solved by mainardi lioville
method ode 848

Internal problem ID [12200]

Internal file name [OUTPUT/10852_Thursday_September_21_2023_05_48_07_AM_16469034/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 52.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_nonlinear_solved_by_mainardi_lioville_method"

Maple gives the following as the ode type

```
[_Liouville, [_2nd_order, _with_linear_symmetries], [_2nd_order, _reducible, _mu_x_y1], [_2nd_order, _reducible, _mu_xy]]
```

$$xyy'' - xy'^2 - yy' = 0$$

2.37.1 Solving as second order nonlinear solved by mainardi lioville method ode

The ode has the Liouville form given by

$$y'' + f(x)y' + g(y)y'^2 = 0 \tag{1A}$$

Where in this problem

$$f(x) = -\frac{1}{x}$$
$$g(y) = -\frac{1}{y}$$

Dividing through by y' then Eq (1A) becomes

$$\frac{y''}{y'} + f + gy' = 0 \tag{2A}$$

But the first term in Eq (2A) can be written as

$$\frac{y''}{y'} = \frac{d}{dx} \ln(y') \quad (3A)$$

And the last term in Eq (2A) can be written as

$$\begin{aligned} g \frac{dy}{dx} &= \left(\frac{d}{dy} \int g dy \right) \frac{dy}{dx} \\ &= \frac{d}{dx} \int g dy \end{aligned} \quad (4A)$$

Substituting (3A,4A) back into (2A) gives

$$\frac{d}{dx} \ln(y') + \frac{d}{dx} \int g dy = -f \quad (5A)$$

Integrating the above w.r.t. x gives

$$\ln(y') + \int g dy = - \int f dx + c_1$$

Where c_1 is arbitrary constant. Taking the exponential of the above gives

$$y' = c_2 e^{\int -g dy} e^{\int -f dx} \quad (6A)$$

Where c_2 is a new arbitrary constant. But since $g = -\frac{1}{y}$ and $f = -\frac{1}{x}$, then

$$\begin{aligned} \int -g dy &= \int \frac{1}{y} dy \\ &= \ln(y) \\ \int -f dx &= \int \frac{1}{x} dx \\ &= \ln(x) \end{aligned}$$

Substituting the above into Eq(6A) gives

$$y' = c_2 y x$$

Which is now solved as first order separable ode. In canonical form the ODE is

$$\begin{aligned} y' &= F(x, y) \\ &= f(x)g(y) \\ &= c_2 y x \end{aligned}$$

Where $f(x) = c_2x$ and $g(y) = y$. Integrating both sides gives

$$\begin{aligned}\frac{1}{y} dy &= c_2x dx \\ \int \frac{1}{y} dy &= \int c_2x dx \\ \ln(y) &= \frac{c_2x^2}{2} + c_3 \\ y &= e^{\frac{c_2x^2}{2} + c_3} \\ &= c_3e^{\frac{c_2x^2}{2}}\end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_3e^{\frac{c_2x^2}{2}} \quad (1)$$

Verification of solutions

$$y = c_3e^{\frac{c_2x^2}{2}}$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying 2nd order Liouville  
<- 2nd_order Liouville successful`
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 17

```
dsolve(x*y(x)*diff(y(x),x$2)-x*diff(y(x),x)^2-y(x)*diff(y(x),x)=0,y(x), singsol=all)
```

$$\begin{aligned}y(x) &= 0 \\ y(x) &= e^{\frac{c_1x^2}{2}} c_2\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.196 (sec). Leaf size: 19

```
DSolve[x*y[x]*y'[x]-x*y'[x]^2-y[x]*y'[x]==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_2 e^{\frac{c_1 x^2}{2}}$$

2.38 problem Problem 53

Internal problem ID [12201]

Internal file name [OUTPUT/10853_Thursday_September_21_2023_05_48_07_AM_23358333/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 53.

ODE order: 6.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _with_linear_symmetries]]
```

$$y^{(6)} - y = e^{2x}$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y^{(6)} - y = 0$$

The characteristic equation is

$$\lambda^6 - 1 = 0$$

The roots of the above equation are

$$\begin{aligned}\lambda_1 &= 1 \\ \lambda_2 &= -1 \\ \lambda_3 &= \frac{\sqrt{-2 - 2i\sqrt{3}}}{2} \\ \lambda_4 &= -\frac{\sqrt{-2 - 2i\sqrt{3}}}{2} \\ \lambda_5 &= \frac{\sqrt{-2 + 2i\sqrt{3}}}{2} \\ \lambda_6 &= -\frac{\sqrt{-2 + 2i\sqrt{3}}}{2}\end{aligned}$$

Therefore the homogeneous solution is

$$y_h(x) = c_1 e^{-x} + c_2 e^x + e^{\frac{\sqrt{-2+2i\sqrt{3}}x}{2}} c_3 + e^{\frac{\sqrt{-2-2i\sqrt{3}}x}{2}} c_4 + e^{-\frac{\sqrt{-2-2i\sqrt{3}}x}{2}} c_5 + e^{-\frac{\sqrt{-2+2i\sqrt{3}}x}{2}} c_6$$

The fundamental set of solutions for the homogeneous solution are the following

$$\begin{aligned}y_1 &= e^{-x} \\ y_2 &= e^x \\ y_3 &= e^{\frac{\sqrt{-2+2i\sqrt{3}}x}{2}} \\ y_4 &= e^{\frac{\sqrt{-2-2i\sqrt{3}}x}{2}} \\ y_5 &= e^{-\frac{\sqrt{-2-2i\sqrt{3}}x}{2}} \\ y_6 &= e^{-\frac{\sqrt{-2+2i\sqrt{3}}x}{2}}\end{aligned}$$

Now the particular solution to the given ODE is found

$$y^{(6)} - y = e^{2x}$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$e^{2x}$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^{2x}\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\left\{ e^x, e^{-x}, e^{-\frac{\sqrt{-2-2i\sqrt{3}}x}{2}}, e^{\frac{\sqrt{-2-2i\sqrt{3}}x}{2}}, e^{-\frac{\sqrt{-2+2i\sqrt{3}}x}{2}}, e^{\frac{\sqrt{-2+2i\sqrt{3}}x}{2}} \right\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 e^{2x}$$

The unknowns $\{A_1\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$63A_1 e^{2x} = e^{2x}$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{63} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^{2x}}{63}$$

Therefore the general solution is

$$y = y_h + y_p = \left(c_1 e^{-x} + c_2 e^x + e^{\frac{\sqrt{-2+2i\sqrt{3}}x}{2}} c_3 + e^{\frac{\sqrt{-2-2i\sqrt{3}}x}{2}} c_4 + e^{-\frac{\sqrt{-2-2i\sqrt{3}}x}{2}} c_5 + e^{-\frac{\sqrt{-2+2i\sqrt{3}}x}{2}} c_6 \right) + \left(\frac{e^{2x}}{63} \right)$$

Which simplifies to

$$y = c_1 e^{-x} + c_2 e^x + e^{\frac{(1+i\sqrt{3})x}{2}} c_3 + e^{-\frac{(i\sqrt{3}-1)x}{2}} c_4 + e^{\frac{(i\sqrt{3}-1)x}{2}} c_5 + e^{-\frac{(1+i\sqrt{3})x}{2}} c_6 + \frac{e^{2x}}{63}$$

Summary

The solution(s) found are the following

$$y = c_1 e^{-x} + c_2 e^x + e^{\frac{(1+i\sqrt{3})x}{2}} c_3 + e^{-\frac{(i\sqrt{3}-1)x}{2}} c_4 + e^{\frac{(i\sqrt{3}-1)x}{2}} c_5 + e^{-\frac{(1+i\sqrt{3})x}{2}} c_6 + \frac{e^{2x}}{63} \quad (1)$$

Verification of solutions

$$y = c_1 e^{-x} + c_2 e^x + e^{\frac{(1+i\sqrt{3})x}{2}} c_3 + e^{-\frac{(i\sqrt{3}-1)x}{2}} c_4 + e^{\frac{(i\sqrt{3}-1)x}{2}} c_5 + e^{-\frac{(1+i\sqrt{3})x}{2}} c_6 + \frac{e^{2x}}{63}$$

Verified OK.

Maple trace

```
`Methods for high order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 6; linear nonhomogeneous with symmetry [0,1]  
trying high order linear exact nonhomogeneous  
trying differential order: 6; missing the dependent variable  
checking if the LODE has constant coefficients  
<- constant coefficients successful`
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 65

```
dsolve(diff(y(x),x$6)-y(x)=exp(2*x),y(x), singsol=all)
```

$$y(x) = e^{-x} \left((c_3 e^{\frac{x}{2}} + c_5 e^{\frac{3x}{2}}) \cos\left(\frac{\sqrt{3}x}{2}\right) + (e^{\frac{x}{2}} c_4 + c_6 e^{\frac{3x}{2}}) \sin\left(\frac{\sqrt{3}x}{2}\right) + e^{2x} c_1 + \frac{e^{3x}}{63} + c_2 \right)$$

✓ Solution by Mathematica

Time used: 0.88 (sec). Leaf size: 85

```
DSolve[y''''''[x]-y[x]==Exp[2*x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{e^{2x}}{63} + c_1 e^x + c_4 e^{-x} + e^{-x/2} (c_2 e^x + c_3) \cos\left(\frac{\sqrt{3}x}{2}\right) + e^{-x/2} (c_6 e^x + c_5) \sin\left(\frac{\sqrt{3}x}{2}\right)$$

2.39 problem Problem 54

Internal problem ID [12202]

Internal file name [OUTPUT/10854_Thursday_September_21_2023_05_48_07_AM_3695680/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 54.

ODE order: 6.

ODE degree: 1.

The type(s) of ODE detected by this program : "**higher_order_linear_constant_coefficients_ODE**"

Maple gives the following as the ode type

```
[[_high_order , _missing_y]]
```

$$y^{(6)} + 2y'''' + y'' = x + e^x$$

This is higher order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE And y_p is a particular solution to the nonhomogeneous ODE. y_h is the solution to

$$y^{(6)} + 2y'''' + y'' = 0$$

The characteristic equation is

$$\lambda^6 + 2\lambda^4 + \lambda^2 = 0$$

The roots of the above equation are

$$\lambda_1 = 0$$

$$\lambda_2 = 0$$

$$\lambda_3 = i$$

$$\lambda_4 = -i$$

$$\lambda_5 = i$$

$$\lambda_6 = -i$$

Therefore the homogeneous solution is

$$y_h(x) = c_2x + c_1 + e^{ix}c_3 + xe^{ix}c_4 + e^{-ix}c_5 + xe^{-ix}c_6$$

The fundamental set of solutions for the homogeneous solution are the following

$$y_1 = 1$$

$$y_2 = x$$

$$y_3 = e^{ix}$$

$$y_4 = xe^{ix}$$

$$y_5 = e^{-ix}$$

$$y_6 = xe^{-ix}$$

Now the particular solution to the given ODE is found

$$y^{(6)} + 2y'''' + y'' = x + e^x$$

The particular solution is found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$x + e^x$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{e^x\}, \{1, x\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{1, x, xe^{ix}, xe^{-ix}, e^{ix}, e^{-ix}\}$$

Since 1 is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{e^x\}, \{x, x^2\}]$$

Since x is duplicated in the UC_set, then this basis is multiplied by extra x . The UC_set becomes

$$[\{e^x\}, \{x^2, x^3\}]$$

Since there was duplication between the basis functions in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis function in the above updated UC_set.

$$y_p = A_1e^x + A_2x^2 + A_3x^3$$

The unknowns $\{A_1, A_2, A_3\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$4A_1e^x + 2A_2 + 6A_3x = x + e^x$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = \frac{1}{4}, A_2 = 0, A_3 = \frac{1}{6} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = \frac{e^x}{4} + \frac{x^3}{6}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_2x + c_1 + e^{ix}c_3 + xe^{ix}c_4 + e^{-ix}c_5 + xe^{-ix}c_6) + \left(\frac{e^x}{4} + \frac{x^3}{6} \right) \end{aligned}$$

Which simplifies to

$$y = (c_6x + c_5)e^{-ix} + (c_4x + c_3)e^{ix} + c_2x + c_1 + \frac{e^x}{4} + \frac{x^3}{6}$$

Summary

The solution(s) found are the following

$$y = (c_6x + c_5)e^{-ix} + (c_4x + c_3)e^{ix} + c_2x + c_1 + \frac{e^x}{4} + \frac{x^3}{6} \quad (1)$$

Verification of solutions

$$y = (c_6x + c_5)e^{-ix} + (c_4x + c_3)e^{ix} + c_2x + c_1 + \frac{e^x}{4} + \frac{x^3}{6}$$

Verified OK.

Maple trace

```
`Methods for high order ODEs:
--- Trying classification methods ---
trying a quadrature
trying high order exact linear fully integrable
trying differential order: 6; linear nonhomogeneous with symmetry [0,1]
-> Calling odsolve with the ODE`, diff(diff(diff(diff(_b(_a), _a), _a), _a), _a) = -2*(diff(
  Methods for high order ODEs:
  --- Trying classification methods ---
  trying a quadrature
  trying high order exact linear fully integrable
  trying differential order: 4; linear nonhomogeneous with symmetry [0,1]
  trying high order linear exact nonhomogeneous
  trying differential order: 4; missing the dependent variable
  checking if the LODE has constant coefficients
  <- constant coefficients successful
<- differential order: 6; linear nonhomogeneous with symmetry [0,1] successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 46

```
dsolve(diff(y(x),x$6)+2*diff(y(x),x$4)+diff(y(x),x$2)=x+exp(x),y(x), singsol=all)
```

$$y(x) = (-c_3x - c_1 - 2c_4) \cos(x) + (-c_4x - c_2 + 2c_3) \sin(x) + \frac{x^3}{6} + c_5x + c_6 + \frac{e^x}{4}$$

✓ Solution by Mathematica

Time used: 0.61 (sec). Leaf size: 58

```
DSolve[y''''''[x]+2*y''''[x]+y''[x]==x+Exp[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{x^3}{6} + \frac{e^x}{4} + c_6x - (c_2x + c_1 + 2c_4) \cos(x) + (-c_4x + 2c_2 - c_3) \sin(x) + c_5$$

2.40 problem Problem 55

Internal problem ID [12203]

Internal file name [OUTPUT/10855_Thursday_September_21_2023_05_48_08_AM_58249007/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 55.

ODE order: 1.

ODE degree: 2.

The type(s) of ODE detected by this program : "**unknown**"

Maple gives the following as the ode type

```
[[_high_order, _missing_x], [_high_order, _missing_y], [
  _high_order, _with_linear_symmetries], [_high_order,
  _reducible, _mu_poly_yn]]
```

Unable to solve or complete the solution.

Unable to parse ODE.

Maple trace

```
`Methods for high order ODEs:
--- Trying classification methods ---
trying 4th order ODE linearizable_by_differentiation
trying high order reducible
trying differential order: 4; mu polynomial in y
-> Calling odsolve with the ODE`,  $-(5/6)*\ln(\text{diff}(\text{diff}(\_b(\_a), \_a), \_a))+\ln(\text{diff}(\text{diff}(\text{diff}(\_b(\_a), \_a), \_a), \_a), \_a))$ 
Methods for third order ODEs:
--- Trying classification methods ---
trying 3rd order ODE linearizable_by_differentiation
differential order: 3; trying a linearization to 4th order
trying differential order: 3; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`,  $\text{diff}(\_g(\_f), \_f) = \_g(\_f)^{(5/6)*\exp(-c\_1)}$ ,  $\_g(\_f)$ , HIN
symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[1, 0], [ $\_f$ ,  $6*\_g$ ]
```

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 25

```
dsolve(6*diff(y(x),x$2)*diff(y(x),x$4)-5*diff(y(x),x$3)^2=0,y(x), singsol=all)
```

$$y(x) = c_1x + c_2$$
$$y(x) = \frac{(c_2 + x)^8 c_1}{2612736} + c_3x + c_4$$

✓ Solution by Mathematica

Time used: 0.266 (sec). Leaf size: 26

```
DSolve[6*y''[x]*y''''[x]-5*y''''[x]^2==0,y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{56}c_2(x - 6c_1)^8 + c_4x + c_3$$

2.41 problem Problem 56

2.41.1 Solving as second order ode missing y ode 862

Internal problem ID [12204]

Internal file name [OUTPUT/10856_Thursday_September_21_2023_05_48_08_AM_39220548/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 56.

ODE order: 2.

ODE degree: 0.

The type(s) of ODE detected by this program : "second_order_ode_missing_y"

Maple gives the following as the ode type

```
[[_2nd_order , _missing_y]]
```

$$xy'' - y' \ln\left(\frac{y'}{x}\right) = 0$$

2.41.1 Solving as second order ode missing y ode

This is second order ode with missing dependent variable y . Let

$$p(x) = y'$$

Then

$$p'(x) = y''$$

Hence the ode becomes

$$p'(x)x - p(x) \ln\left(\frac{p(x)}{x}\right) = 0$$

Which is now solve for $p(x)$ as first order ode. Using the change of variables $p(x) = u(x)x$ on the above ode results in new ode in $u(x)$

$$(u'(x)x + u(x))x - u(x)x \ln(u(x)) = 0$$

In canonical form the ODE is

$$\begin{aligned}u' &= F(x, u) \\ &= f(x)g(u) \\ &= \frac{u(\ln(u) - 1)}{x}\end{aligned}$$

Where $f(x) = \frac{1}{x}$ and $g(u) = u(\ln(u) - 1)$. Integrating both sides gives

$$\begin{aligned}\frac{1}{u(\ln(u) - 1)} du &= \frac{1}{x} dx \\ \int \frac{1}{u(\ln(u) - 1)} du &= \int \frac{1}{x} dx \\ \ln(\ln(u) - 1) &= \ln(x) + c_2\end{aligned}$$

Raising both side to exponential gives

$$\ln(u) - 1 = e^{\ln(x) + c_2}$$

Which simplifies to

$$\ln(u) - 1 = c_3 x$$

Therefore the solution $p(x)$ is

$$\begin{aligned}p(x) &= xu \\ &= x e^{1 + c_3 e^{c_2 x}}\end{aligned}$$

Since $p = y'$ then the new first order ode to solve is

$$y' = x e^{1 + c_3 e^{c_2 x}}$$

Integrating both sides gives

$$\begin{aligned}y &= \int x e^{1 + c_3 e^{c_2 x}} dx \\ &= \frac{(c_3 e^{c_2 x} - 1) e^{1 + c_3 e^{c_2 x}} e^{-2c_2}}{c_3^2} + c_4\end{aligned}$$

Summary

The solution(s) found are the following

$$y = \frac{(c_3 e^{c_2 x} - 1) e^{1 + c_3 e^{c_2 x}} e^{-2c_2}}{c_3^2} + c_4 \quad (1)$$

Verification of solutions

$$y = \frac{(c_3 e^{c_2 x} - 1) e^{1+c_3 e^{c_2 x}} e^{-2c_2}}{c_3^2} + c_4$$

Verified OK.

Maple trace

```
`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
differential order: 2; trying a linearization to 3rd order
trying 2nd order ODE linearizable_by_differentiation
trying 2nd order, 2 integrating factors of the form mu(x,y)
trying differential order: 2; missing variables
`, `-> Computing symmetries using: way = 3
-> Calling odsolve with the ODE`, diff(_b(_a), _a) = _b(_a)*ln(_b(_a)/_a)/_a, _b(_a), HINT =
symmetry methods on request
`, `1st order, trying reduction of order with given symmetries: `[ _a, _b]
```

✓ Solution by Maple

Time used: 0.016 (sec). Leaf size: 31

```
dsolve(x*diff(y(x),x$2)=diff(y(x),x)*ln(diff(y(x),x)/x),y(x), singsol=all)
```

$$y(x) = \frac{e^{c_1 x + 1} c_1 x + c_2 c_1^2 - e^{c_1 x + 1}}{c_1^2}$$

✓ Solution by Mathematica

Time used: 0.905 (sec). Leaf size: 31

```
DSolve[x*y''[x]==y'[x]*Log[y'[x]/x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow e^{e^{c_1 x + 1} - 2c_1} (-1 + e^{c_1 x}) + c_2$$

2.42 problem Problem 57

2.42.1 Solving as second order linear constant coeff ode	865
2.42.2 Solving using Kovacic algorithm	868
2.42.3 Maple step by step solution	873

Internal problem ID [12205]

Internal file name [OUTPUT/10857_Thursday_September_21_2023_05_48_09_AM_58321793/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 57.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "kovacic", "second_order_linear_constant_coeff"

Maple gives the following as the ode type

```
[[_2nd_order , _linear , _nonhomogeneous]]
```

$$y'' + y = \sin(3x) \cos(x)$$

2.42.1 Solving as second order linear constant coeff ode

This is second order non-homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = f(x)$$

Where $A = 1, B = 0, C = 1, f(x) = \sin(3x) \cos(x)$. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the non-homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

This is second order with constant coefficients homogeneous ODE. In standard form the ODE is

$$Ay''(x) + By'(x) + Cy(x) = 0$$

Where in the above $A = 1, B = 0, C = 1$. Let the solution be $y = e^{\lambda x}$. Substituting this into the ODE gives

$$\lambda^2 e^{\lambda x} + e^{\lambda x} = 0 \quad (1)$$

Since exponential function is never zero, then dividing Eq(2) throughout by $e^{\lambda x}$ gives

$$\lambda^2 + 1 = 0 \quad (2)$$

Equation (2) is the characteristic equation of the ODE. Its roots determine the general solution form. Using the quadratic formula

$$\lambda_{1,2} = \frac{-B}{2A} \pm \frac{1}{2A} \sqrt{B^2 - 4AC}$$

Substituting $A = 1, B = 0, C = 1$ into the above gives

$$\begin{aligned} \lambda_{1,2} &= \frac{0}{(2)(1)} \pm \frac{1}{(2)(1)} \sqrt{0^2 - (4)(1)(1)} \\ &= \pm i \end{aligned}$$

Hence

$$\lambda_1 = +i$$

$$\lambda_2 = -i$$

Which simplifies to

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

Since roots are complex conjugate of each others, then let the roots be

$$\lambda_{1,2} = \alpha \pm i\beta$$

Where $\alpha = 0$ and $\beta = 1$. Therefore the final solution, when using Euler relation, can be written as

$$y = e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$$

Which becomes

$$y = e^0 (c_1 \cos(x) + c_2 \sin(x))$$

Or

$$y = c_1 \cos(x) + c_2 \sin(x)$$

Therefore the homogeneous solution y_h is

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(3x) \cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(2x), \sin(2x)\}, \{\cos(4x), \sin(4x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(2x) + A_2 \sin(2x) + A_3 \cos(4x) + A_4 \sin(4x)$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-3A_1 \cos(2x) - 3A_2 \sin(2x) - 15A_3 \cos(4x) - 15A_4 \sin(4x) = \sin(3x) \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = -\frac{1}{6}, A_3 = 0, A_4 = -\frac{1}{30} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{\sin(2x)}{6} - \frac{\sin(4x)}{30}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{\sin(2x)}{6} - \frac{\sin(4x)}{30} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{\sin(2x)}{6} - \frac{\sin(4x)}{30} \quad (1)$$




Figure 123: Slope field plot

Verification of solutions

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{\sin(2x)}{6} - \frac{\sin(4x)}{30}$$

Verified OK.

2.42.2 Solving using Kovacic algorithm

Writing the ode as

$$y'' + y = 0 \quad (1)$$

$$Ay'' + By' + Cy = 0 \quad (2)$$

Comparing (1) and (2) shows that

$$\begin{aligned}A &= 1 \\B &= 0 \\C &= 1\end{aligned}\tag{3}$$

Applying the Liouville transformation on the dependent variable gives

$$z(x) = ye^{\int \frac{B}{2A} dx}$$

Then (2) becomes

$$z''(x) = rz(x)\tag{4}$$

Where r is given by

$$\begin{aligned}r &= \frac{s}{t} \\&= \frac{2AB' - 2BA' + B^2 - 4AC}{4A^2}\end{aligned}\tag{5}$$

Substituting the values of A, B, C from (3) in the above and simplifying gives

$$r = \frac{-1}{1}\tag{6}$$

Comparing the above to (5) shows that

$$\begin{aligned}s &= -1 \\t &= 1\end{aligned}$$

Therefore eq. (4) becomes

$$z''(x) = -z(x)\tag{7}$$

Equation (7) is now solved. After finding $z(x)$ then y is found using the inverse transformation

$$y = z(x) e^{-\int \frac{B}{2A} dx}$$

The first step is to determine the case of Kovacic algorithm this ode belongs to. There are 3 cases depending on the order of poles of r and the order of r at ∞ . The following table summarizes these cases.

Case	Allowed pole order for r	Allowed value for $\mathcal{O}(\infty)$
1	$\{0, 1, 2, 4, 6, 8, \dots\}$	$\{\dots, -6, -4, -2, 0, 2, 3, 4, 5, 6, \dots\}$
2	Need to have at least one pole that is either order 2 or odd order greater than 2. Any other pole order is allowed as long as the above condition is satisfied. Hence the following set of pole orders are all allowed. $\{1, 2\}, \{1, 3\}, \{2\}, \{3\}, \{3, 4\}, \{1, 2, 5\}$.	no condition
3	$\{1, 2\}$	$\{2, 3, 4, 5, 6, 7, \dots\}$

Table 94: Necessary conditions for each Kovacic case

The order of r at ∞ is the degree of t minus the degree of s . Therefore

$$\begin{aligned}
 \mathcal{O}(\infty) &= \deg(t) - \deg(s) \\
 &= 0 - 0 \\
 &= 0
 \end{aligned}$$

There are no poles in r . Therefore the set of poles Γ is empty. Since there is no odd order pole larger than 2 and the order at ∞ is 0 then the necessary conditions for case one are met. Therefore

$$L = [1]$$

Since $r = -1$ is not a function of x , then there is no need run Kovacic algorithm to obtain a solution for transformed ode $z'' = rz$ as one solution is

$$z_1(x) = \cos(x)$$

Using the above, the solution for the original ode can now be found. The first solution to the original ode in y is found from

$$y_1 = z_1 e^{\int -\frac{1}{2} \frac{B}{A} dx}$$

Since $B = 0$ then the above reduces to

$$\begin{aligned}
 y_1 &= z_1 \\
 &= \cos(x)
 \end{aligned}$$

Which simplifies to

$$y_1 = \cos(x)$$

The second solution y_2 to the original ode is found using reduction of order

$$y_2 = y_1 \int \frac{e^{\int -\frac{B}{A} dx}}{y_1^2} dx$$

Since $B = 0$ then the above becomes

$$\begin{aligned} y_2 &= y_1 \int \frac{1}{y_1^2} dx \\ &= \cos(x) \int \frac{1}{\cos(x)^2} dx \\ &= \cos(x) (\tan(x)) \end{aligned}$$

Therefore the solution is

$$\begin{aligned} y &= c_1 y_1 + c_2 y_2 \\ &= c_1 (\cos(x)) + c_2 (\cos(x) (\tan(x))) \end{aligned}$$

This is second order nonhomogeneous ODE. Let the solution be

$$y = y_h + y_p$$

Where y_h is the solution to the homogeneous ODE $Ay''(x) + By'(x) + Cy(x) = 0$, and y_p is a particular solution to the nonhomogeneous ODE $Ay''(x) + By'(x) + Cy(x) = f(x)$. y_h is the solution to

$$y'' + y = 0$$

The homogeneous solution is found using the Kovacic algorithm which results in

$$y_h = c_1 \cos(x) + c_2 \sin(x)$$

The particular solution is now found using the method of undetermined coefficients. Looking at the RHS of the ode, which is

$$\sin(3x) \cos(x)$$

Shows that the corresponding undetermined set of the basis functions (UC_set) for the trial solution is

$$[\{\cos(2x), \sin(2x)\}, \{\cos(4x), \sin(4x)\}]$$

While the set of the basis functions for the homogeneous solution found earlier is

$$\{\cos(x), \sin(x)\}$$

Since there is no duplication between the basis function in the UC_set and the basis functions of the homogeneous solution, the trial solution is a linear combination of all the basis in the UC_set.

$$y_p = A_1 \cos(2x) + A_2 \sin(2x) + A_3 \cos(4x) + A_4 \sin(4x)$$

The unknowns $\{A_1, A_2, A_3, A_4\}$ are found by substituting the above trial solution y_p into the ODE and comparing coefficients. Substituting the trial solution into the ODE and simplifying gives

$$-3A_1 \cos(2x) - 3A_2 \sin(2x) - 15A_3 \cos(4x) - 15A_4 \sin(4x) = \sin(3x) \cos(x)$$

Solving for the unknowns by comparing coefficients results in

$$\left[A_1 = 0, A_2 = -\frac{1}{6}, A_3 = 0, A_4 = -\frac{1}{30} \right]$$

Substituting the above back in the above trial solution y_p , gives the particular solution

$$y_p = -\frac{\sin(2x)}{6} - \frac{\sin(4x)}{30}$$

Therefore the general solution is

$$\begin{aligned} y &= y_h + y_p \\ &= (c_1 \cos(x) + c_2 \sin(x)) + \left(-\frac{\sin(2x)}{6} - \frac{\sin(4x)}{30} \right) \end{aligned}$$

Summary

The solution(s) found are the following

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{\sin(2x)}{6} - \frac{\sin(4x)}{30} \quad (1)$$




Figure 124: Slope field plot

Verification of solutions

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{\sin(2x)}{6} - \frac{\sin(4x)}{30}$$

Verified OK.

2.42.3 Maple step by step solution

Let's solve

$$y'' + y = \sin(3x) \cos(x)$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Characteristic polynomial of homogeneous ODE

$$r^2 + 1 = 0$$

- Use quadratic formula to solve for r

$$r = \frac{0 \pm (\sqrt{-4})}{2}$$

- Roots of the characteristic polynomial

$$r = (-I, I)$$

- 1st solution of the homogeneous ODE

$$y_1(x) = \cos(x)$$

- 2nd solution of the homogeneous ODE

$$y_2(x) = \sin(x)$$

- General solution of the ODE

$$y = c_1 y_1(x) + c_2 y_2(x) + y_p(x)$$

- Substitute in solutions of the homogeneous ODE

$$y = c_1 \cos(x) + c_2 \sin(x) + y_p(x)$$

- Find a particular solution $y_p(x)$ of the ODE

- Use variation of parameters to find y_p here $f(x)$ is the forcing function

$$\left[y_p(x) = -y_1(x) \left(\int \frac{y_2(x)f(x)}{W(y_1(x),y_2(x))} dx \right) + y_2(x) \left(\int \frac{y_1(x)f(x)}{W(y_1(x),y_2(x))} dx \right), f(x) = \sin(3x) \cos(x) \right]$$

- Wronskian of solutions of the homogeneous equation

$$W(y_1(x), y_2(x)) = \begin{bmatrix} \cos(x) & \sin(x) \\ -\sin(x) & \cos(x) \end{bmatrix}$$

- Compute Wronskian

$$W(y_1(x), y_2(x)) = 1$$

- Substitute functions into equation for $y_p(x)$

$$y_p(x) = -\frac{\cos(x) \left(\int (\cos(x) - \cos(5x)) dx \right)}{4} + \sin(x) \left(\int \cos(x)^2 \sin(3x) dx \right)$$

- Compute integrals

$$y_p(x) = -\frac{\sin(x) \cos(x) (4 \cos(x)^2 + 3)}{15}$$

- Substitute particular solution into general solution to ODE

$$y = c_1 \cos(x) + c_2 \sin(x) - \frac{\sin(x) \cos(x) (4 \cos(x)^2 + 3)}{15}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying a quadrature  
trying high order exact linear fully integrable  
trying differential order: 2; linear nonhomogeneous with symmetry [0,1]  
trying a double symmetry of the form [xi=0, eta=F(x)]  
-> Try solving first the homogeneous part of the ODE  
    checking if the LODE has constant coefficients  
    <- constant coefficients successful  
<- solving first the homogeneous part of the ODE successful`
```

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 25

```
dsolve(diff(y(x),x$2)+y(x)=sin(3*x)*cos(x),y(x), singsol=all)
```

$$y(x) = \sin(x) c_2 + c_1 \cos(x) - \frac{\sin(2x)}{6} - \frac{\sin(4x)}{30}$$

✓ Solution by Mathematica

Time used: 0.187 (sec). Leaf size: 30

```
DSolve[y''[x]+y[x]==Sin[3*x]*Cos[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow c_1 \cos(x) - \frac{1}{15} \sin(x)(6 \cos(x) + \cos(3x) - 15c_2)$$

2.43 problem Problem 58

- 2.43.1 Solving as second order ode can be made integrable ode 876
- 2.43.2 Solving as second order ode missing x ode 878
- 2.43.3 Maple step by step solution 881

Internal problem ID [12206]

Internal file name [OUTPUT/10858_Thursday_September_21_2023_05_48_11_AM_81150646/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 58.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "second_order_ode_missing_x", "second_order_ode_can_be_made_integrable"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1]]
```

$$y'' - 2y^3 = 0$$

With initial conditions

$$[y(1) = 1, y'(1) = 1]$$

2.43.1 Solving as second order ode can be made integrable ode

Multiplying the ode by y' gives

$$y'y'' - 2y^3y' = 0$$

Integrating the above w.r.t x gives

$$\int (y'y'' - 2y^3y') dx = 0$$
$$\frac{y'^2}{2} - \frac{y^4}{2} = c_2$$

Which is now solved for y . Solving the given ode for y' results in 2 differential equations to solve. Each one of these will generate a solution. The equations generated are

$$y' = \sqrt{y^4 + 2c_1} \quad (1)$$

$$y' = -\sqrt{y^4 + 2c_1} \quad (2)$$

Now each one of the above ODE is solved.

Solving equation (1)

Integrating both sides gives

$$\int \frac{1}{\sqrt{y^4 + 2c_1}} dy = \int dx$$

$$\int^y \frac{1}{\sqrt{-a^4 + 2c_1}} d_a = x + c_2$$

Solving equation (2)

Integrating both sides gives

$$\int -\frac{1}{\sqrt{y^4 + 2c_1}} dy = \int dx$$

$$\int^y -\frac{1}{\sqrt{-a^4 + 2c_1}} d_a = c_3 + x$$

Initial conditions are used to solve for the constants of integration.

Looking at the First solution

$$\int^y \frac{1}{\sqrt{-a^4 + 2c_1}} d_a = x + c_2 \quad (1)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 1$ in the above gives

$$\int^1 \frac{1}{\sqrt{-a^4 + 2c_1}} d_a = 1 + c_2 \quad (1A)$$

Taking derivative of the solution gives

$$y' = \sqrt{\text{RootOf} \left(- \left(\int^{-z} \frac{1}{\sqrt{-a^4 + 2c_1}} d_a \right) + x + c_2 \right)^4 + 2c_1}$$

substituting $y' = 1$ and $x = 1$ in the above gives

$$1 = \sqrt{\text{RootOf} \left(- \left(\int^{-z} \frac{1}{\sqrt{-a^4 + 2c_1}} d_{-a} \right) + 1 + c_2 \right)^4 + 2c_1} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_2\}$. There is no solution for the constants of integrations. This solution is removed.

Looking at the Second solution

$$\int^y -\frac{1}{\sqrt{-a^4 + 2c_1}} d_{-a} = c_3 + x \quad (2)$$

Initial conditions are now substituted in the above solution. This will generate the required equations to solve for the integration constants. substituting $y = 1$ and $x = 1$ in the above gives

$$-\left(\int^1 \frac{1}{\sqrt{-a^4 + 2c_1}} d_{-a} \right) = 1 + c_3 \quad (1A)$$

Taking derivative of the solution gives

$$y' = -\sqrt{\text{RootOf} \left(- \left(\int^{-z} -\frac{1}{\sqrt{-a^4 + 2c_1}} d_{-a} \right) + c_3 + x \right)^4 + 2c_1}$$

substituting $y' = 1$ and $x = 1$ in the above gives

$$1 = -\sqrt{\text{RootOf} \left(\int^{-z} \frac{1}{\sqrt{-a^4 + 2c_1}} d_{-a} + c_3 + 1 \right)^4 + 2c_1} \quad (2A)$$

Equations {1A,2A} are now solved for $\{c_1, c_3\}$. There is no solution for the constants of integrations. This solution is removed.

Verification of solutions N/A

2.43.2 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned}y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy}\end{aligned}$$

Hence the ode becomes

$$p(y) \left(\frac{d}{dy} p(y) \right) - 2y^3 = 0$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned}p' &= F(y, p) \\ &= f(y)g(p) \\ &= \frac{2y^3}{p}\end{aligned}$$

Where $f(y) = 2y^3$ and $g(p) = \frac{1}{p}$. Integrating both sides gives

$$\begin{aligned}\frac{1}{\frac{1}{p}} dp &= 2y^3 dy \\ \int \frac{1}{\frac{1}{p}} dp &= \int 2y^3 dy \\ \frac{p^2}{2} &= \frac{y^4}{2} + c_1\end{aligned}$$

The solution is

$$\frac{p(y)^2}{2} - \frac{y^4}{2} - c_1 = 0$$

Initial conditions are used to solve for c_1 . Substituting $y = 1$ and $p = 1$ in the above solution gives an equation to solve for the constant of integration.

$$-c_1 = 0$$

$$c_1 = 0$$

Substituting c_1 found above in the general solution gives

$$\frac{p^2}{2} - \frac{y^4}{2} = 0$$

Solving for $p(y)$ from the above gives

$$p(y) = y^2$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = y^2$$

Integrating both sides gives

$$\int \frac{1}{y^2} dy = x + c_2$$
$$-\frac{1}{y} = x + c_2$$

Solving for y gives these solutions

$$y_1 = -\frac{1}{x + c_2}$$

Initial conditions are used to solve for c_2 . Substituting $x = 1$ and $y = 1$ in the above solution gives an equation to solve for the constant of integration.

$$1 = -\frac{1}{1 + c_2}$$

$$c_2 = -2$$

Substituting c_2 found above in the general solution gives

$$y = -\frac{1}{x - 2}$$

Initial conditions are used to solve for the constants of integration.

Summary

The solution(s) found are the following

$$y = -\frac{1}{x - 2} \tag{1}$$




Figure 125: Solution plot

Verification of solutions

$$y = -\frac{1}{x-2}$$

Verified OK.

2.43.3 Maple step by step solution

Let's solve

$$\left[y'' - 2y^3 = 0, y(1) = 1, y'|_{\{x=1\}} = 1 \right]$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Define new dependent variable u

$$u(x) = y'$$

- Compute y''

$$u'(x) = y''$$

- Use chain rule on the lhs

$$y' \left(\frac{d}{dy} u(y) \right) = y''$$

- Substitute in the definition of u

$$u(y) \left(\frac{d}{dy} u(y) \right) = y''$$

- Make substitutions $y' = u(y)$, $y'' = u(y) \left(\frac{d}{dy} u(y) \right)$ to reduce order of ODE

$$u(y) \left(\frac{d}{dy} u(y) \right) - 2y^3 = 0$$

- Integrate both sides with respect to y

$$\int \left(u(y) \left(\frac{d}{dy} u(y) \right) - 2y^3 \right) dy = \int 0 dy + c_1$$

- Evaluate integral

$$\frac{u(y)^2}{2} - \frac{y^4}{2} = c_1$$

- Solve for $u(y)$

$$\left\{ u(y) = \sqrt{y^4 + 2c_1}, u(y) = -\sqrt{y^4 + 2c_1} \right\}$$

- Solve 1st ODE for $u(y)$

$$u(y) = \sqrt{y^4 + 2c_1}$$

- Revert to original variables with substitution $u(y) = y'$, $y = y$

$$y' = \sqrt{y^4 + 2c_1}$$

- Separate variables

$$\frac{y'}{\sqrt{y^4 + 2c_1}} = 1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\sqrt{y^4 + 2c_1}} dx = \int 1 dx + c_2$$

- Evaluate integral

$$\frac{\sqrt{2} \sqrt{4 - \frac{21y^2\sqrt{2}}{\sqrt{c_1}}} \sqrt{4 + \frac{21y^2\sqrt{2}}{\sqrt{c_1}}} \text{EllipticF} \left(\frac{y\sqrt{2} \sqrt{\frac{1\sqrt{2}}{\sqrt{c_1}}}}{2}, \mathbf{I} \right)}{4 \sqrt{\frac{1\sqrt{2}}{\sqrt{c_1}}} \sqrt{y^4 + 2c_1}} = x + c_2$$

- Solve for y

$$\left\{ \frac{\text{JacobiSN} \left(\sqrt{1\sqrt{c_1}} \sqrt{2} (x+c_2), \mathbf{I} \right) \sqrt{2}}{\sqrt{\frac{1\sqrt{2} \text{RootOf} \left(-Z^2 - c_1, \text{index}=1 \right)}{c_1}}}, - \frac{\text{JacobiSN} \left(\sqrt{1\sqrt{c_1}} \sqrt{2} (x+c_2), \mathbf{I} \right) \sqrt{2}}{\sqrt{\frac{1\sqrt{2} \text{RootOf} \left(-Z^2 - c_1, \text{index}=1 \right)}{c_1}}} \right\}$$

- Solve 2nd ODE for $u(y)$

$$u(y) = -\sqrt{y^4 + 2c_1}$$

- Revert to original variables with substitution $u(y) = y', y = y$

$$y' = -\sqrt{y^4 + 2c_1}$$

- Separate variables

$$\frac{y'}{\sqrt{y^4 + 2c_1}} = -1$$

- Integrate both sides with respect to x

$$\int \frac{y'}{\sqrt{y^4 + 2c_1}} dx = \int (-1) dx + c_2$$

- Evaluate integral

$$\frac{\sqrt{2} \sqrt{4 - \frac{2Iy^2\sqrt{2}}{\sqrt{c_1}}} \sqrt{4 + \frac{2Iy^2\sqrt{2}}{\sqrt{c_1}}} \text{EllipticF}\left(\frac{y\sqrt{2}\sqrt{\frac{I\sqrt{2}}{\sqrt{c_1}}}, I\right)}{4\sqrt{\frac{I\sqrt{2}}{\sqrt{c_1}}} \sqrt{y^4 + 2c_1}} = -x + c_2$$

- Solve for y

$$\left\{ \frac{\text{JacobiSN}\left(\sqrt{I\sqrt{c_1}}\sqrt{2}(-x+c_2), I\right)\sqrt{2}}{\sqrt{\frac{I\sqrt{2}\text{RootOf}\left(-Z^2 - c_1, index=1\right)}{c_1}}}, -\frac{\text{JacobiSN}\left(\sqrt{I\sqrt{c_1}}\sqrt{2}(-x+c_2), I\right)\sqrt{2}}{\sqrt{\frac{I\sqrt{2}\text{RootOf}\left(-Z^2 - c_1, index=1\right)}{c_1}}} \right\}$$

□ Check validity of solution $\frac{\text{JacobiSN}\left(\sqrt{I\sqrt{c_1}}\sqrt{2}(-x+c_2), I\right)\sqrt{2}}{\sqrt{\frac{I\sqrt{2}\text{RootOf}\left(-Z^2 - c_1, index=1\right)}{c_1}}}$

- Use initial condition $y(1) = 1$

$$\frac{\text{JacobiSN}\left(\sqrt{I\sqrt{c_1}}\sqrt{2}(-1+c_2), I\right)\sqrt{2}}{\sqrt{\frac{I\sqrt{2}\text{RootOf}\left(-Z^2 - c_1, index=1\right)}{c_1}}}$$

- Compute derivative of the solution

$$\frac{\sqrt{I\sqrt{c_1}}\sqrt{2}\text{JacobiCN}\left(\sqrt{I\sqrt{c_1}}\sqrt{2}(-x+c_2), I\right)\text{JacobiDN}\left(\sqrt{I\sqrt{c_1}}\sqrt{2}(-x+c_2), I\right)\sqrt{2}}{\sqrt{\frac{I\sqrt{2}\text{RootOf}\left(-Z^2 - c_1, index=1\right)}{c_1}}}$$

- Use the initial condition $y'|_{\{x=1\}} = 1$

$$\frac{\sqrt{I\sqrt{c_1}}\sqrt{2}\text{JacobiCN}\left(\sqrt{I\sqrt{c_1}}\sqrt{2}(-1+c_2), I\right)\text{JacobiDN}\left(\sqrt{I\sqrt{c_1}}\sqrt{2}(-1+c_2), I\right)\sqrt{2}}{\sqrt{\frac{I\sqrt{2}\text{RootOf}\left(-Z^2 - c_1, index=1\right)}{c_1}}}$$

- Solve for c_1 and c_2

- The solution does not satisfy the initial conditions
- Check validity of solution
$$\frac{\text{JacobiSN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(x+c_2),I\right)\sqrt{2}}{\sqrt{\frac{I\sqrt{2}\text{RootOf}\left(-Z^2-c_1,index=1\right)}{c_1}}}$$
- Use initial condition $y(1) = 1$
- $$\frac{\text{JacobiSN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(1+c_2),I\right)\sqrt{2}}{\sqrt{\frac{I\sqrt{2}\text{RootOf}\left(-Z^2-c_1,index=1\right)}{c_1}}}$$
- Compute derivative of the solution
- $$\frac{\sqrt{I\sqrt{c_1}\sqrt{2}}\text{JacobiCN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(x+c_2),I\right)\text{JacobiDN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(x+c_2),I\right)\sqrt{2}}{\sqrt{\frac{I\sqrt{2}\text{RootOf}\left(-Z^2-c_1,index=1\right)}{c_1}}}$$
- Use the initial condition $y' \Big|_{\{x=1\}} = 1$
- $$\frac{\sqrt{I\sqrt{c_1}\sqrt{2}}\text{JacobiCN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(1+c_2),I\right)\text{JacobiDN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(1+c_2),I\right)\sqrt{2}}{\sqrt{\frac{I\sqrt{2}\text{RootOf}\left(-Z^2-c_1,index=1\right)}{c_1}}}$$
- Solve for c_1 and c_2
- The solution does not satisfy the initial conditions
- Check validity of solution
$$-\frac{\text{JacobiSN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(-x+c_2),I\right)\sqrt{2}}{\sqrt{\frac{I\sqrt{2}\text{RootOf}\left(-Z^2-c_1,index=1\right)}{c_1}}}$$
- Use initial condition $y(1) = 1$
- $$-\frac{\text{JacobiSN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(-1+c_2),I\right)\sqrt{2}}{\sqrt{\frac{I\sqrt{2}\text{RootOf}\left(-Z^2-c_1,index=1\right)}{c_1}}}$$
- Compute derivative of the solution
- $$\frac{\sqrt{I\sqrt{c_1}\sqrt{2}}\text{JacobiCN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(-x+c_2),I\right)\text{JacobiDN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(-x+c_2),I\right)\sqrt{2}}{\sqrt{\frac{I\sqrt{2}\text{RootOf}\left(-Z^2-c_1,index=1\right)}{c_1}}}$$
- Use the initial condition $y' \Big|_{\{x=1\}} = 1$

$$\frac{\sqrt{I\sqrt{c_1}\sqrt{2}} \operatorname{JacobiCN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(-1+c_2), I\right) \operatorname{JacobiDN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(-1+c_2), I\right) \sqrt{2}}{\sqrt{\frac{I\sqrt{2} \operatorname{RootOf}\left(-Z^2 - c_1, \operatorname{index}=1\right)}{c_1}}}$$

- Solve for c_1 and c_2
- The solution does not satisfy the initial conditions

□ Check validity of solution $-\frac{\operatorname{JacobiSN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(x+c_2), I\right) \sqrt{2}}{\sqrt{\frac{I\sqrt{2} \operatorname{RootOf}\left(-Z^2 - c_1, \operatorname{index}=1\right)}{c_1}}}$

- Use initial condition $y(1) = 1$

$$\frac{\operatorname{JacobiSN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(1+c_2), I\right) \sqrt{2}}{\sqrt{\frac{I\sqrt{2} \operatorname{RootOf}\left(-Z^2 - c_1, \operatorname{index}=1\right)}{c_1}}}$$

- Compute derivative of the solution

$$\frac{\sqrt{I\sqrt{c_1}\sqrt{2}} \operatorname{JacobiCN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(x+c_2), I\right) \operatorname{JacobiDN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(x+c_2), I\right) \sqrt{2}}{\sqrt{\frac{I\sqrt{2} \operatorname{RootOf}\left(-Z^2 - c_1, \operatorname{index}=1\right)}{c_1}}}$$

- Use the initial condition $y'|_{\{x=1\}} = 1$

$$\frac{\sqrt{I\sqrt{c_1}\sqrt{2}} \operatorname{JacobiCN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(1+c_2), I\right) \operatorname{JacobiDN}\left(\sqrt{I\sqrt{c_1}\sqrt{2}}(1+c_2), I\right) \sqrt{2}}{\sqrt{\frac{I\sqrt{2} \operatorname{RootOf}\left(-Z^2 - c_1, \operatorname{index}=1\right)}{c_1}}}$$

- Solve for c_1 and c_2
- The solution does not satisfy the initial conditions

Maple trace

```

`Methods for second order ODEs:
--- Trying classification methods ---
trying 2nd order Liouville
trying 2nd order WeierstrassP
trying 2nd order JacobiSN
<- 2nd_order JacobiSN successful`

```

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 11

```
dsolve([diff(y(x),x$2)=2*y(x)^3,y(1) = 1, D(y)(1) = 1],y(x), singsol=all)
```

$$y(x) = -\frac{1}{x-2}$$

✓ Solution by Mathematica

Time used: 0.143 (sec). Leaf size: 12

```
DSolve[{y'[x]==2*y[x]^3,{y[1]==1,y'[1]==1}},y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1}{2-x}$$

2.44 problem Problem 59

- 2.44.1 Solving as second order ode missing x ode 887
- 2.44.2 Maple step by step solution 889

Internal problem ID [12207]

Internal file name [OUTPUT/10859_Thursday_September_21_2023_05_48_12_AM_51619486/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 2, DIFFERENTIAL EQUATIONS OF THE SECOND ORDER AND HIGHER. Problems page 172

Problem number: Problem 59.

ODE order: 2.

ODE degree: 1.

The type(s) of ODE detected by this program : "**second_order_ode_missing_x**"

Maple gives the following as the ode type

```
[[_2nd_order, _missing_x], [_2nd_order, _reducible, _mu_x_y1],  
 [_2nd_order, _reducible, _mu_xy]]
```

$$yy'' - y'^2 - y' = 0$$

2.44.1 Solving as second order ode missing x ode

This is missing independent variable second order ode. Solved by reduction of order by using substitution which makes the dependent variable y an independent variable. Using

$$y' = p(y)$$

Then

$$\begin{aligned} y'' &= \frac{dp}{dx} \\ &= \frac{dy}{dx} \frac{dp}{dy} \\ &= p \frac{dp}{dy} \end{aligned}$$

Hence the ode becomes

$$yp(y) \left(\frac{d}{dy} p(y) \right) + (-p(y) - 1) p(y) = 0$$

Which is now solved as first order ode for $p(y)$. In canonical form the ODE is

$$\begin{aligned} p' &= F(y, p) \\ &= f(y)g(p) \\ &= \frac{p+1}{y} \end{aligned}$$

Where $f(y) = \frac{1}{y}$ and $g(p) = p + 1$. Integrating both sides gives

$$\begin{aligned} \frac{1}{p+1} dp &= \frac{1}{y} dy \\ \int \frac{1}{p+1} dp &= \int \frac{1}{y} dy \\ \ln(p+1) &= \ln(y) + c_1 \end{aligned}$$

Raising both side to exponential gives

$$p+1 = e^{\ln(y)+c_1}$$

Which simplifies to

$$p+1 = c_2 y$$

Which simplifies to

$$p(y) = c_2 y e^{c_1} - 1$$

For solution (1) found earlier, since $p = y'$ then we now have a new first order ode to solve which is

$$y' = c_2 y e^{c_1} - 1$$

Integrating both sides gives

$$\begin{aligned} \int \frac{1}{c_2 y e^{c_1} - 1} dy &= \int dx \\ \frac{\ln(c_2 y e^{c_1} - 1) e^{-c_1}}{c_2} &= c_3 + x \end{aligned}$$

Raising both side to exponential gives

$$e^{\frac{\ln(c_2 y e^{c_1} - 1) e^{-c_1}}{c_2}} = e^{c_3 + x}$$

Which simplifies to

$$(c_2 y e^{c_1} - 1)^{\frac{e^{-c_1}}{c_2}} = e^x c_4$$

Summary

The solution(s) found are the following

$$y = \frac{\left((e^x c_4)^{e^{c_1} c_2} + 1 \right) e^{-c_1}}{c_2} \quad (1)$$

Verification of solutions

$$y = \frac{\left((e^x c_4)^{e^{c_1} c_2} + 1 \right) e^{-c_1}}{c_2}$$

Verified OK.

2.44.2 Maple step by step solution

Let's solve

$$y y'' + (-y' - 1) y' = 0$$

- Highest derivative means the order of the ODE is 2

$$y''$$

- Define new dependent variable u

$$u(x) = y'$$

- Compute y''

$$u'(x) = y''$$

- Use chain rule on the lhs

$$y' \left(\frac{d}{dy} u(y) \right) = y''$$

- Substitute in the definition of u

$$u(y) \left(\frac{d}{dy} u(y) \right) = y''$$

- Make substitutions $y' = u(y)$, $y'' = u(y) \left(\frac{d}{dy} u(y) \right)$ to reduce order of ODE

$$yu(y) \left(\frac{d}{dy} u(y) \right) + (-u(y) - 1) u(y) = 0$$

- Separate variables

$$\frac{\frac{d}{dy} u(y)}{-u(y)-1} = -\frac{1}{y}$$

- Integrate both sides with respect to y

$$\int \frac{\frac{d}{dy} u(y)}{-u(y)-1} dy = \int -\frac{1}{y} dy + c_1$$

- Evaluate integral

$$-\ln(-u(y) - 1) = -\ln(y) + c_1$$

- Solve for $u(y)$

$$u(y) = -\frac{e^{c_1+y}}{e^{c_1}}$$

- Solve 1st ODE for $u(y)$

$$u(y) = -\frac{e^{c_1+y}}{e^{c_1}}$$

- Revert to original variables with substitution $u(y) = y', y = y$

$$y' = -\frac{e^{c_1+y}}{e^{c_1}}$$

- Separate variables

$$\frac{y'}{e^{c_1+y}} = -\frac{1}{e^{c_1}}$$

- Integrate both sides with respect to x

$$\int \frac{y'}{e^{c_1+y}} dx = \int -\frac{1}{e^{c_1}} dx + c_2$$

- Evaluate integral

$$\ln(e^{c_1} + y) = -\frac{x}{e^{c_1}} + c_2$$

- Solve for y

$$y = e^{\frac{e^{c_1} c_2 - x}{e^{c_1}}} - e^{c_1}$$

Maple trace

```
`Methods for second order ODEs:  
--- Trying classification methods ---  
trying 2nd order Liouville  
trying 2nd order WeierstrassP  
trying 2nd order JacobiSN  
differential order: 2; trying a linearization to 3rd order  
trying 2nd order ODE linearizable_by_differentiation  
trying 2nd order, 2 integrating factors of the form mu(x,y)  
trying differential order: 2; missing variables  
, ` -> Computing symmetries using: way = 3  
-> Calling odsolve with the ODE`, (diff(_b(_a), _a))*_b(_a)-_b(_a)*(_b(_a)+1)/_a = 0, _b(_a)  
symmetry methods on request  
, `1st order, trying reduction of order with given symmetries: `[a, 0]
```

✓ Solution by Maple

Time used: 0.078 (sec). Leaf size: 20

```
dsolve(y(x)*diff(y(x),x$2)-diff(y(x),x)^2=diff(y(x),x),y(x), singsol=all)
```

$$y(x) = 0$$
$$y(x) = \frac{e^{c_1(c_2+x)} + 1}{c_1}$$

✓ Solution by Mathematica

Time used: 2.51 (sec). Leaf size: 26

```
DSolve[y[x]*y'[x]-y'[x]^2==y'[x],y[x],x,IncludeSingularSolutions -> True]
```

$$y(x) \rightarrow \frac{1 + e^{c_1(x+c_2)}}{c_1}$$
$$y(x) \rightarrow \text{Indeterminate}$$

3 Chapter 3, SYSTEMS OF DIFFERENTIAL EQUATIONS. Problems page 209

3.1	problem Problem 1	893
3.2	problem Problem 3	900
3.3	problem Problem 4	912
3.4	problem Problem 5	926

3.1 problem Problem 1

- 3.1.1 Solution using Matrix exponential method 893
- 3.1.2 Solution using explicit Eigenvalue and Eigenvector method . . . 894

Internal problem ID [12208]

Internal file name [OUTPUT/10860_Thursday_September_21_2023_05_48_12_AM_8843035/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 3, SYSTEMS OF DIFFERENTIAL EQUATIONS. Problems page 209

Problem number: Problem 1.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$\begin{aligned}x'(t) &= y(t) \\y'(t) &= -x(t)\end{aligned}$$

With initial conditions

$$[x(0) = 0, y(0) = 1]$$

3.1.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \cos(t) & \sin(t) \\ -\sin(t) & \cos(t) \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned}\vec{x}_h(t) &= e^{At}\vec{x}_0 \\ &= \begin{bmatrix} \cos(t) & \sin(t) \\ -\sin(t) & \cos(t) \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} \\ &= \begin{bmatrix} \sin(t) \\ \cos(t) \end{bmatrix}\end{aligned}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

3.1.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det\left(\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\right) = 0$$

Therefore

$$\det\left(\begin{bmatrix} -\lambda & 1 \\ -1 & -\lambda \end{bmatrix}\right) = 0$$

Which gives the characteristic equation

$$\lambda^2 + 1 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = i$$

$$\lambda_2 = -i$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
i	1	complex eigenvalue
$-i$	1	complex eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -i$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} - (-i) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} i & 1 \\ -1 & i \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} i & 1 & 0 \\ -1 & i & 0 \end{array} \right]$$

$$R_2 = -iR_1 + R_2 \implies \left[\begin{array}{cc|c} i & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} i & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = it\}$

Hence the solution is

$$\begin{bmatrix} It \\ t \end{bmatrix} = \begin{bmatrix} it \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} It \\ t \end{bmatrix} = t \begin{bmatrix} i \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} It \\ t \end{bmatrix} = \begin{bmatrix} i \\ 1 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = i$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} - (i) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -i & 1 \\ -1 & -i \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{cc|c} -i & 1 & 0 \\ -1 & -i & 0 \end{array} \right]$$

$$R_2 = iR_1 + R_2 \implies \left[\begin{array}{cc|c} -i & 1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -i & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = -it\}$

Hence the solution is

$$\begin{bmatrix} -It \\ t \end{bmatrix} = \begin{bmatrix} -it \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -It \\ t \end{bmatrix} = t \begin{bmatrix} -i \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -It \\ t \end{bmatrix} = \begin{bmatrix} -i \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
i	1	1	No	$\begin{bmatrix} -i \\ 1 \end{bmatrix}$
$-i$	1	1	No	$\begin{bmatrix} i \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} -ie^{it} \\ e^{it} \end{bmatrix} + c_2 \begin{bmatrix} ie^{-it} \\ e^{-it} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} i(c_2 e^{-it} - c_1 e^{it}) \\ c_1 e^{it} + c_2 e^{-it} \end{bmatrix}$$

Since initial conditions are given, the solution above needs to be updated by solving for the constants of integrations using the given initial conditions

$$\begin{bmatrix} x(0) = 0 \\ y(0) = 1 \end{bmatrix} \tag{1}$$

Substituting initial conditions into the above solution at $t = 0$ gives

$$\begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -i(c_1 - c_2) \\ c_1 + c_2 \end{bmatrix}$$

Solving for the constants of integrations gives

$$\begin{bmatrix} c_1 = \frac{1}{2} \\ c_2 = \frac{1}{2} \end{bmatrix}$$

Substituting these constants back in original solution in Eq. (1) gives

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} i\left(\frac{e^{-it}}{2} - \frac{e^{it}}{2}\right) \\ \frac{e^{it}}{2} + \frac{e^{-it}}{2} \end{bmatrix}$$

The following is the phase plot of the system.




Figure 126: Phase plot

The following are plots of each solution.

✓ Solution by Maple

Time used: 0.0 (sec). Leaf size: 12

```
dsolve([diff(x(t),t) = y(t), diff(y(t),t) = -x(t), x(0) = 0, y(0) = 1], singsol=all)
```

$$\begin{aligned}x(t) &= \sin(t) \\ y(t) &= \cos(t)\end{aligned}$$

✓ Solution by Mathematica

Time used: 0.003 (sec). Leaf size: 31

```
DSolve[{x'[t]==y[t],y'[t]==-x[t]},{},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$\begin{aligned}x(t) &\rightarrow c_1 \cos(t) + c_2 \sin(t) \\ y(t) &\rightarrow c_2 \cos(t) - c_1 \sin(t)\end{aligned}$$

3.2 problem Problem 3

3.2.1	Solution using Matrix exponential method	900
3.2.2	Solution using explicit Eigenvalue and Eigenvector method . . .	902
3.2.3	Maple step by step solution	908

Internal problem ID [12209]

Internal file name [OUTPUT/10861_Thursday_September_21_2023_05_48_13_AM_19860575/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 3, SYSTEMS OF DIFFERENTIAL EQUATIONS. Problems page 209

Problem number: Problem 3.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "system of linear ODEs"

Solve

$$\begin{aligned}x'(t) &= -5x(t) - y(t) + e^t \\y'(t) &= x(t) + 3y(t) + e^{2t}\end{aligned}$$

3.2.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} -5 & -1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} + \begin{bmatrix} e^t \\ e^{2t} \end{bmatrix}$$

Since the system is nonhomogeneous, then the solution is given by

$$\vec{x}(t) = \vec{x}_h(t) + \vec{x}_p(t)$$

Where $\vec{x}_h(t)$ is the homogeneous solution to $\vec{x}'(t) = A\vec{x}(t)$ and $\vec{x}_p(t)$ is a particular solution to $\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$. The particular solution will be found using variation

of parameters method applied to the fundamental matrix. For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{(4\sqrt{15}+15)e^{-(1+\sqrt{15})t}}{30} + \frac{(-4\sqrt{15}+15)e^{-(1+\sqrt{15})t}}{30} & \frac{(-e^{-(1+\sqrt{15})t} + e^{-(1+\sqrt{15})t})\sqrt{15}}{30} \\ -\frac{(e^{-(1+\sqrt{15})t} + e^{-(1+\sqrt{15})t})\sqrt{15}}{30} & \frac{(-4\sqrt{15}+15)e^{-(1+\sqrt{15})t}}{30} + \frac{e^{-(1+\sqrt{15})t}(4\sqrt{15}+15)}{30} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\begin{aligned} \vec{x}_h(t) &= e^{At}\vec{c} \\ &= \begin{bmatrix} \frac{(4\sqrt{15}+15)e^{-(1+\sqrt{15})t}}{30} + \frac{(-4\sqrt{15}+15)e^{-(1+\sqrt{15})t}}{30} & \frac{(-e^{-(1+\sqrt{15})t} + e^{-(1+\sqrt{15})t})\sqrt{15}}{30} \\ -\frac{(e^{-(1+\sqrt{15})t} + e^{-(1+\sqrt{15})t})\sqrt{15}}{30} & \frac{(-4\sqrt{15}+15)e^{-(1+\sqrt{15})t}}{30} + \frac{e^{-(1+\sqrt{15})t}(4\sqrt{15}+15)}{30} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \\ &= \begin{bmatrix} \left(\frac{(4\sqrt{15}+15)e^{-(1+\sqrt{15})t}}{30} + \frac{(-4\sqrt{15}+15)e^{-(1+\sqrt{15})t}}{30} \right) c_1 + \frac{(-e^{-(1+\sqrt{15})t} + e^{-(1+\sqrt{15})t})\sqrt{15}c_2}{30} \\ -\frac{(e^{-(1+\sqrt{15})t} + e^{-(1+\sqrt{15})t})\sqrt{15}c_1}{30} + \left(\frac{(-4\sqrt{15}+15)e^{-(1+\sqrt{15})t}}{30} + \frac{e^{-(1+\sqrt{15})t}(4\sqrt{15}+15)}{30} \right) c_2 \end{bmatrix} \\ &= \begin{bmatrix} \frac{((4c_1+c_2)\sqrt{15}+15c_1)e^{-(1+\sqrt{15})t}}{30} - \frac{2e^{-(1+\sqrt{15})t}((c_1+\frac{c_2}{4})\sqrt{15}-\frac{15c_1}{4})}{15} \\ \frac{((-c_1-4c_2)\sqrt{15}+15c_2)e^{-(1+\sqrt{15})t}}{30} + \frac{e^{-(1+\sqrt{15})t}((c_1+4c_2)\sqrt{15}+15c_2)}{30} \end{bmatrix} \end{aligned}$$

The particular solution given by

$$\vec{x}_p(t) = e^{At} \int e^{-At} \vec{G}(t) dt$$

But

$$\begin{aligned} e^{-At} &= (e^{At})^{-1} \\ &= \begin{bmatrix} \frac{e^{2t}(4\sqrt{15}e^{-(1+\sqrt{15})t} - 4\sqrt{15}e^{-(1+\sqrt{15})t} - 15e^{-(1+\sqrt{15})t} - 15e^{-(1+\sqrt{15})t})}{30} & -\frac{\sqrt{15}e^{2t}(-e^{-(1+\sqrt{15})t} + e^{-(1+\sqrt{15})t})}{30} \\ \frac{\sqrt{15}e^{2t}(-e^{-(1+\sqrt{15})t} + e^{-(1+\sqrt{15})t})}{30} & \frac{e^{2t}(4\sqrt{15}+15)e^{-(1+\sqrt{15})t}}{30} - \frac{2e^{2t}e^{-(1+\sqrt{15})t}}{15} \end{bmatrix} \end{aligned}$$

Hence

$$\begin{aligned} \vec{x}_p(t) &= \begin{bmatrix} \frac{(4\sqrt{15}+15)e^{-(1+\sqrt{15})t}}{30} + \frac{(-4\sqrt{15}+15)e^{(-1+\sqrt{15})t}}{30} & \frac{(-e^{(-1+\sqrt{15})t} + e^{-(1+\sqrt{15})t})\sqrt{15}}{30} \\ -\frac{(e^{(-1+\sqrt{15})t} + e^{-(1+\sqrt{15})t})\sqrt{15}}{30} & \frac{(-4\sqrt{15}+15)e^{-(1+\sqrt{15})t}}{30} + \frac{e^{(-1+\sqrt{15})t}(4\sqrt{15}+15)}{30} \end{bmatrix} \int \begin{bmatrix} e \\ -35e \end{bmatrix} \\ &= \begin{bmatrix} \frac{(4\sqrt{15}+15)e^{-(1+\sqrt{15})t}}{30} + \frac{(-4\sqrt{15}+15)e^{(-1+\sqrt{15})t}}{30} & \frac{(-e^{(-1+\sqrt{15})t} + e^{-(1+\sqrt{15})t})\sqrt{15}}{30} \\ -\frac{(e^{(-1+\sqrt{15})t} + e^{-(1+\sqrt{15})t})\sqrt{15}}{30} & \frac{(-4\sqrt{15}+15)e^{-(1+\sqrt{15})t}}{30} + \frac{e^{(-1+\sqrt{15})t}(4\sqrt{15}+15)}{30} \end{bmatrix} \begin{bmatrix} 5 \\ -35 \end{bmatrix} \\ &= \begin{bmatrix} \frac{e^{2t}}{6} + \frac{2e^t}{11} \\ -\frac{7e^{2t}}{6} - \frac{e^t}{11} \end{bmatrix} \end{aligned}$$

Hence the complete solution is

$$\begin{aligned} \vec{x}(t) &= \vec{x}_h(t) + \vec{x}_p(t) \\ &= \begin{bmatrix} \frac{((4c_1+c_2)\sqrt{15}+15c_1)e^{-(1+\sqrt{15})t}}{30} + \frac{((-4c_1-c_2)\sqrt{15}+15c_1)e^{(-1+\sqrt{15})t}}{30} + \frac{2e^t}{11} + \frac{e^{2t}}{6} \\ \frac{((-c_1-4c_2)\sqrt{15}+15c_2)e^{-(1+\sqrt{15})t}}{30} + \frac{e^{(-1+\sqrt{15})t}((c_1+4c_2)\sqrt{15}+15c_2)}{30} - \frac{e^t}{11} - \frac{7e^{2t}}{6} \end{bmatrix} \end{aligned}$$

3.2.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \end{bmatrix} = \begin{bmatrix} -5 & -1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} + \begin{bmatrix} e^t \\ e^{2t} \end{bmatrix}$$

Since the system is nonhomogeneous, then the solution is given by

$$\vec{x}(t) = \vec{x}_h(t) + \vec{x}_p(t)$$

Where $\vec{x}_h(t)$ is the homogeneous solution to $\vec{x}'(t) = A\vec{x}(t)$ and $\vec{x}_p(t)$ is a particular solution to $\vec{x}'(t) = A\vec{x}(t) + \vec{G}(t)$. The particular solution will be found using variation of parameters method applied to the fundamental matrix.

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{bmatrix} -5 & -1 \\ 1 & 3 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} -5 - \lambda & -1 \\ 1 & 3 - \lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^2 + 2\lambda - 14 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = -1 + \sqrt{15}$$

$$\lambda_2 = -1 - \sqrt{15}$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
$-1 + \sqrt{15}$	1	real eigenvalue
$-1 - \sqrt{15}$	1	real eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = -1 - \sqrt{15}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} -5 & -1 \\ 1 & 3 \end{bmatrix} - (-1 - \sqrt{15}) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -4 + \sqrt{15} & -1 \\ 1 & 4 + \sqrt{15} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\begin{bmatrix} -4 + \sqrt{15} & -1 & 0 \\ 1 & 4 + \sqrt{15} & 0 \end{bmatrix}$$

$$R_2 = R_2 - \frac{R_1}{-4 + \sqrt{15}} \implies \begin{bmatrix} -4 + \sqrt{15} & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -4 + \sqrt{15} & -1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{ v_1 = \frac{t}{-4 + \sqrt{15}} \right\}$

Hence the solution is

$$\begin{bmatrix} \frac{t}{-4 + \sqrt{15}} \\ t \end{bmatrix} = \begin{bmatrix} \frac{t}{-4 + \sqrt{15}} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{t}{-4 + \sqrt{15}} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{1}{-4 + \sqrt{15}} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{1}{-4 + \sqrt{15}} \\ 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{-4 + \sqrt{15}} \\ 1 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = -1 + \sqrt{15}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} -5 & -1 \\ 1 & 3 \end{bmatrix} - (-1 + \sqrt{15}) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -4 - \sqrt{15} & -1 \\ 1 & 4 - \sqrt{15} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\begin{bmatrix} -4 - \sqrt{15} & -1 & 0 \\ 1 & 4 - \sqrt{15} & 0 \end{bmatrix}$$

$$R_2 = R_2 - \frac{R_1}{-4 - \sqrt{15}} \implies \begin{bmatrix} -4 - \sqrt{15} & -1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -4 - \sqrt{15} & -1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_2\}$ and the leading variables are $\{v_1\}$. Let $v_2 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{ v_1 = -\frac{t}{4+\sqrt{15}} \right\}$

Hence the solution is

$$\begin{bmatrix} -\frac{t}{4+\sqrt{15}} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{t}{4+\sqrt{15}} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -\frac{t}{4+\sqrt{15}} \\ t \end{bmatrix} = t \begin{bmatrix} -\frac{1}{4+\sqrt{15}} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -\frac{1}{4+\sqrt{15}} \\ 1 \end{bmatrix} = \begin{bmatrix} -\frac{1}{4+\sqrt{15}} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} -\frac{1}{4+\sqrt{15}} \\ 1 \end{bmatrix} = \begin{bmatrix} -\frac{1}{4+\sqrt{15}} \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
$-1 + \sqrt{15}$	1	1	No	$\begin{bmatrix} -\frac{1}{4+\sqrt{15}} \\ 1 \end{bmatrix}$
$-1 - \sqrt{15}$	1	1	No	$\begin{bmatrix} -\frac{1}{4-\sqrt{15}} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue $-1 + \sqrt{15}$ is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_1(t) &= \vec{v}_1 e^{(-1+\sqrt{15})t} \\ &= \begin{bmatrix} -\frac{1}{4+\sqrt{15}} \\ 1 \end{bmatrix} e^{(-1+\sqrt{15})t}\end{aligned}$$

Since eigenvalue $-1 - \sqrt{15}$ is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned}\vec{x}_2(t) &= \vec{v}_2 e^{(-1-\sqrt{15})t} \\ &= \begin{bmatrix} -\frac{1}{4-\sqrt{15}} \\ 1 \end{bmatrix} e^{(-1-\sqrt{15})t}\end{aligned}$$

Therefore the homogeneous solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = c_1 \begin{bmatrix} -\frac{e^{(-1+\sqrt{15})t}}{4+\sqrt{15}} \\ e^{(-1+\sqrt{15})t} \end{bmatrix} + c_2 \begin{bmatrix} -\frac{e^{(-1-\sqrt{15})t}}{4-\sqrt{15}} \\ e^{(-1-\sqrt{15})t} \end{bmatrix}$$

Now that we found homogeneous solution above, we need to find a particular solution $\vec{x}_p(t)$. We will use Variation of parameters. The fundamental matrix is

$$\Phi = \begin{bmatrix} \vec{x}_1 & \vec{x}_2 & \cdots \end{bmatrix}$$

Where \vec{x}_i are the solution basis found above. Therefore the fundamental matrix is

$$\Phi(t) = \begin{bmatrix} -\frac{e^{(-1+\sqrt{15})t}}{4+\sqrt{15}} & -\frac{e^{(-1-\sqrt{15})t}}{4-\sqrt{15}} \\ e^{(-1+\sqrt{15})t} & e^{(-1-\sqrt{15})t} \end{bmatrix}$$

The particular solution is then given by

$$\vec{x}_p(t) = \Phi \int \Phi^{-1} \vec{G}(t) dt$$

But

$$\Phi^{-1} = \begin{bmatrix} \frac{\sqrt{15}e^{-(1+\sqrt{15})t}}{30} & \frac{\sqrt{15}(4+\sqrt{15})e^{-(1+\sqrt{15})t}}{30} \\ -\frac{e^{(1+\sqrt{15})t}\sqrt{15}}{30} & \frac{e^{(1+\sqrt{15})t}\sqrt{15}(-4+\sqrt{15})}{30} \end{bmatrix}$$

Hence

$$\begin{aligned} \vec{x}_p(t) &= \begin{bmatrix} -\frac{e^{(-1+\sqrt{15})t}}{4+\sqrt{15}} & -\frac{e^{(-1-\sqrt{15})t}}{4-\sqrt{15}} \\ e^{(-1+\sqrt{15})t} & e^{(-1-\sqrt{15})t} \end{bmatrix} \int \begin{bmatrix} \frac{\sqrt{15}e^{-(1+\sqrt{15})t}}{30} & \frac{\sqrt{15}(4+\sqrt{15})e^{-(1+\sqrt{15})t}}{30} \\ -\frac{e^{(1+\sqrt{15})t}\sqrt{15}}{30} & \frac{e^{(1+\sqrt{15})t}\sqrt{15}(-4+\sqrt{15})}{30} \end{bmatrix} \begin{bmatrix} e^t \\ e^{2t} \end{bmatrix} dt \\ &= \begin{bmatrix} -\frac{e^{(-1+\sqrt{15})t}}{4+\sqrt{15}} & -\frac{e^{(-1-\sqrt{15})t}}{4-\sqrt{15}} \\ e^{(-1+\sqrt{15})t} & e^{(-1-\sqrt{15})t} \end{bmatrix} \int \begin{bmatrix} \frac{e^{-t(-3+\sqrt{15})}(4\sqrt{15}+15)}{30} + \frac{\sqrt{15}e^{-t(-2+\sqrt{15})}}{30} \\ \frac{(-4\sqrt{15}+15)e^{t(3+\sqrt{15})}}{30} - \frac{\sqrt{15}e^{t(2+\sqrt{15})}}{30} \end{bmatrix} dt \\ &= \begin{bmatrix} -\frac{e^{(-1+\sqrt{15})t}}{4+\sqrt{15}} & -\frac{e^{(-1-\sqrt{15})t}}{4-\sqrt{15}} \\ e^{(-1+\sqrt{15})t} & e^{(-1-\sqrt{15})t} \end{bmatrix} \begin{bmatrix} -\frac{\sqrt{15}(7e^{-t(-3+\sqrt{15})} + \sqrt{15}e^{-t(-2+\sqrt{15})} + 2\sqrt{15}e^{-t(-3+\sqrt{15})} - 3e^{-t(-2+\sqrt{15})})}{30(-2+\sqrt{15})(-3+\sqrt{15})} \\ -\frac{((3+\sqrt{15})e^{t(2+\sqrt{15})} + e^{t(3+\sqrt{15})}(2\sqrt{15}-7))\sqrt{15}}{30(3+\sqrt{15})(2+\sqrt{15})} \end{bmatrix} \\ &= \begin{bmatrix} \frac{e^{2t}}{6} + \frac{2e^t}{11} \\ -\frac{7e^{2t}}{6} - \frac{e^t}{11} \end{bmatrix} \end{aligned}$$

Now that we found particular solution, the final solution is

$$\begin{aligned} \vec{x}(t) &= \vec{x}_h(t) + \vec{x}_p(t) \\ \begin{bmatrix} x(t) \\ y(t) \end{bmatrix} &= \begin{bmatrix} -\frac{c_1 e^{(-1+\sqrt{15})t}}{4+\sqrt{15}} \\ c_1 e^{(-1+\sqrt{15})t} \end{bmatrix} + \begin{bmatrix} -\frac{c_2 e^{(-1-\sqrt{15})t}}{4-\sqrt{15}} \\ c_2 e^{(-1-\sqrt{15})t} \end{bmatrix} + \begin{bmatrix} \frac{e^{2t}}{6} + \frac{2e^t}{11} \\ -\frac{7e^{2t}}{6} - \frac{e^t}{11} \end{bmatrix} \end{aligned}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} -c_2(4 + \sqrt{15})e^{-(1+\sqrt{15})t} + c_1(-4 + \sqrt{15})e^{(-1+\sqrt{15})t} + \frac{2e^t}{11} + \frac{e^{2t}}{6} \\ c_1e^{(-1+\sqrt{15})t} + c_2e^{-(1+\sqrt{15})t} - \frac{7e^{2t}}{6} - \frac{e^t}{11} \end{bmatrix}$$

3.2.3 Maple step by step solution

Let's solve

$$\left[x'(t) = -5x(t) - y(t) + e^t, y'(t) = x(t) + 3y(t) + (e^t)^2 \right]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} -5 & -1 \\ 1 & 3 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} e^t \\ (e^t)^2 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} -5 & -1 \\ 1 & 3 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} e^t \\ (e^t)^2 \end{bmatrix}$$

- Define the forcing function

$$\vec{f}(t) = \begin{bmatrix} e^t \\ (e^t)^2 \end{bmatrix}$$

- Define the coefficient matrix

$$A = \begin{bmatrix} -5 & -1 \\ 1 & 3 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t) + \vec{f}$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[-1 - \sqrt{15}, \begin{bmatrix} -\frac{1}{4-\sqrt{15}} \\ 1 \end{bmatrix} \right], \left[-1 + \sqrt{15}, \begin{bmatrix} -\frac{1}{4+\sqrt{15}} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[-1 - \sqrt{15}, \begin{bmatrix} -\frac{1}{4-\sqrt{15}} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^{(-1-\sqrt{15})t} \cdot \begin{bmatrix} -\frac{1}{4-\sqrt{15}} \\ 1 \end{bmatrix}$$

- Consider eigenpair

$$\left[-1 + \sqrt{15}, \begin{bmatrix} -\frac{1}{4+\sqrt{15}} \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_2 = e^{(-1+\sqrt{15})t} \cdot \begin{bmatrix} -\frac{1}{4+\sqrt{15}} \\ 1 \end{bmatrix}$$

- General solution of the system of ODEs can be written in terms of the particular solution $\vec{x}_p(t)$

$$\vec{x}(t) = c_1 \vec{x}_1 + c_2 \vec{x}_2 + \vec{x}_p(t)$$

- Fundamental matrix

- Let $\phi(t)$ be the matrix whose columns are the independent solutions of the homogeneous system

$$\phi(t) = \begin{bmatrix} -\frac{e^{(-1-\sqrt{15})t}}{4-\sqrt{15}} & -\frac{e^{(-1+\sqrt{15})t}}{4+\sqrt{15}} \\ e^{(-1-\sqrt{15})t} & e^{(-1+\sqrt{15})t} \end{bmatrix}$$

- The fundamental matrix, $\Phi(t)$ is a normalized version of $\phi(t)$ satisfying $\Phi(0) = I$ where I is the identity matrix

$$\Phi(t) = \phi(t) \cdot \frac{1}{\phi(0)}$$

- Substitute the value of $\phi(t)$ and $\phi(0)$

$$\Phi(t) = \begin{bmatrix} -\frac{e^{(-1-\sqrt{15})t}}{4-\sqrt{15}} & -\frac{e^{(-1+\sqrt{15})t}}{4+\sqrt{15}} \\ e^{(-1-\sqrt{15})t} & e^{(-1+\sqrt{15})t} \end{bmatrix} \cdot \frac{1}{\begin{bmatrix} -\frac{1}{4-\sqrt{15}} & -\frac{1}{4+\sqrt{15}} \\ 1 & 1 \end{bmatrix}}$$

- Evaluate and simplify to get the fundamental matrix

$$\Phi(t) = \begin{bmatrix} \frac{\sqrt{15} \left((4+\sqrt{15})e^{-(1+\sqrt{15})t} + e^{(-1+\sqrt{15})t}(-4+\sqrt{15}) \right)}{30} & \frac{\left(-e^{(-1+\sqrt{15})t} + e^{-(1+\sqrt{15})t} \right) \sqrt{15}}{30} \\ -\frac{\left(-e^{(-1+\sqrt{15})t} + e^{-(1+\sqrt{15})t} \right) \sqrt{15}}{30} & \frac{(-4\sqrt{15}+15)e^{-(1+\sqrt{15})t}}{30} + \frac{e^{(-1+\sqrt{15})t}(4\sqrt{15}+15)}{30} \end{bmatrix}$$

□

Find a particular solution of the system of ODEs using variation of parameters

- Let the particular solution be the fundamental matrix multiplied by $\vec{v}(t)$ and solve for $\vec{v}(t)$

$$\vec{x}_p(t) = \Phi(t) \cdot \vec{v}(t)$$

- Take the derivative of the particular solution

$$\vec{x}'_p(t) = \Phi'(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t)$$

- Substitute particular solution and its derivative into the system of ODEs

$$\Phi'(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t) = A \cdot \Phi(t) \cdot \vec{v}(t) + \vec{f}(t)$$

- The fundamental matrix has columns that are solutions to the homogeneous system so its derivative is $A \cdot \Phi(t) \cdot \vec{v}(t)$

$$A \cdot \Phi(t) \cdot \vec{v}(t) + \Phi(t) \cdot \vec{v}'(t) = A \cdot \Phi(t) \cdot \vec{v}(t) + \vec{f}(t)$$

- Cancel like terms

$$\Phi(t) \cdot \vec{v}'(t) = \vec{f}(t)$$

- Multiply by the inverse of the fundamental matrix

$$\vec{v}'(t) = \frac{1}{\Phi(t)} \cdot \vec{f}(t)$$

- Integrate to solve for $\vec{v}(t)$

$$\vec{v}(t) = \int_0^t \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds$$

- Plug $\vec{v}(t)$ into the equation for the particular solution

$$\vec{x}_p(t) = \Phi(t) \cdot \left(\int_0^t \frac{1}{\Phi(s)} \cdot \vec{f}(s) ds \right)$$

- Plug in the fundamental matrix and the forcing function and compute

$$\vec{x}_p(t) = \begin{bmatrix} \frac{(-3\sqrt{15}-115)e^{-(1+\sqrt{15})t}}{660} + \frac{(3\sqrt{15}-115)e^{(-1+\sqrt{15})t}}{660} + \frac{2e^t}{11} + \frac{e^{2t}}{6} \\ \frac{(-103\sqrt{15}+415)e^{-(1+\sqrt{15})t}}{660} + \frac{(103\sqrt{15}+415)e^{(-1+\sqrt{15})t}}{660} - \frac{e^t}{11} - \frac{7e^{2t}}{6} \end{bmatrix}$$

- Plug particular solution back into general solution

$$\vec{x}(t) = c_1 \vec{x}_1 + c_2 \vec{x}_2 + \begin{bmatrix} \frac{(-3\sqrt{15}-115)e^{-(1+\sqrt{15})t}}{660} + \frac{(3\sqrt{15}-115)e^{(-1+\sqrt{15})t}}{660} + \frac{2e^t}{11} + \frac{e^{2t}}{6} \\ \frac{(-103\sqrt{15}+415)e^{-(1+\sqrt{15})t}}{660} + \frac{(103\sqrt{15}+415)e^{(-1+\sqrt{15})t}}{660} - \frac{e^t}{11} - \frac{7e^{2t}}{6} \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} \frac{((-660c_1-3)\sqrt{15}-2640c_1-115)e^{-(1+\sqrt{15})t}}{660} + \frac{((660c_2+3)\sqrt{15}-2640c_2-115)e^{(-1+\sqrt{15})t}}{660} + \frac{2e^t}{11} + \frac{e^{2t}}{6} \\ \frac{(660c_1-103\sqrt{15}+415)e^{-(1+\sqrt{15})t}}{660} + \frac{(660c_2+103\sqrt{15}+415)e^{(-1+\sqrt{15})t}}{660} - \frac{e^t}{11} - \frac{7e^{2t}}{6} \end{bmatrix}$$

- Solution to the system of ODEs

$$\begin{cases} x(t) = \frac{((-660c_1-3)\sqrt{15}-2640c_1-115)e^{-(1+\sqrt{15})t}}{660} + \frac{((660c_2+3)\sqrt{15}-2640c_2-115)e^{(-1+\sqrt{15})t}}{660} + \frac{2e^t}{11} + \frac{e^{2t}}{6}, y(t) = \end{cases}$$

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 102

```
dsolve([diff(x(t),t)+5*x(t)+y(t)=exp(t),diff(y(t),t)-x(t)-3*y(t)=exp(2*t)],singsol=all)
```

$$x(t) = e^{(\sqrt{15}-1)t} c_2 + e^{-(1+\sqrt{15})t} c_1 + \frac{e^{2t}}{6} + \frac{2e^t}{11}$$

$$y(t) = -e^{(\sqrt{15}-1)t} c_2 \sqrt{15} + e^{-(1+\sqrt{15})t} c_1 \sqrt{15} - 4e^{(\sqrt{15}-1)t} c_2 - 4e^{-(1+\sqrt{15})t} c_1 - \frac{e^t}{11} - \frac{7e^{2t}}{6}$$

✓ Solution by Mathematica

Time used: 4.39 (sec). Leaf size: 206

```
DSolve[{x'[t]+5*x[t]+y[t]==Exp[t],y'[t]-x[t]-3*y[t]==Exp[2*t]},{x[t],y[t]},t,IncludeSingular
```

$$x(t) \rightarrow \frac{1}{330} e^{-((1+\sqrt{15})t)} \left(60e^{(2+\sqrt{15})t} + 55e^{(3+\sqrt{15})t} - 11 \left((4\sqrt{15} - 15) c_1 + \sqrt{15} c_2 \right) e^{2\sqrt{15}t} + 11 \left((15 + 4\sqrt{15}) c_1 + \sqrt{15} c_2 \right) \right)$$

$$y(t) \rightarrow -\frac{1}{330} e^{-((1+\sqrt{15})t)} \left(30e^{(2+\sqrt{15})t} + 385e^{(3+\sqrt{15})t} - 11 \left(\sqrt{15} c_1 + (15 + 4\sqrt{15}) c_2 \right) e^{2\sqrt{15}t} + 11 \left(\sqrt{15} c_1 + (4\sqrt{15} - 15) c_2 \right) \right)$$

3.3 problem Problem 4

3.3.1	Solution using Matrix exponential method	912
3.3.2	Solution using explicit Eigenvalue and Eigenvector method . . .	913
3.3.3	Maple step by step solution	921

Internal problem ID [12210]

Internal file name [OUTPUT/10862_Thursday_September_21_2023_05_48_14_AM_31237680/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 3, SYSTEMS OF DIFFERENTIAL EQUATIONS. Problems page 209

Problem number: Problem 4.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**"

Solve

$$x'(t) = y(t)$$

$$y'(t) = z(t)$$

$$z'(t) = x(t)$$

3.3.1 Solution using Matrix exponential method

In this method, we will assume we have found the matrix exponential e^{At} already. There are different methods to determine this but will not be shown here. This is a system of linear ODE's given as

$$\vec{x}'(t) = A \vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \\ z'(t) \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix}$$

For the above matrix A , the matrix exponential can be found to be

$$e^{At} = \begin{bmatrix} \frac{e^t}{3} + \frac{2e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} & -\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^t}{3} & -\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} - \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} \\ -\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} - \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^t}{3} & \frac{e^t}{3} + \frac{2e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} & -\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} \\ -\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^t}{3} & -\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} - \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^t}{3} & \frac{e^t}{3} + \frac{2e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} \end{bmatrix}$$

Therefore the homogeneous solution is

$$\vec{x}_h(t) = e^{At} \vec{c}$$

$$= \begin{bmatrix} \frac{e^t}{3} + \frac{2e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} & -\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^t}{3} & -\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} - \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} \\ -\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} - \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^t}{3} & \frac{e^t}{3} + \frac{2e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} & -\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} \\ -\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^t}{3} & -\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} - \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^t}{3} & \frac{e^t}{3} + \frac{2e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} \end{bmatrix} \begin{bmatrix} c_1 \\ c_2 \\ c_3 \end{bmatrix}$$

$$= \begin{bmatrix} \left(\frac{e^t}{3} + \frac{2e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3}\right) c_1 + \left(-\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^t}{3}\right) c_2 + \left(-\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} - \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3}\right) c_3 \\ \left(-\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} - \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^t}{3}\right) c_1 + \left(\frac{e^t}{3} + \frac{2e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3}\right) c_2 + \left(-\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3}\right) c_3 \\ \left(-\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^t}{3}\right) c_1 + \left(-\frac{e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} - \frac{e^{-\frac{t}{2}} \sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^t}{3}\right) c_2 + \left(\frac{e^t}{3} + \frac{2e^{-\frac{t}{2}} \cos\left(\frac{\sqrt{3}t}{2}\right)}{3}\right) c_3 \end{bmatrix}$$

$$= \begin{bmatrix} \frac{2e^{-\frac{t}{2}}(c_1 - \frac{c_2}{2} - \frac{c_3}{2}) \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{\sqrt{3}e^{-\frac{t}{2}}(-c_3 + c_2) \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^t(c_1 + c_2 + c_3)}{3} \\ -\frac{e^{-\frac{t}{2}}(c_1 - 2c_2 + c_3) \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} - \frac{\sqrt{3}e^{-\frac{t}{2}}(-c_3 + c_1) \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^t(c_1 + c_2 + c_3)}{3} \\ -\frac{e^{-\frac{t}{2}}(c_1 + c_2 - 2c_3) \cos\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{\sqrt{3}e^{-\frac{t}{2}}(c_1 - c_2) \sin\left(\frac{\sqrt{3}t}{2}\right)}{3} + \frac{e^t(c_1 + c_2 + c_3)}{3} \end{bmatrix}$$

Since no forcing function is given, then the final solution is $\vec{x}_h(t)$ above.

3.3.2 Solution using explicit Eigenvalue and Eigenvector method

This is a system of linear ODE's given as

$$\vec{x}'(t) = A\vec{x}(t)$$

Or

$$\begin{bmatrix} x'(t) \\ y'(t) \\ z'(t) \end{bmatrix} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix}$$

The first step is find the homogeneous solution. We start by finding the eigenvalues of A . This is done by solving the following equation for the eigenvalues λ

$$\det(A - \lambda I) = 0$$

Expanding gives

$$\det \left(\begin{pmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} - \lambda \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) = 0$$

Therefore

$$\det \left(\begin{bmatrix} -\lambda & 1 & 0 \\ 0 & -\lambda & 1 \\ 1 & 0 & -\lambda \end{bmatrix} \right) = 0$$

Which gives the characteristic equation

$$\lambda^3 - 1 = 0$$

The roots of the above are the eigenvalues.

$$\lambda_1 = 1$$

$$\lambda_2 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$$

$$\lambda_3 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$$

This table summarises the above result

eigenvalue	algebraic multiplicity	type of eigenvalue
1	1	real eigenvalue
$-\frac{1}{2} - \frac{i\sqrt{3}}{2}$	1	complex eigenvalue
$-\frac{1}{2} + \frac{i\sqrt{3}}{2}$	1	complex eigenvalue

Now the eigenvector for each eigenvalue are found.

Considering the eigenvalue $\lambda_1 = 1$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} - (1) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \\ 1 & 0 & -1 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{ccc|c} -1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 1 & 0 & -1 & 0 \end{array} \right]$$

$$R_3 = R_3 + R_1 \implies \left[\begin{array}{ccc|c} -1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 1 & -1 & 0 \end{array} \right]$$

$$R_3 = R_3 + R_2 \implies \left[\begin{array}{ccc|c} -1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_3\}$ and the leading variables are $\{v_1, v_2\}$. Let $v_3 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\{v_1 = t, v_2 = t\}$

Hence the solution is

$$\begin{bmatrix} t \\ t \\ t \end{bmatrix} = \begin{bmatrix} t \\ t \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} t \\ t \\ t \end{bmatrix} = t \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} t \\ t \\ t \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Considering the eigenvalue $\lambda_2 = -\frac{1}{2} - \frac{i\sqrt{3}}{2}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} - \left(-\frac{1}{2} - \frac{i\sqrt{3}}{2} \right) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} \frac{1}{2} + \frac{i\sqrt{3}}{2} & 1 & 0 \\ 0 & \frac{1}{2} + \frac{i\sqrt{3}}{2} & 1 \\ 1 & 0 & \frac{1}{2} + \frac{i\sqrt{3}}{2} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{ccc|c} \frac{1}{2} + \frac{i\sqrt{3}}{2} & 1 & 0 & 0 \\ 0 & \frac{1}{2} + \frac{i\sqrt{3}}{2} & 1 & 0 \\ 1 & 0 & \frac{1}{2} + \frac{i\sqrt{3}}{2} & 0 \end{array} \right]$$

$$R_3 = R_3 - \frac{R_1}{\frac{1}{2} + \frac{i\sqrt{3}}{2}} \implies \left[\begin{array}{ccc|c} \frac{1}{2} + \frac{i\sqrt{3}}{2} & 1 & 0 & 0 \\ 0 & \frac{1}{2} + \frac{i\sqrt{3}}{2} & 1 & 0 \\ 0 & -\frac{2}{1+i\sqrt{3}} & \frac{1}{2} + \frac{i\sqrt{3}}{2} & 0 \end{array} \right]$$

$$R_3 = R_3 + \frac{2R_2}{(1+i\sqrt{3})\left(\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)} \Rightarrow \left[\begin{array}{ccc|c} \frac{1}{2} + \frac{i\sqrt{3}}{2} & 1 & 0 & 0 \\ 0 & \frac{1}{2} + \frac{i\sqrt{3}}{2} & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} \frac{1}{2} + \frac{i\sqrt{3}}{2} & 1 & 0 \\ 0 & \frac{1}{2} + \frac{i\sqrt{3}}{2} & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_3\}$ and the leading variables are $\{v_1, v_2\}$. Let $v_3 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{ v_1 = \frac{2t}{i\sqrt{3}-1}, v_2 = -\frac{2t}{1+i\sqrt{3}} \right\}$

Hence the solution is

$$\begin{bmatrix} \frac{2t}{i\sqrt{3}-1} \\ -\frac{2t}{1+i\sqrt{3}} \\ t \end{bmatrix} = \begin{bmatrix} \frac{2t}{i\sqrt{3}-1} \\ -\frac{2t}{1+i\sqrt{3}} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} \frac{2t}{i\sqrt{3}-1} \\ -\frac{2t}{1+i\sqrt{3}} \\ t \end{bmatrix} = t \begin{bmatrix} \frac{2}{i\sqrt{3}-1} \\ -\frac{2}{1+i\sqrt{3}} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} \frac{2t}{i\sqrt{3}-1} \\ -\frac{2t}{1+i\sqrt{3}} \\ t \end{bmatrix} = \begin{bmatrix} \frac{2}{i\sqrt{3}-1} \\ -\frac{2}{1+i\sqrt{3}} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} \frac{2t}{i\sqrt{3}-1} \\ -\frac{2t}{1+i\sqrt{3}} \\ t \end{bmatrix} = \begin{bmatrix} \frac{2}{i\sqrt{3}-1} \\ -\frac{2}{1+i\sqrt{3}} \\ 1 \end{bmatrix}$$

Considering the eigenvalue $\lambda_3 = -\frac{1}{2} + \frac{i\sqrt{3}}{2}$

We need to solve $A\vec{v} = \lambda\vec{v}$ or $(A - \lambda I)\vec{v} = \vec{0}$ which becomes

$$\left(\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} - \left(-\frac{1}{2} + \frac{i\sqrt{3}}{2} \right) \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \right) \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} \frac{1}{2} - \frac{i\sqrt{3}}{2} & 1 & 0 \\ 0 & \frac{1}{2} - \frac{i\sqrt{3}}{2} & 1 \\ 1 & 0 & \frac{1}{2} - \frac{i\sqrt{3}}{2} \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

Now forward elimination is applied to solve for the eigenvector \vec{v} . The augmented matrix is

$$\left[\begin{array}{ccc|c} \frac{1}{2} - \frac{i\sqrt{3}}{2} & 1 & 0 & 0 \\ 0 & \frac{1}{2} - \frac{i\sqrt{3}}{2} & 1 & 0 \\ 1 & 0 & \frac{1}{2} - \frac{i\sqrt{3}}{2} & 0 \end{array} \right]$$

$$R_3 = R_3 - \frac{R_1}{\frac{1}{2} - \frac{i\sqrt{3}}{2}} \Rightarrow \left[\begin{array}{ccc|c} \frac{1}{2} - \frac{i\sqrt{3}}{2} & 1 & 0 & 0 \\ 0 & \frac{1}{2} - \frac{i\sqrt{3}}{2} & 1 & 0 \\ 0 & \frac{2}{i\sqrt{3}-1} & \frac{1}{2} - \frac{i\sqrt{3}}{2} & 0 \end{array} \right]$$

$$R_3 = R_3 - \frac{2R_2}{(i\sqrt{3}-1)\left(\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)} \Rightarrow \left[\begin{array}{ccc|c} \frac{1}{2} - \frac{i\sqrt{3}}{2} & 1 & 0 & 0 \\ 0 & \frac{1}{2} - \frac{i\sqrt{3}}{2} & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

Therefore the system in Echelon form is

$$\begin{bmatrix} \frac{1}{2} - \frac{i\sqrt{3}}{2} & 1 & 0 \\ 0 & \frac{1}{2} - \frac{i\sqrt{3}}{2} & 1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \\ v_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

The free variables are $\{v_3\}$ and the leading variables are $\{v_1, v_2\}$. Let $v_3 = t$. Now we start back substitution. Solving the above equation for the leading variables in terms of free variables gives equation $\left\{ v_1 = -\frac{2t}{1+i\sqrt{3}}, v_2 = \frac{2t}{i\sqrt{3}-1} \right\}$

Hence the solution is

$$\begin{bmatrix} -\frac{2t}{1+i\sqrt{3}} \\ \frac{2t}{i\sqrt{3}-1} \\ t \end{bmatrix} = \begin{bmatrix} -\frac{2t}{1+i\sqrt{3}} \\ \frac{2t}{i\sqrt{3}-1} \\ t \end{bmatrix}$$

Since there is one free Variable, we have found one eigenvector associated with this eigenvalue. The above can be written as

$$\begin{bmatrix} -\frac{2t}{1+i\sqrt{3}} \\ \frac{2t}{i\sqrt{3}-1} \\ t \end{bmatrix} = t \begin{bmatrix} -\frac{2}{1+i\sqrt{3}} \\ \frac{2}{i\sqrt{3}-1} \\ 1 \end{bmatrix}$$

Let $t = 1$ the eigenvector becomes

$$\begin{bmatrix} -\frac{2}{1+i\sqrt{3}} \\ \frac{2}{i\sqrt{3}-1} \\ 1 \end{bmatrix} = \begin{bmatrix} -\frac{2}{1+i\sqrt{3}} \\ \frac{2}{i\sqrt{3}-1} \\ 1 \end{bmatrix}$$

Which is normalized to

$$\begin{bmatrix} -\frac{2}{1+i\sqrt{3}} \\ \frac{2}{i\sqrt{3}-1} \\ 1 \end{bmatrix} = \begin{bmatrix} -\frac{2}{1+i\sqrt{3}} \\ \frac{2}{i\sqrt{3}-1} \\ 1 \end{bmatrix}$$

The following table gives a summary of this result. It shows for each eigenvalue the algebraic multiplicity m , and its geometric multiplicity k and the eigenvectors associated with the eigenvalue. If $m > k$ then the eigenvalue is defective which means the number of normal linearly independent eigenvectors associated with this eigenvalue (called the geometric multiplicity k) does not equal the algebraic multiplicity m , and we need to determine an additional $m - k$ generalized eigenvectors for this eigenvalue.

eigenvalue	multiplicity		defective?	eigenvectors
	algebraic m	geometric k		
1	1	1	No	$\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$
$-\frac{1}{2} + \frac{i\sqrt{3}}{2}$	1	1	No	$\begin{bmatrix} \frac{1}{\left(-\frac{1}{2} + \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} + \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix}$
$-\frac{1}{2} - \frac{i\sqrt{3}}{2}$	1	1	No	$\begin{bmatrix} \frac{1}{\left(-\frac{1}{2} - \frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix}$

Now that we found the eigenvalues and associated eigenvectors, we will go over each eigenvalue and generate the solution basis. The only problem we need to take care of is if the eigenvalue is defective. Since eigenvalue 1 is real and distinct then the corresponding eigenvector solution is

$$\begin{aligned} \vec{x}_1(t) &= \vec{v}_1 e^t \\ &= \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} e^t \end{aligned}$$

Therefore the final solution is

$$\vec{x}_h(t) = c_1 \vec{x}_1(t) + c_2 \vec{x}_2(t) + c_3 \vec{x}_3(t)$$

Which is written as

$$\begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix} = c_1 \begin{bmatrix} e^t \\ e^t \\ e^t \end{bmatrix} + c_2 \begin{bmatrix} \frac{e^{(-\frac{1}{2} + \frac{i\sqrt{3}}{2})t}}{(-\frac{1}{2} + \frac{i\sqrt{3}}{2})^2} \\ \frac{e^{(-\frac{1}{2} + \frac{i\sqrt{3}}{2})t}}{-\frac{1}{2} + \frac{i\sqrt{3}}{2}} \\ e^{(-\frac{1}{2} + \frac{i\sqrt{3}}{2})t} \end{bmatrix} + c_3 \begin{bmatrix} \frac{e^{(-\frac{1}{2} - \frac{i\sqrt{3}}{2})t}}{(-\frac{1}{2} - \frac{i\sqrt{3}}{2})^2} \\ \frac{e^{(-\frac{1}{2} - \frac{i\sqrt{3}}{2})t}}{-\frac{1}{2} - \frac{i\sqrt{3}}{2}} \\ e^{(-\frac{1}{2} - \frac{i\sqrt{3}}{2})t} \end{bmatrix}$$

Which becomes

$$\begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix} = \begin{bmatrix} \frac{c_3(-i\sqrt{3}-1)e^{-\frac{(1+i\sqrt{3})t}{2}}}{2} + \frac{c_2(i\sqrt{3}-1)e^{\frac{(i\sqrt{3}-1)t}{2}}}{2} + c_1e^t \\ \frac{c_3(i\sqrt{3}-1)e^{-\frac{(1+i\sqrt{3})t}{2}}}{2} + \frac{c_2(-i\sqrt{3}-1)e^{\frac{(i\sqrt{3}-1)t}{2}}}{2} + c_1e^t \\ c_1e^t + c_2e^{\frac{(i\sqrt{3}-1)t}{2}} + c_3e^{-\frac{(1+i\sqrt{3})t}{2}} \end{bmatrix}$$

3.3.3 Maple step by step solution

Let's solve

$$[x'(t) = y(t), y'(t) = z(t), z'(t) = x(t)]$$

- Define vector

$$\vec{x}(t) = \begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix}$$

- Convert system into a vector equation

$$\vec{x}'(t) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \cdot \vec{x}(t) + \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

- System to solve

$$\vec{x}'(t) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \cdot \vec{x}(t)$$

- Define the coefficient matrix

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$$

- Rewrite the system as

$$\vec{x}'(t) = A \cdot \vec{x}(t)$$

- To solve the system, find the eigenvalues and eigenvectors of A
- Eigenpairs of A

$$\left[\left[\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right], \left[-\frac{1}{2} - \frac{\text{I}\sqrt{3}}{2}, \begin{bmatrix} \frac{1}{\left(-\frac{1}{2} - \frac{\text{I}\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{\text{I}\sqrt{3}}{2}} \\ 1 \end{bmatrix} \right] \right], \left[-\frac{1}{2} + \frac{\text{I}\sqrt{3}}{2}, \begin{bmatrix} \frac{1}{\left(-\frac{1}{2} + \frac{\text{I}\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} + \frac{\text{I}\sqrt{3}}{2}} \\ 1 \end{bmatrix} \right] \right]$$

- Consider eigenpair

$$\left[1, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right]$$

- Solution to homogeneous system from eigenpair

$$\vec{x}_1 = e^t \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

- Consider complex eigenpair, complex conjugate eigenvalue can be ignored

$$\left[-\frac{1}{2} - \frac{\text{I}\sqrt{3}}{2}, \begin{bmatrix} \frac{1}{\left(-\frac{1}{2} - \frac{\text{I}\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2} - \frac{\text{I}\sqrt{3}}{2}} \\ 1 \end{bmatrix} \right]$$

- Solution from eigenpair

$$e^{\left(-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)t} \cdot \begin{bmatrix} \frac{1}{\left(-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2}-\frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix}$$

- Use Euler identity to write solution in terms of sin and cos

$$e^{-\frac{t}{2}} \cdot \left(\cos\left(\frac{\sqrt{3}t}{2}\right) - i \sin\left(\frac{\sqrt{3}t}{2}\right) \right) \cdot \begin{bmatrix} \frac{1}{\left(-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)^2} \\ \frac{1}{-\frac{1}{2}-\frac{i\sqrt{3}}{2}} \\ 1 \end{bmatrix}$$

- Simplify expression

$$e^{-\frac{t}{2}} \cdot \begin{bmatrix} \frac{\cos\left(\frac{\sqrt{3}t}{2}\right) - i \sin\left(\frac{\sqrt{3}t}{2}\right)}{\left(-\frac{1}{2}-\frac{i\sqrt{3}}{2}\right)^2} \\ \frac{\cos\left(\frac{\sqrt{3}t}{2}\right) - i \sin\left(\frac{\sqrt{3}t}{2}\right)}{-\frac{1}{2}-\frac{i\sqrt{3}}{2}} \\ \cos\left(\frac{\sqrt{3}t}{2}\right) - i \sin\left(\frac{\sqrt{3}t}{2}\right) \end{bmatrix}$$

- Both real and imaginary parts are solutions to the homogeneous system

$$\vec{x}_2(t) = e^{-\frac{t}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}t}{2}\right)}{2} - \frac{\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{2} \\ -\frac{\cos\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{2} \\ \cos\left(\frac{\sqrt{3}t}{2}\right) \end{bmatrix}, \vec{x}_3(t) = e^{-\frac{t}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}t}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}t}{2}\right)}{2} \\ \frac{\cos\left(\frac{\sqrt{3}t}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}t}{2}\right)}{2} \\ -\sin\left(\frac{\sqrt{3}t}{2}\right) \end{bmatrix}$$

- General solution to the system of ODEs

$$\vec{x} = c_1 \vec{x}_1 + c_2 \vec{x}_2(t) + c_3 \vec{x}_3(t)$$

- Substitute solutions into the general solution

$$\vec{x} = c_1 e^t \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + c_2 e^{-\frac{t}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}t}{2}\right)}{2} - \frac{\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{2} \\ -\frac{\cos\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{\sqrt{3} \sin\left(\frac{\sqrt{3}t}{2}\right)}{2} \\ \cos\left(\frac{\sqrt{3}t}{2}\right) \end{bmatrix} + c_3 e^{-\frac{t}{2}} \cdot \begin{bmatrix} -\frac{\cos\left(\frac{\sqrt{3}t}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}t}{2}\right)}{2} \\ \frac{\cos\left(\frac{\sqrt{3}t}{2}\right)\sqrt{3}}{2} + \frac{\sin\left(\frac{\sqrt{3}t}{2}\right)}{2} \\ -\sin\left(\frac{\sqrt{3}t}{2}\right) \end{bmatrix}$$

- Substitute in vector of dependent variables

$$\begin{bmatrix} x(t) \\ y(t) \\ z(t) \end{bmatrix} = \begin{bmatrix} -\frac{e^{-\frac{t}{2}}(c_3\sqrt{3}+c_2)\cos\left(\frac{\sqrt{3}t}{2}\right)}{2} - \frac{e^{-\frac{t}{2}}(c_2\sqrt{3}-c_3)\sin\left(\frac{\sqrt{3}t}{2}\right)}{2} + c_1e^t \\ -\frac{e^{-\frac{t}{2}}(-c_3\sqrt{3}+c_2)\cos\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{e^{-\frac{t}{2}}(c_2\sqrt{3}+c_3)\sin\left(\frac{\sqrt{3}t}{2}\right)}{2} + c_1e^t \\ c_1e^t + c_2e^{-\frac{t}{2}}\cos\left(\frac{\sqrt{3}t}{2}\right) - c_3e^{-\frac{t}{2}}\sin\left(\frac{\sqrt{3}t}{2}\right) \end{bmatrix}$$

- Solution to the system of ODEs

$$\left\{ \begin{aligned} x(t) &= -\frac{e^{-\frac{t}{2}}(c_3\sqrt{3}+c_2)\cos\left(\frac{\sqrt{3}t}{2}\right)}{2} - \frac{e^{-\frac{t}{2}}(c_2\sqrt{3}-c_3)\sin\left(\frac{\sqrt{3}t}{2}\right)}{2} + c_1e^t, \\ y(t) &= -\frac{e^{-\frac{t}{2}}(-c_3\sqrt{3}+c_2)\cos\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{e^{-\frac{t}{2}}(c_2\sqrt{3}+c_3)\sin\left(\frac{\sqrt{3}t}{2}\right)}{2} + c_1e^t \end{aligned} \right.$$

✓ Solution by Maple

Time used: 0.047 (sec). Leaf size: 176

```
dsolve([diff(x(t),t)=y(t),diff(y(t),t)=z(t),diff(z(t),t)=x(t)],singsol=all)
```

$$\begin{aligned} x(t) &= c_1e^t + c_2e^{-\frac{t}{2}}\sin\left(\frac{\sqrt{3}t}{2}\right) + c_3e^{-\frac{t}{2}}\cos\left(\frac{\sqrt{3}t}{2}\right) \\ y(t) &= c_1e^t - \frac{c_2e^{-\frac{t}{2}}\sin\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{c_2e^{-\frac{t}{2}}\sqrt{3}\cos\left(\frac{\sqrt{3}t}{2}\right)}{2} \\ &\quad - \frac{c_3e^{-\frac{t}{2}}\cos\left(\frac{\sqrt{3}t}{2}\right)}{2} - \frac{c_3e^{-\frac{t}{2}}\sqrt{3}\sin\left(\frac{\sqrt{3}t}{2}\right)}{2} \\ z(t) &= c_1e^t - \frac{c_2e^{-\frac{t}{2}}\sin\left(\frac{\sqrt{3}t}{2}\right)}{2} - \frac{c_2e^{-\frac{t}{2}}\sqrt{3}\cos\left(\frac{\sqrt{3}t}{2}\right)}{2} \\ &\quad - \frac{c_3e^{-\frac{t}{2}}\cos\left(\frac{\sqrt{3}t}{2}\right)}{2} + \frac{c_3e^{-\frac{t}{2}}\sqrt{3}\sin\left(\frac{\sqrt{3}t}{2}\right)}{2} \end{aligned}$$

✓ Solution by Mathematica

Time used: 0.063 (sec). Leaf size: 234

```
DSolve[{x'[t]==y[t],y'[t]==z[t],z'[t]==x[t]},{x[t],y[t],z[t]},t,IncludeSingularSolutions ->
```

$$\begin{aligned}x(t) &\rightarrow \frac{1}{3}e^{-t/2} \left((c_1 + c_2 + c_3)e^{3t/2} + (2c_1 - c_2 - c_3) \cos\left(\frac{\sqrt{3}t}{2}\right) + \sqrt{3}(c_2 - c_3) \sin\left(\frac{\sqrt{3}t}{2}\right) \right) \\y(t) &\rightarrow \frac{1}{3}e^{-t/2} \left((c_1 + c_2 + c_3)e^{3t/2} - (c_1 - 2c_2 + c_3) \cos\left(\frac{\sqrt{3}t}{2}\right) - \sqrt{3}(c_1 - c_3) \sin\left(\frac{\sqrt{3}t}{2}\right) \right) \\z(t) &\rightarrow \frac{1}{3}e^{-t/2} \left((c_1 + c_2 + c_3)e^{3t/2} - (c_1 + c_2 - 2c_3) \cos\left(\frac{\sqrt{3}t}{2}\right) + \sqrt{3}(c_1 - c_2) \sin\left(\frac{\sqrt{3}t}{2}\right) \right)\end{aligned}$$

3.4 problem Problem 5

Internal problem ID [12211]

Internal file name [OUTPUT/10863_Thursday_September_21_2023_05_48_15_AM_75472716/index.tex]

Book: Differential equations and the calculus of variations by L. EISGOLTS. MIR PUBLISHERS, MOSCOW, Third printing 1977.

Section: Chapter 3, SYSTEMS OF DIFFERENTIAL EQUATIONS. Problems page 209

Problem number: Problem 5.

ODE order: 1.

ODE degree: 1.

The type(s) of ODE detected by this program : "**system of linear ODEs**" Unable to solve or complete t

Solve

$$\begin{aligned}x'(t) &= y(t) \\ y'(t) &= \frac{y(t)^2}{x(t)}\end{aligned}$$

Does not currently support non linear system of equations. This is the phase plot of the system.




Figure 127: Phase plot

✓ Solution by Maple

Time used: 0.062 (sec). Leaf size: 20

```
dsolve([diff(x(t),t)=y(t),diff(y(t),t)=y(t)^2/x(t)],singsol=all)
```

$$\begin{cases} x(t) = e^{c_1 t} c_2 \\ y(t) = \frac{d}{dt} x(t) \end{cases}$$

✓ Solution by Mathematica

Time used: 0.022 (sec). Leaf size: 28

```
DSolve[{x'[t]==y[t],y'[t]==y[t]^2/x[t]},{x[t],y[t]},t,IncludeSingularSolutions -> True]
```

$$\begin{aligned} y(t) &\rightarrow c_1 c_2 e^{c_1 t} \\ x(t) &\rightarrow c_2 e^{c_1 t} \end{aligned}$$